



Sebaceous Cyst Excision

General Anaesthetic Pre-Operative Instructions

Pre-operative Information for General Anaesthetic procedures

These pre-operative instructions are to help you prepare for your surgery. Please let me or my team know if you have any new medical problems or medications prior to surgery.

Smokers

You should cease smoking for a minimum of four weeks before the surgery, although discontinuing earlier is preferable. Nicotine diminishes blood oxygen levels and can markedly elevate the likelihood of post-operative healing complications. Since it's the nicotine and not the smoke itself that poses issues, this caution **extends to nicotine replacement therapies** such as chewing gum, e-cigarettes, and patches.

Sickle cell trait or sickle cell disease

If you have either condition, it is essential that we are made aware of this prior to your surgery to ensure that you are optimised and have blood transfusions and anticoagulation if necessary. This will be fully discussed at your initial consultation.

Dressing allergies

Please let me know if you have an allergy to tissue glue or any dressings so I can select an appropriate dressing for your incisions.

Medications

Certain medications should be stopped prior to surgery as they can cause blood clots or excessive bleeding. These include:

- **Aspirin / Clopidogrel:** To be stopped 2 weeks prior to surgery as they can cause excessive bleeding and haematoma formation.
- **Warfarin:** This should be stopped 4 to 5 days before surgery, depending on your INR and the indications for the Warfarin. We will start a blood thinning injection to protect you whilst the Warfarin has been discontinued.



Anne Dancey
PLASTIC SURGERY

Sebaceous Cyst Excision

General Anaesthetic Pre-Operative Instructions

Pre-operative assessment

Prior to your surgery, you will be given a pre-operative assessment with the anaesthetist. This can be either over the phone or in the clinic, if the anaesthetist wishes to see you face to face prior to surgery. Depending on your age and general health, routine blood tests, MRSA swabs and a heart tracing (ECG) may be taken.

You will also have an opportunity to ask any questions you may have or raise any concerns.

If you are unsure whether you wish to proceed with a procedure, then it is strongly advised not to proceed with surgery.

Admission to hospital

This will normally be on the day of your operation. If you are having a general anaesthetic, you will be asked not to eat anything 6 hours before surgery. Black tea/coffee or non-fizzy drinks (nothing containing milk) can be taken up to 4 hours before surgery. Chewing gum should also be avoided for 6 hours before your operation.

Sometimes delays do occur, however, you will be cared for whilst waiting under the direction of the anaesthetist and the nursing team.

I will come and see you with the anaesthetist before you go to the operating room. I will take pre-operative photographs and go over the operation again to remind you of any risks or complications.

If you have any questions or concerns, there will always be sufficient time made to answer them.

At any time if you are unsure about anything concerning surgery, then please speak to me.



Anne Dancey
PLASTIC SURGERY

Sebaceous Cyst Excision

General Anaesthetic Pre-Operative Instructions



I hope you find this information useful. If you have any questions or require a little more information, then please do not hesitate to contact me.

Anne Dancey

Dr Anne Dancey

Plastic and Reconstructive Surgeon FRCS(Plast), MBChB(Hons), MmedSci(Hons) and MCh(PASP)