



Anne Dancey  
PLASTIC SURGERY

## Accessory Breast Tissue Removal Post-Operative Instructions

### Sutures

Any small stab incision is closed with dissolvable sutures, which do not require removal and will dissolve over a period of approximately 7 to 14 days.

Patients with very large amounts of accessory tissue with stretched skin may require skin removal via an incision in the axilla. The additional sutures will also dissolve over a period of approximately 7 to 14 days.

### Dressings

I use waterproof dressings to cover the stab incisions. You can shower as soon as you like after surgery and do not require any dressing changes until we see you in clinic. However, you will be given some spare dressings in case they fall off.

### Medication

During your surgery I will use a long-acting local anaesthetic block. This reduces any pain that you may have for 72 hours after surgery, but you may need additional analgesia in the form of Tylenol and Ibuprofen.

I may consider prescribing Tramadol if you are in significant pain, but most patients do not need this. The medications should be given to you before you go home, so please check with the nurse if you do not receive any pain relief.

You do not need routine antibiotics after surgery as accessory breast tissue removal surgery has a low risk of infection.

The hospital will provide you with TED garments to wear during and after your surgery to reduce the risk of a DVT in your legs. It is recommended that you continue to wear them until you are fully mobile. You may also be given anticoagulants to take if you are at a higher risk of a DVT or PE, although most patients do not require this.

### Scar advice

Once the dressings are removed, I recommend that you massage the incisions using small circular motions at least twice a day for 5 minutes. You can use any moisturising cream of your choice and apply firm pressure until it blanches.



Anne Dancey  
PLASTIC SURGERY

## Accessory Breast Tissue Removal Post-Operative Instructions

The scars would also benefit from Silicone scar gel (ScarAway® or Kelo-Cote®) twice a day, which will soften and fade the scars as well as applying sunblock for 12 months to provide sun protection.

ScarAway® can be purchased from Healthcare Pharmacy at Governors Square in Grand Cayman.

### Follow up

Before you are discharged from Hospital you will be given a follow up appointment to see the nurse after one week post-operatively.

I will normally see you in the clinic 1-2 weeks following your procedure, depending on the type of surgery.

You should also be given contact numbers in the event of any problem. If you have any questions or concerns about your recovery, please do not hesitate to phone or email me and I will endeavour to get an answer to you the same day.

If it is more urgent then please contact the Clinic/Hospital. In the unlikely event of an emergency, you must visit your closest Emergency Department.



I hope you find this information useful. If you have any questions or require a little more information then please do not hesitate to contact me.

*Anne Dancey*

**Dr Anne Dancey**

Plastic and Reconstructive Surgeon FRCS(Plast), MBChB(Hons), MmedSci(Hons) and MCh(PASP)