

Grant Code: AP7075

Title: Weed seedbank control in rotational crops as a proactive herbicide resistance management strategy

Personnel: Albert Adjesiwor, University of Idaho

Address: Albert Adjesiwor; Kimberly Research & Extension Center; 3806 N 3600 E; Kimberly, ID 83341; Office: 208-423-6616; Email: aadjesiwor@uidaho.edu

Justification/Rationale: Herbicide-resistant weed populations are evolving rapidly and threatening the sustainability of crop production. In Idaho, there are 11 documented cases of herbicide resistance in seven common weed species. These resistant cases encompass at least seven herbicide sites of action, which are among herbicides commonly used in economically important crops such as wheat. Herbicide-resistant weeds can impact the economics of wheat production by reducing crop yield and contaminating the harvested grain. In a bid to manage the current threat of herbicide-resistant weeds, weed scientists continue to recommend crop rotations that include competitive crops like wheat and alfalfa.

It is known that crop rotations do not eliminate weeds but can be used to reduce the build-up of weed populations, especially herbicide-resistant weeds. Idaho has a wide variety of crops which offers more options for crop rotation sequences. Nevertheless, for crop rotations to be effective weed management tools, the selection of crops must take into account the type of weeds present. This proposed work seeks to answer the following questions: (1) what happens to the weed seeds in the soil when wheat is planted in rotation with alfalfa? (2) is it better to rotate wheat with alfalfa or other annual crops to manage troublesome weed seeds in the soil? (3) are there effective residual herbicides to manage weed seeds in the soil during the alfalfa crop before rotating back to wheat? Answering these questions would provide the necessary information to growers wishing to diversify their weed management program and reduce the selection or build-up of herbicide-resistant weeds.

Objectives: The goal of this research is to provide the foundational knowledge needed by stakeholders to adopt and integrate effective residual herbicide programs for weed control in wheat-alfalfa rotations. Specifically, we are proposing to:

1. Compare weed seedbank densities in wheat-alfalfa and wheat-corn/dry bean rotations
2. Evaluate residual herbicide programs for effective weed seedbank management within wheat-alfalfa rotations
3. Assess the economic impact of using herbicide mixtures and crop rotations for proactive herbicide resistance management

Methods/Plan of work: The field study was established under sprinkler irrigation at the University of Idaho Kimberly Research and Extension Center, Kimberly, ID. Before seeding weeds and crops in spring 2021, soil samples were collected across the research field to determine the weed seed bank composition. This experiment was laid out as a split-plot randomized complete block design with four replications.

Main plot (Crop rotation): Main plots are 45 ft wide by 30 ft long and consist of four crop rotations ranging in diversity and complexity (Figure 1).

Literature Review:

Crop rotation for weed management: Crop rotations have long been recognized as one of the most effective practices for weed management ([Pavlychenko and Harrington, 1934](#)). Diverse crop rotations provide some flexibility in the use of different weed control practices ([Goplen et al., 2017](#)). A four-year crop rotation study has shown that a diverse crop rotation that included small grains reduced herbicide-resistant kochia seed population in the soil by 34 to 68%, compared to continuous corn and less diverse rotations ([Mosqueda, 2019](#)). Aside from the competitiveness of small grains against most annual weeds, small grains are often harvested earlier in the summer before annual weed seed production ([Goplen et al., 2016](#)). This reduces the chances of weed escapes and enrichment of the seedbank. Including perennial forage crops like alfalfa in rotations also allow for the elimination of annual weed seed production through multiple crop harvest ([Goplen et al., 2017](#); [Meiss et al., 2010](#)). Thus, a crop rotation program that includes competitive small grain crops, perennial forage, and residual herbicide programs has the potential to deplete the seedbank of herbicide-resistant annual weeds.

Economics of proactive herbicide resistance management: The economic impact of herbicide resistance management has gained interest among scientists and farmers. A benchmark study has found that weed control costs were about 31% higher for best management practices recommended by academics compared to standard practices used by farmers ([Edwards et al., 2014](#)). However, crop yields tended to be greater in academic best management practices, thereby offsetting the additional cost incurred ([Edwards et al., 2014](#)). Using a bioeconomic model, [Livingston et al. \(2016\)](#) showed that although herbicide resistance management reduced first-year farm profits, net farm returns increased from the second year and subsequent 18 years. Thus, if implemented effectively, gains from herbicide resistance management would outweigh the short-term cost of resistance management.

References:

1. Edwards CB, Jordan DL, Owen MD, Dixon PM, Young BG, Wilson RG, Weller SC, Shaw DR (2014) Benchmark study on glyphosate-resistant crop systems in the United States. Economics of herbicide resistance management practices in a 5 year field-scale study. *Pest Management Science* 70:1924-1929
2. Goplen JJ, Sheaffer CC, Becker RL, Coulter JA, Breitenbach FR, Behnken LM, Johnson GA, Gunsolus JL (2016) Giant ragweed (*Ambrosia trifida*) seed production and retention in soybean and field margins. *Weed Technology* 30:246-253
3. Goplen JJ, Sheaffer CC, Becker RL, Coulter JA, Breitenbach FR, Behnken LM, Johnson GA, Gunsolus JL (2017) Seedbank depletion and emergence patterns of giant ragweed (*Ambrosia trifida*) in Minnesota cropping systems. *Weed Science* 65:52-60
4. Livingston M, Fernandez-Cornejo J, Frisvold GB (2016) Economic returns to herbicide resistance management in the short and long run: the role of neighbor effects. *Weed Science* 64:595-608
5. Meiss H, Médiène S, Waldhardt R, Caneill J, Munier-Jolain N (2010) Contrasting weed species composition in perennial alfalfas and six annual crops: implications for integrated weed management. *Agronomy for sustainable development* 30:657-666
6. Mosqueda EG (2019) Efficacy and economics of cultural and mechanical weed control practices for herbicide-resistant weed management. Ph.D. Dissertations. Laramie WY: University of Wyoming. 111p.
7. Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Science* 60:31-62
8. Pavlychenko T, Harrington J (1934) Competitive efficiency of weeds and cereal crops. *Canadian Journal of Research* 10:77-94

**FY2025
COMMODITY COMMISSION BUDGET
Principal Investigator: Albert Adjesiwor**

Allocated by <u>Idaho</u> Wheat Commission	during FY2023	\$	19,448
(Commission/Organization)			
Allocated by <u>Idaho</u> Wheat Commission	during FY2024	\$	19,626
(Commission/Organization)			

REQUESTED SUPPORT	<u>Awarded for FY2024</u>	<u>Requested for FY2025</u>
Budget Categories		
(10) Salary (staff, post-docs, et NOTE: Faculty salary/fringe NOT allowed)	\$ 11,205	\$ 11,000
(12) Temporary Help/IH	\$ -	\$ -
(11) Fringe Benefits	\$ 403	\$ 275
(20) Travel	\$ 800	\$ 800
(30) Other Expenses	\$ 1,554	\$ 1,200
(40) Capital Outlay >\$5k	\$ -	\$ -
(45) Capital Outlay <\$5k	\$ -	\$ -
(70) Graduate Student		
Tuition/Fees	\$ 5,664	\$ 6,354
TOTALS	\$ 19,626	\$ 19,629

TOTAL BUDGET REQUESTED FOR FY2025:	\$ 19,629
---	------------------

BREAKDOWN FOR MULTIPLE INDEXES:				
Budget Categories	(Insert Co-PI Name)	(Insert Co-PI Name)	(Insert Co-PI Name)	(Insert Co-PI Name)
(10) Salary (staff, post-docs, et	\$ -	\$ -	\$ -	\$ -
(12) Temporary Help	\$ -	\$ -	\$ -	\$ -
(11) Fringe Benefits	\$ -	\$ -	\$ -	\$ -
(20) Travel	\$ -	\$ -	\$ -	\$ -
(30) Other Expenses	\$ -	\$ -	\$ -	\$ -
(40) Capital Outlay >\$5k	\$ -	\$ -	\$ -	\$ -
(45) Capital Outlay <\$5k	\$ -	\$ -	\$ -	\$ -
(70) Graduate Student				
Tuition/Fees	\$ -	\$ -	\$ -	\$ -
TOTALS	\$ -	\$ -	\$ -	\$ -
Total Sub-budgets				\$ -

Annual Report

Grant Code: AP7075

Title: Weed seedbank control in rotational crops as a proactive herbicide resistance management strategy

Personnel: Albert Adjesiwor, University of Idaho

Address: Albert Adjesiwor; Kimberly Research & Extension Center; 3806 N 3600 E; Kimberly, ID 83341; Office: 208-423-6616; Email: aadjesiwor@uidaho.edu

Abstract:

Herbicide-resistant weed populations are evolving rapidly and threatening the sustainability of crop production in Idaho. A 4-year crop rotation study was initiated in 2021 at the University of Idaho Kimberly Research and Extension Center to evaluate weed control and seedbank dynamics in wheat-alfalfa vs wheat-annual crop (corn and dry bean) rotations. There were three herbicide treatments: untreated, postemergence (POST) only, and preemergence (PRE) + POST. It was observed that weed seedbank density was reduced from 8,737 to as low as 470 seeds per 10 square ft in some treatments. Weed seedbank density tended to be higher in the untreated checks and there was a trend of preemergence (PRE) + postemergence (POST) treatments slightly reducing weed seedbank density compared to POST only treatment. Including alfalfa in the crop rotation significantly reduced weed seedbank density, irrespective of the herbicide treatment. On the contrary, dry bean in the rotation significantly increased weed seedbank density. Weed density within the crops during the growing season was influenced by the type of crop as well as the herbicide treatment. Both POST only and PRE + POST treatments reduced weed density compared to the untreated and the PRE + POST treatments reduced weed density in each crop compared to the POST only treatment. Weed control treatments had no effect on alfalfa yield. However, herbicide application (POST only and PRE + POST) improved corn and dry bean yield. The combination of fewer weeds and greater crop yield in the PRE + POST treatments holds promise for reducing weed seedbank and potentially improving crop productivity and economics.

Background: Herbicide-resistant weed populations are evolving rapidly and threatening the sustainability of crop production. In a bid to manage the current threat of herbicide-resistant weeds, weed scientists continue to recommend crop rotations that include competitive crops like wheat and alfalfa. This research project seeks to answer the following questions: (1) what happens to the weed seeds in the soil when wheat is planted in rotation with alfalfa? (2) is it better to rotate wheat with alfalfa or other annual crops to manage troublesome weed seeds in the soil?

Objectives: The objectives of this study were to:

1. Compare weed seedbank densities in wheat-alfalfa and wheat-corn/dry bean rotations
2. Evaluate residual herbicide programs for effective weed seedbank management within wheat-alfalfa rotations
3. Assess the economic impact of using herbicide mixtures and crop rotations for proactive herbicide resistance management

Results/Accomplishments

2023 field trial and weed seedbank density:

The soil samples collected from fall 2021 were thawed in spring 2022 for exhaustive germination in the greenhouse to estimate weed seedbank density. Seed density was counted weekly throughout the summer until there was no weed emergence. It was observed that weed seedbank density ranged from 470 to 8,737 seeds per 10 square ft (Figure 1). Weed seedbank density tended to be higher in the untreated checks and there was a trend of preemergence (PRE) + postemergence (POST) treatments slightly reducing weed seedbank density (Figure 1). Including alfalfa in the crop rotation significantly reduced weed seedbank density, irrespective of the herbicide treatment. On the contrary, dry bean in the rotation significantly increased weed seedbank density (Figure 1).

Weed density within the crops during the growing season was influenced by the type of crop as well as the herbicide treatment (Figure 2). Both POST only and PRE + POST treatments reduced weed density compared to the untreated. In addition, PRE + POST treatments reduced weed density in each crop compared to the POST only treatment. This has implications for the number of weeds that will go to seed at the end of the growing season.

Weed control treatments had minimal effect on alfalfa yield (Figure 3). However, herbicide application (POST only and PRE + POST) improved dry bean and wheat yield. The combination of fewer weeds and greater crop yield in the PRE + POST treatments holds promise for reducing weed seedbank and potentially improving crop productivity and economics.

Nest Steps/Projections: This study will be continued in 2024 and results will be made available to the Idaho Wheat Commission and Idaho wheat growers. Results from this study will be presented at the 2024 Weed Tour at Kimberly and the 2023 and 2024 Western Society of Weed Science Conference.

Publication/ Outreach: This study was showcased at the 2023 Weed Tour at Kimberly.

1. Chandra L. Montgomery, **Adjesiwor**, A.T. 2023. Weed Seedbank Control in Rotational Crops for Proactive Herbicide Resistance Management. Proceedings of the WSWs Annual Meeting. Page 24. <https://wsweedscience.org/wp-content/uploads/WSWS-2023-Proceedings.pdf>




Figure 1. Effect of postemergence (POST) and preemergence (PRE) POST herbicide treatments on weed density in spring wheat rotations. Soils collected from fall 2022 at Kimberly, Idaho.




Figure 2 Effect of postemergence (POST) and preemergence (PRE) POST herbicide treatments on weed density in spring wheat rotations in 2023 at Kimberly, Idaho.




Figure 3 Effect of postemergence (POST) and preemergence (PRE) POST herbicide treatments on crop yield in spring wheat rotations in 2023 at Kimberly, Idaho