

E-PRESCRIBING GUIDE – DIAZEPAM / BACLOFEN SUPPOSITORIES

SCHEDULE IV CONTROLLED SUBSTANCE — E-PRESCRIBE ONLY (EPCS)

Diazepam prescriptions cannot be faxed, phoned, or handwritten. Paper order forms are not accepted — this guide replaces them.

How to E-Prescribe

1. **Search for our pharmacy in your EPCS-enabled EHR.** Search the pharmacy directory for “Bayview Pharmacy” — or by ZIP code 02886, Warwick, RI. Your platform must be EPCS-certified with identity proofing and two-factor authentication.
2. **Verify you have the right location.** Confirm the listing matches the details in the box below before selecting it.
3. **Pick a drug entry that validates as controlled.** Your EHR’s drug database does not contain compounded suppositories. Select a **diazepam** entry — “Diazepam (bulk) powder” where your system offers it, otherwise any diazepam product — so the EHR classifies the prescription as Schedule IV and EPCS validates it. Then describe the actual suppository in the body.
4. **Enter the formulation and transmit.** Lead with the Bayview code, then the full formulation, strength, quantity, directions, days’ supply, and refills as permitted for Schedule IV (up to 5 refills within 6 months, where state law allows). State the **route (vaginal or rectal)** and include an ICD-10 diagnosis code.

IMPORTANT Put the formulation in the **sig / prescription body**, not in “Notes to Pharmacy”. Some EHRs treat that field as internal and never transmit it — the prescription arrives with no formula and cannot be filled.

Pharmacy Name	Bayview Pharmacy
Address	3844 Post Road, Warwick, RI 02886
NCPDP ID	4106882
NPI	1750455143
Phone	(401) 284-4505
Fax	(401) 284-4506 (not valid for diazepam prescriptions)

PRESCRIBE WITH BAYVIEW CODES

Every validated formulation has a short Bayview code (like **DB-5-10**) — listed on the validated formulations sheet on page 2. Lead with the code in your compound / free-text entry and write the strength where the formulation is custom:

Bayview DB-5-10 — diazepam 5 mg / baclofen 10 mg vaginal suppository, #30 — insert 1 suppository vaginally at bedtime as needed for pelvic floor spasm, max 1 per day, 30-day supply. Refills: 1 (4-week trial, then reassess). ICD-10: N94.1

Route must be stated — vaginal or rectal. The label and the base will match what you write.

Refills: Schedule IV allows a maximum of 5 within 6 months. Consider a defined trial duration with a reassessment or taper plan rather than open-ended refills; vaginal diazepam produces measurable serum levels, so systemic benzodiazepine effects apply.

Baclofen-only suppositories (no diazepam) are **not controlled** and may be faxed on the separate **Baclofen Suppositories** order form. Ketamine-containing suppositories are Schedule III — select a ketamine entry instead and see the ketamine guide.

Questions? Call (401) 284-4505 and one of our pharmacists will walk you through it.

Validated Formulations – Diazepam / Baclofen Suppositories

Companion to the E-Prescribing Guide • Codes identify the form and strength — the prescriber may set a custom strength within the validated range • E-prescribe only (EPCS) • Schedule IV

DIAZEPAM / BACLOFEN — vaginal or rectal suppository (state route)		dispensed per suppository
DB-5-10	Diazepam 5 mg / baclofen 10 mg suppository	15 30
DB-10-10	Diazepam 10 mg / baclofen 10 mg suppository	15 30
DB-CUST	Diazepam ____ mg / baclofen ____ mg suppository — validated range diazepam 2–10 mg, baclofen 5–20 mg	15 30

DIAZEPAM — single agent, vaginal or rectal suppository		dispensed per suppository
DB-D5	Diazepam 5 mg suppository	15 30
DB-D10	Diazepam 10 mg suppository	15 30

DIAZEPAM / BACLOFEN / KETAMINE — Schedule III (select a ketamine entry for EPCS)		dispensed per suppository
DB-K	Diazepam 5 mg / baclofen 5 mg / ketamine ____ mg suppository — validated range ketamine 17.25–25 mg	30
Ketamine-containing versions are Schedule III and carry the FDA's October 10, 2023 compounded-ketamine warning. Not controlled: baclofen-only (DB-B10) and baclofen / lidocaine (DB-BL) — use the Baclofen Suppositories fax form.		

Prescribing checklist: (a) Bayview code or full formulation and strength; (b) route — vaginal or rectal; (c) quantity (15 or 30); (d) directions/SIG; (e) days' supply; (f) refills (max 5 within 6 months) with a defined trial length; (g) ICD-10 diagnosis code (e.g. N94.1, N94.2, R10.2). **Not listed?** Additional strengths may be available — call the pharmacy to confirm before prescribing.

EVIDENCE Randomized trials of vaginal diazepam for hypertonic pelvic floor dysfunction have not shown benefit over placebo; no society guideline endorses these suppositories as standard therapy. Bayview prepares them only on prescriber request and makes no efficacy claim.