

This document is one component of the full-length and abbreviated PPTA donor history questionnaire documents. The full-length and abbreviated PPTA donor history questionnaire documents must be used collectively.

| Medication List | | | | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|-----------------------------------------------------|----------------|---------------------------------------------------------------------------------------------------------------------|
| Please review this list carefully and tell us if you are taking or have taken any of these medications. Do NOT discontinue your prescribed medications, including PrEP or PEP, in order to donate plasma. | | | | |
| Medication | Which is also known as... | And is usually used for... | In the last... | You should not donate because the medication may... |
| Absorica Accutane Amnesteem Claravis Myorisan Sotret Zenatane | isotretinoin | Severe acne | 1 month | Harm patients who receive medications prepared from your plasma by causing birth defects or death in unborn babies. |
| Propecia | finasteride | Baldness | | |
| Proscar | | Enlarged prostate gland | | |
| Thalomid Revlimid | thalidomide lenalidomide | Multiple myeloma | | |
| Rinvoq | upadacitinib | Rheumatoid arthritis | | |
| Depakote Depakote ER Convulex Eplim Stavzor | valproate valproic acid sodium valproate | Epilepsy Bipolar disorder Migraine prevention | | |
| CellCept | mycophenolate mofetil | Preventing organ rejection | 6 weeks | |
| Any medication taken by mouth (oral) for HIV prevention (also known as PrEP or PEP), examples include Truvada, Descovy | | | 3 months | Delay detection of HIV infection by tests performed for plasma donations. |
| HepaGam B HyperHEP B S/D Nabi- HB | hepatitis B immune globulin | Exposure to hepatitis B | | Not prevent you from contracting hepatitis B infection following a possible or known exposure. |
| Avodart Jalyn | dutasteride | Enlarged prostate gland | 6 months | Harm patients who receive medications prepared from your plasma by causing birth defects or death in unborn babies. |
| Any experimental or unlicensed vaccine or medication | | | 12 months | Harm patients who receive medications prepared from your plasma. Medical evaluation needed. |
| Arava | leflunomide | Rheumatoid arthritis | 2 years | Harm patients who receive medications prepared from your plasma by causing birth defects or death in unborn babies. |
| Aubagio | teriflunomide | Relapsing multiple sclerosis | | |
| Erivedge | vismodegib | Basal cell skin cancer | | |
| Odomzo | sonidegib | | | |
| Apretude Yeztugo | cabotegravir lenacapavir (injectable) | HIV prevention (also known as PrEP or PEP) | | Delay detection of HIV infection by tests performed for plasma donations. |
| Soriatane | acitretin | Severe psoriasis | 3 years | Harm patients who receive medications prepared from your plasma by causing birth defects or death in unborn babies. |
| Tegison | etretinate | Severe psoriasis | Ever | |
| Any medication to treat HIV. May also be called antiretroviral therapy (ART) | | | | |
| If you are taking or have taken these medications, you may not be eligible to donate plasma, whole blood, or platelets. | | | | |