

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Active Pharmaceutical Ingredient	Autoclave*	Dry Heat**	General Information on Thermal Stability
Acetylcysteine ¹	No	No	Studies on thermal degradation of acetylcysteine have found significant degradation of stock solutions at 80 °C over 3 hours.
Alprostadiol ²	No	No	Alprostadil is significantly susceptible to thermal degradation
Amikacin Sulfate ^{3,4}	Yes	Unknown	Information exists to suggest amikacin aqueous solutions are acceptable for autoclave sterilization. With regards to dry heat sterilization, information was unavailable. One study noted a small amount of degradation with amikacin dry powder exposed to 70 °C over 48 hours, but no studies were available at higher temperatures consistent with dry heat sterilization
Amphotericin B ^{5,6}	No	No	Though some information suggests limited amphotericin B stability, and even potentially enhanced activity at temperatures up to 41 °C in some trials, amphotericin B is not known to be stable to temperatures consistent with autoclave or dry heat sterilization.
Anastrozole ^{30,31}	Yes	Unknown	One study on bulk drug powder and anastrozole tablets found minimal degradation after exposure to 105 °C for 6 hours, but higher temperatures were not evaluated. Another study of anastrozole and testosterone subcutaneous pellets stated the method of sterilization was autoclave at 121 °C for 25 minutes at 20-25psi
Arginine HCl ^{7,8}	Yes	Unknown	Though information on dry heat sterilization of arginine HCl was unavailable, some information exists to support its stability to autoclave. Some studies have evaluated the effect of arginine at 75 °C for the prevention of protein aggregation as well in some preparations.
Ascorbic Acid ⁹	No	No	Ascorbic acid has known thermal instability

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Atropine Sulfate ^{7,10}	Yes	No	The presence of a degradant (apoatropine sulphate) as been noted after exposure to 150 ° C, so while some information supports autoclave stability, dry heat sterilization is not recommended
Biotin ¹¹	Unknown	No	Limited information is available regarding biotin thermal stability, some sources suggest good stability in solution in excess of 100 ° C, making this API potentially a candidate for autoclave
Bupivacaine HCl ^{12,13}	Yes	Unknown	Bupivacaine HCl solutions are stable to autoclave, information on dry heat sterilization was not available, but some studies on heat degradation suggest that there may be elevated thermal stability
Calcium Chloride ⁷	Yes	Unknown	Handbook on Pharmaceutical Manufacturing Formulations contains autoclave sterilized formulations, suggesting stability to autoclave
Calcium Gluconate ^{7,15}	Yes	No	Calcium gluconate undergoes thermal degradation at temperatures greater than 150 ° C
Cholecalciferol	No	No	One study of vitamin D3 in oil at various temperatures found that samples treated at 100 ° C for 30 min showed no significant reduction, while samples treated at 150 and 180 ° C were significantly degraded
Cyanocobalamin ¹⁷⁻¹⁹	Yes	No	While the package insert for cyanocobalamin states that it may be sterilized via autoclave without significantly effecting potency, data on higher temperatures suggests that degradation may occur if dry heat sterilization is attempted
Cyclosporine ^{20,21}	Yes	Unknown	Some studies evaluation aqueous preparations of cyclosporine have noted stability to autoclave at 121 ° C for 15 minutes at 15 psi

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Cysteine ^{22,23,59}	Unknown	Unknown	Information is limited, one study on degradation of dry powder amino acids found cysteine degradation starting at 200 °C suggesting some potential thermal stability, other studies have corroborated similar temperatures. However, specific information regarding sterilization under steam heat was not found in the literature
Dexamethasone Sodium Phosphate ²⁴	No	No	The package insert for dexamethasone sodium phosphate recommends against autoclaving due to dexamethasone sodium phosphate thermal instability
Dexpanthenol ^{25,26}	Unknown	Unknown	Information regarding dexpanthenol thermal stability at temperatures needed for terminal sterilization was unavailable, one study found good thermal stability at 40 °C for 6 hours, and another found no degradation in solution at 80 °C over 5 hours, but higher temperatures were not evaluated
Edetate Disodium ²⁷	No	Unknown	Information suggests EDTA degradation at temperatures greater than 200 °C
Estradiol ^{61,62}	No	No	One study of estradiol found nearly 10% degradation of bulk drug hormone and hormone in an aqueous cream formulation when exposed to 90 °C for 24 hours, another found approximately 30% degradation after exposure to 180 °C for 30min
Estradiol Valerate	Unknown	Unknown	Insufficient information to determine thermal stability
Finasteride ⁵⁵	No	No	While finasteride does have some thermal stability, greater than 10% degradation has been noted with exposure to 60-70 °C for 90 minutes
Folic Acid ^{28,29}	Yes	No	While some data exists to support stability to autoclave, some data on dry heat suggests degradation at temperatures consistent with those for dry heat sterilization

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Glutathione ³²	No	No	Little information is available on glutathione stability in solution under heat, but some data from food studies suggests complete degradation at temperatures of 75 °C or above.
Hydroxocobalamin ³³	No	No	Thermal studies have noted degradation at temperatures below those required for autoclave sterilization
Lidocaine	Unknown	Yes	One study found lidocaine degradation to occur beginning at approximately 196 °C. Note that though lidocaine has a high temperature for degradation, its melting point is below the temperature typically used for dry heat sterilization. Studies on the base form of lidocaine regarding stability to steam heat were unavailable.
Lidocaine HCl ³⁴	Yes	Unknown	The package insert for lidocaine HCl Injection specifies that the product may be autoclaved at 121 °C for 15min at 15psi. Studies on lidocaine exposure to dry heat have shown stability at least to 120 °C for 72 hours
Lysine ³⁴	No	No	One study on lysine at water activity 0.3, 0.5, and 0.7 found significant degradation when exposed to temperatures in excess of 75 °C
Magnesium Chloride ^{7,36}	Yes	Yes	Thermal degradation studies on magnesium chloride note hydrolysis starting at 203 °C, higher than typical for dry heat sterilization
Magnesium Sulfate ^{37,38}	Yes	Yes	Decomposition studies on magnesium sulfate suggest temperatures much higher than those used for dry heat sterilization are needed to degrade magnesium sulfate
Methionine ³⁹	Unknown	Unknown	One study noted methionine stability to autoclave, another noted degradation between 190-240 °C

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Methylcobalamin ³³	No	No	Thermal degradation studies between 100-121 °C have noted significant methylcobalamin degradation
Minoxidil ⁵⁵	No	No	While minoxidil does have some thermal stability, some limited degradation has been noted with exposure to 60-70 °C for 90 minutes, suggesting thermal degradation does occur
Nandrolone Decanoate ⁶³	Unknown	Unknown	Though some degradation studies on other nandrolone salts, such as nandrolone propionate, demonstrated stability at 150 °C and degradation at higher temperatures of 200 °C, specific information on nandrolone decanoate in oil was unavailable
Niacinamide ^{7,41}	Unknown	No	Some B-complex formulations containing thiamine, riboflavin-r-phosphate, pyridoxine HCl, and niacinamide have procedures requiring autoclave sterilization at 121 °C for 20 minutes per the Handbook on Pharmaceutical Manufacturing Formulations, suggesting stability to autoclave. Niacinamide has a melting point beginning at 129 °C
Papaverine HCl ⁷	Unknown	No	Insufficient information available to determine if autoclave sterilization is appropriate. One formula from the Handbook on Injectable Drugs sterilized a papaverine containing formula at 121 °C for 30 minutes
Phentolamine Mesylate	Unknown	Unknown	Insufficient information to determine thermal stability
Progesterone ^{7,51,52}	Unknown	Possible	One study evaluating progesterone at 60 °C over one week didn't note degradation. Another study of progesterone in pharmaceutical waste noted that temperatures of 300 °C for 15 minutes were needed to degrade progesterone completely (temperatures between 200- 350 °C were evaluated).
Pyridoxine HCl ^{7,41}	Yes	No	Information suggests that pyridoxine HCl is stable at temperatures required for autoclave, tests at 121 °C for 15 minutes have not noted pyridoxine degradation

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Riboflavin-5-Phosphate ⁷	Unknown	No	Some B-complex formulations containing thiamine, riboflavin-r-phosphate, pyridoxine HCl, and niacinamide have procedures requiring autoclave sterilization at 121 °C for 20 minutes per the Handbook on Pharmaceutical Manufacturing Formulations, suggesting stability to autoclave.
Semaglutide ⁴²	Unknown	Unknown	One study exposed semaglutide solutions to 105 °C for 15 hours and noted less than 3% degradation
Sodium Bicarbonate ⁴⁴	No	No	Some studies note that exposure of sodium bicarbonate to heat and moisture can result in the formation of carbon dioxide, water, and sodium carbonate
Tacrolimus ^{53,54}	No	No	One study evaluating stability of tacrolimus aqueous solution noted minimal degradation after exposure to 60 °C for 24 hours suggesting some thermal stability, however, one study analyzing stability to autoclave of several aqueous tacrolimus preparations noted significant degradation.
Testosterone ^{30,46,47,48}	Yes	Unknown	<p>Testosterone: One study of anastrozole and testosterone subcutaneous pellets stated the method of sterilization was autoclave at 121 °C for 25 minutes at 20-25psi for the combination and 40 minutes, all other parameters being the same, for testosterone pellets alone. One study of testosterone exposed to 100 °C for 4 hours though did not note significant degradation</p> <p>Testosterone Cypionate: one study observed no degradation when testosterone was exposed to 80 °C for 48 hours</p> <p>Testosterone Undecanoate: one study on testosterone undecanoate capsules exposed to 105 °C over 48 hours found over 18% degradation.</p>
Tetracaine HCl ⁵⁸	Yes	Unknown	The tetracaine HCl injection package insert states that the vial may be autoclaved at 121 °C at 15psi for 15 minutes

Active Ingredient Summary Table

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

Thiamine HCl ^{7,45}	Unknown	No	Some B-complex formulations containing thiamine, riboflavin-r-phosphate, pyridoxine HCl, and niacinamide have procedures requiring autoclave sterilization at 121 °C for 20 minutes per the Handbook on Pharmaceutical Manufacturing Formulations, suggesting stability to autoclave. Information suggest that thiamine heat stability is temperature related, with solutions at pH 3 being much more stable than those at pH 6 in one study.
Tirzepatide ⁶⁰	Unknown	Unknown	One study noted just over 3% degradation with dry heat exposure to 70 °C over 24 hours
Tocopherol Acetate (Vitamin E Acetate) ⁴⁹	Unknown	Unknown	One study of vitamin E in a cream found thermal stability up to 250 °C at which point there was a significant degradation curve
Vancomycin HCl ⁵⁰	No	No	One study noted vancomycin HCl degradation after exposure to 80 °C for 300 minutes.
Zinc ^{56,57}	Yes	Yes	<p>Zinc Sulfate - the Zinc Sulfate Compounded Injection USP Monograph notes autoclave as the method for sterilization. One study notes zinc sulfate stability up to 680 °C</p> <p>Zinc Chloride - Information not available on autoclave, but one study found zinc chloride to be stable up to temperatures of 320C</p> <p>Zinc Acetate - information not available</p>

*Autoclave refers to steam heat conditions of 121 °C or greater under pressure, typically about 15psi

**Dry heat refers to non-aqueous heat exposure typically above 160 °C

References:

1. Primas N, Lano G, Brun D, et al. Stability Study of Parenteral N-Acetylcysteine, and Chemical Inhibition of Its Dimerization. Pharmaceuticals (Basel). 2023;16(1):72. Published 2023 Jan 3. doi:10.3390/ph16010072

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

2. Younger EW, Szabo RM. The stability of prostaglandin E1 in dilute physiological solutions at 37 degrees C. Prostaglandins. 1986;31(5):923-927. doi:10.1016/0090-6980(86)90024-9
3. Allen L, Amikacin 250mg/ml Injection. Amikacin 250 mg/mL Injection (uspharmacist.com). Published November 16th, 2011, Accessed 11/13/2023.
4. Francis D. Pien, Peter W. L. Ho, Antimicrobial spectrum, pharmacology, adverse effects, and therapeutic use of amikacin sulfate, American Journal of Hospital Pharmacy, Volume 38, Issue 7, 1 July 1981, Pages 981–989, <https://doi.org/10.1093/ajhp/38.7.981>
5. Kalyani L, Venkata C, Rao N. Stability indicating RP-HPLC method development and validation of cefepime and amikacin in pure and pharmaceutical dosage forms. Braz. J. Pharm. Sci. 2018; 54(3):e17258. <https://doi.org/10.1590/s2175-97902018000317258>
6. Hamilton-Miller, Jeremy. (1973). The effect of pH and temperature on the stability and bioactivity of nystatin and amphotericin B. The Journal of pharmacy and pharmacology. 25. 401-7. 10.1111/j.2042-7158.1973.tb10035.x.
7. Niazi S. Handbook of Pharmaceutical Manufacturing Formulations: Sterile Products. Informa Healthcare. Deerfield, IL. 2009.
8. Steven Ren, Effects of arginine in therapeutic protein formulations: a decade review and perspectives, Antibody Therapeutics, 2023;, tbad022, <https://doi.org/10.1093/abt/tbad022>
9. Yuan JP, Chen F. Degradation of ascorbic acid in aqueous solution. ACS Publications. 1998: 46(12):5078-5082.
10. Rosenblum E, Taylor W. Melting points with decomposition and the heat stability of atropine sulphate. Journal of Pharmacy and Pharmacology. 1954; 6(1):. 656-659.
11. Fiume MZ; Cosmetic Ingredient Review Expert Panel. Final report on the safety assessment of biotin. Int J Toxicol. 2001;20 Suppl 4:1-12.
12. Bupivacaine HCl [package insert]. Baxter Corporation. Mississauga, Ontario. 2019.
13. Mcvay G. Handbook on Injectable Drugs. 19th Ed. American Society of Health Systems Pharmacists. Pg 177.
14. Sykuta-Zajac, A., Łodyga-Chruścińska, E., Patecz, B. et al. Thermal and X-ray analysis of racemic bupivacaine hydrochloride. J Therm Anal Calorim 105, 1031–1036 (2011). <https://doi.org/10.1007/s10973-011-1425-9>
15. Labuschagne, Johan & Focke, Walter & Verryn, Sabine. (2001). Identification of the Thermal Decomposition Products of Calcium Gluconate Monohydrate by X-Ray Diffraction. 10.13140/RG.2.2.11630.23361.
16. Zareie M, Abbasi A, Faghhi S. Thermal Stability and Kinetic Study on Thermal Degradation of Vitamin D3 in Fortified Canola Oil. J Food Sci. 2019;84(9):2475-2481.doi:10.1111/1750-3841.14764
17. Cyanocobalamin [package insert] Hikma Pharmaceuticals. Berkly Heights, NJ. 2022.
18. Jagerstad M, Arkbage K. Cobalamins: Properties and Determination. Encyclopedia of Food Sciences and Nutrition. 2nd Ed. 2002. Pg 1419-1427.
19. Lee W, Lee YB, Huh M, Choi J. Determination of the chemical stability of cyanocobalamin in medical food by a validated immunoaffinity column-linked HPLC method. Journal of Food Quality. 2022. <https://doi.org/10.1155/2022/1619936>
20. Başaran E, Yenilmez E, Berkman MS, Büyükköroğlu G, Yazan Y. Chitosan nanoparticles for ocular delivery of cyclosporine A. J Microencapsul. 2014;31(1):49-57. doi:10.3109/02652048.2013.805839
21. Lallemand F, Schmitt M, Bourges JL, Gurny R, Benita S, Garrigue JS. Cyclosporine A delivery to the eye: A comprehensive review of academic and industrial efforts. Eur J Pharm Biopharm. 2017;117:14-28. doi:10.1016/j.ejpb.2017.03.006
22. Weiss, I.M., Muth, C., Drumm, R. et al. Thermal decomposition of the amino acids glycine, cysteine, aspartic acid, asparagine, glutamic acid, glutamine, arginine and histidine. BMC Biophys 11, 2 (2018). <https://doi.org/10.1186/s13628-018-0042-4>

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

23. Pokorný V, Štefja V, Havlín J, Fulem M, Růžička K. Heat Capacities of L-Cysteine, L-Serine, L-Threonine, L-Lysine, and L-Methionine. *Molecules*. 2023; 28(1):451. <https://doi.org/10.3390/molecules28010451>
24. Dexamethasone Sodium Phosphate [package insert]. Pfizer Laboratories. New York, NY. 2011.
25. Mahboubi A, Gholamreza Alviri M, Afshar M, Farhangi M. Development and Validation of A Fast, Simple And Specific Stability Indicating RP-HPLC Method for Determination of Dexpanthenol in Eye Gel Formulation. *Iran J Pharm Res*. 2019 Spring;18(2):670-676. doi: 10.22037/ijpr.2019.1100681. PMID: 31531050; PMCID: PMC6706706.
26. A Doganay, B Koksel, S O Gundogdu, Y Capan, Simultaneous Determination of Dexpanthenol, Lidocaine Hydrochloride, Mepyramine Maleate and their Related Substances by a RP-HPLC Method in Topical Dosage Forms, *Journal of Chromatographic Science*, Volume 56, Issue 10, November 2018, Pages 903– 911, <https://doi.org/10.1093/chromsci/bmy067>
27. Boles JS, Ritchie K, Crerar DA. Reducing the potential for migration of radioactive waste: aqueous thermal degradation of the chelating agent disodium edta. *Nuclear Chemical Waste Management*. 1987; 7(2): 89-93. .
28. Daniel E, Kline OL. Factors effecting folic acid determination. Food and Drug Administration, Federal Security Agency. 1947. 739-746.
29. Neves D, Lobato K, Angelica R et al. Thermal and in vitro digestion stability of folic acid in bread. *Journal of Food Composition and Analysis*. 2019; 84: 103311.
30. Glaser R, York A. Subcutaneous testosterone anastrozole therapy in men: rationale, dosing, and levels on therapy. *IJPC*. 2019; 23(4). 325-339.
31. Omini Y, Desireddy R, Kumar A. development of new validation RP-HPLC method estimation of anastrozole in bulk and tablet dosage forms. 2020; 8(3): 2321-3647.
32. Ohta Y, Chaleckis R, Katsumata M, Yokota K, Ayabe S, Oshima T, et al. Prolonged 20 h heating of chicken meat in water at 75°C softens the meat, increases the levels of endogenous and degraded metabolites in the stock, but depletes reduced glutathione and glutamine. *ChemRxiv*. Cambridge: Cambridge Open Engage;2022;
33. Gouldstein S, Duca A. Contributions to the analytical chemistry of vitamin B12. The thermal stability of cyanocobalamin, hydroxocobalamin and cobinamide in the solid state. *Thermochimica Acta*. 1982; 59(2): 211-220.
34. Lidocaine Hydrochloride [package insert]. Hospira Inc. Lake Forest, IL. 2021.
35. Desrosiers T, Savoie L. Extent of damage to amino acid availability of whey protein heated with sugar. *J Dairy Res*. 1991;58(4):431-441. doi:10.1017/s002202990003003x
36. Huang Q, Lu G, Wang J, Yu J. Thermal decomposition mechanisms of MgCl₂·6H₂O and MgCl·H₂O. *Journal of Analytical and Applied Pyrolysis*. 2011; 91(1): 159-164.
37. Magnesium Sulfate Injection, 500mg/ml in 2mL and 10mL ampoule. Manual for Procurement and Supply of Quality-Assured MNCH Commodities. MNCH Commodities Procurement-Magnesium Sulfate.pdf (ghsupplychain.org). Accessed 11/21/23.
38. Scheidema M, Taskinen P. Decomposition thermodynamics of magnesium sulfate. *American Chemical Society*. 2011; 50(16): 9550-9556.
39. Taira H. Part X. The influence of added sugars on the heat destruction of basic and sulphur containing amino acids in soybean products. Food Research Institute, Ministry of Agriculture and Forestry. 1966: 847-855.
40. Yablokov VA, Vasina YA, Zelyaev IA, Mitrofanova SV. Kinetics of thermal decomposition of sulfur containing amino acids. *Russian Journal of General Chemistry*. 2009; 79: 1141-1145.

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

41. Moreschi E, Matos J, Almeida-Muradian L. Thermal análisis of vitamin PP niacin and niacinamide. *Journal of Thermal Analysis and Calorimetry*. 2009; 98(1): 161-164.
42. Manasa M, Aanandhi V. Stability indicating method development and validation of semaglutide by RP-HPLC in pharmaceutical substance and pharmaceutical product. *Research Journal of Pharmacy and Technology*. 2021; 14(3). DOI: 10.5958/0974-360X.2021.00247.X
43. Hochberg M, Melnick D, Oser B. On the stability of pyridoxine. *Food Research Laboratories Inc*. 1944. 129-136.
44. Voelker, A.L., Taylor, L.S. & Mauer, L.J. Effect of pH and concentration on the chemical stability and reaction kinetics of thiamine mononitrate and thiamine chloride hydrochloride in solution. *BMC Chemistry* 15, 47 (2021). <https://doi.org/10.1186/s13065-021-00773-y>
45. SHOHL AT. PREPARATION OF SODIUM BICARBONATE FOR INTRAPERITONEAL USE. *Am J Dis Child*. 1929;38(5):927-928. doi:10.1001/archpedi.1929.01930110026003
46. Didarul M, Hasan M, Mohluddin T, Hassan M, Latif A, Haque P. Analytical method validation of testosterone undecanoate soft gelatin capsule by RP-HPLC method. *Journal of Developing Drugs*. 2018; 7(1): DOI: 10.4172/2329-6631.1000184
47. Bharti A, Kumbhare R, Jeyaseelan C. HPLC Method development and validation for the quantification of related impurities in testosterone cypionate active pharmaceutical ingredient. *Indian Journal of Pharmaceutical Education and Research*. 2022; 56(1): 240-246
48. Basha D, Reddy G, Rani T, Susithra E, Chekkara R, Vijay R, Shravan R. Simultaneous determination of impurities and degradation products by rapid RP-UPLC-MS method. *Scholars Research Library*. 2016; 8(5):442-450.
49. Mandelli de Almeida M, Ribeiro de Castro Lima C, Quenca-Guillen J et al. Stability evaluation of tocopheryl acetate and ascorbyl tetraisoalmitate in isolation and incorporated in cosmetic formulations using thermal analysis. *Brazilian Journal of Pharmaceutical Sciences*. 2010; 46(1): 129-134.
50. Masse M, Genay S, Martin Mena A, et al. Evaluation of the stability of vancomycin solutions at concentrations used in clinical services. *Eur J Hosp Pharm*. 2020;27(e1):e87-e92. doi:10.1136/ejhpharm-2019-002076
51. Konduru N, Kethe V, Gundla R, Katari N, Mallavarapu R. Determination of progesterone (steroid drug) in the semi-solid dosage form (vaginal gel) using a stability indicating method by RP-HPLC/PDA detector. *Biomedical Chromatography*. 2021; Doi: 10.1002/bmx.5246.
52. Javid, Faisal & Ang, Teck Nam & Hanning, Sara & Svirskis, Darren & Burrell, Rob & Taylor, Matt & Wright, L. & Baroutian, Saeid. (2021). Subcritical Hydrothermal Deconstruction of Two Hormones (Adrenaline and Progesterone) in Pharmaceutical Waste. *The Journal of Supercritical Fluids*. 179. 105388. 10.1016/j.supflu.2021.105388.
53. Peterka, T., Lušin, T., Bergles, J., Ham, Z., Grahek, R. & Urleb, U. (2019). Forced degradation of tacrolimus and the development of a UHPLC method for impurities determination. *Acta Pharmaceutica*, 69(3) 363-380. <https://doi.org/10.2478/acph-2019-0025>
54. Prajapati M, Eriksson F, Loftsson T. Stability characterization, kinetics and mechanism of tacrolimus degradation in cyclodextrin solutions. *International Journal of Pharmaceutics*. 2020; 586(30); 119579
55. Patel J, Tandel J, Chhalotiya U, Patel K. Stability indicating planar chromatographic method for estimation of minoxidil and finasteride combination used in the treatment of hair loss. *Journal of medicinal and chemical sciences*. 2021; 17-28.
56. Zinc Sulfate Compounded Injection. *Pharmacopeial Volume No 45(3)*. Current DocID: GUID-92CE31FB-3538-4A65-B205-87F81409BB71_2_en-US
57. Jones F, Tran H, Lindberg D, Zhao L. Thermal stability of zinc compounds. *Energy & Fuels*. 2013; 27(10): 5663.
58. Tetracaine HCl [package insert] Burbank, CA. Cameron Pharmaceuticals LLC. 2021.

Thermal Stability

Note that stability to elevated temperature can be contingent on other factors including other excipients, solvents, preservatives found in the preparation or chemical characteristics such as pH of the solution. The below is meant to provide information on potential temperature tolerances of various APIs. Stability testing is recommended on specific preparations or combinations to verify the suitability of the sterilization method.

59. Senapati, Samir Shyambabu, "In-Situ Gel of Cyclosporine A – Loaded Solid Lipid Nanoparticles for Topical Occular Delivery for Dry Eye Disease" (2019). Electronic Theses and Dissertations. 1688.
60. Godela R. Qbd approach for analysis of tirzepatide in its bulk and marketed formulation by stability indicating RP-HPLC. International Journal of Pharmaceutical Quality Assurance. 2023; 14(2): 404-409.
61. Gautam P. Method development and validation of stability indicating RP-HPLC method for the determination of female hormones in cream concentrates. Pharmaceutical Analytical Chemistry. 2017; 3(1).
62. Saha, N., McGaughy, K., Held, M.A. et al. Hydrothermal degradation of β -estradiol and oxytetracycline at selective reaction severities. SN Appl. Sci. 2, 1598 (2020). <https://doi.org/10.1007/s42452-020-03436-0>
63. Hadeif Y, Kaloustian J, Nicolay A, Portugal H. Thermal stability evaluation of doping compounds before GC-MS analysis by DSC. Journal of Thermal Analysis and Calorimetry. 2009; 93(2): 553-560.