

English

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: (2) Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Hindi

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Gujarati

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Marathi

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Telugu

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Kannada

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Bengali

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



Punjabi

Phyto Blend

Gazette Notification Number and Date: S.O. 3922(E).—
12-9-2024

Title of Biostimulant: Sargassum tenerrimum 2%(Granular)

Composition of Biostimulant: Seaweed Extract of Sargassum tenerrimum 2.0% w/w, Alginic Acid per cent by weight minimum 0.2% w/w, Phytase 1.0% w/w, Bulk density (g/cc) 0.95-1.1% w/w, pH (1:2 aqueous solution) 6.0-8.5 w/w and Granule Q.S.

Crops: Paddy and Tomato

Dosage: 5 kg/acre, Soil Application



English

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Hindi

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Gujarati

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Marathi

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Telugu

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Kannada

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Bengali

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.



Punjabi

Humizen Ultra

Gazette Notification Number and Date: S.O. 3922(E).—12-9-2024

Title of Biostimulant: Potassium Humate 49% (Powder)

Composition of Biostimulant: Humic Acid per cent. by weight minimum 21% w/w and pH (1 % aq solution) 6.0 – 8.0 w/w

Crops: Paddy and Tomato

Dosage: 400 gm/acre of powder

Single soil application by broadcast method.
It can be also mixed and broadcasted with organic and inorganic fertilisers during application with suitable soil moisture conditions . **Do not apply with bactericides.**
Give sufficient intervals between the applications.

