



Total Hip Replacement

Patient Information and Advice Booklet

Contents

Ulster Independent Clinic	3
The Patient Journey.....	4
Attendance at pre-operative assessment clinic	5
Preparation at home	6
Admission for your operation	9
Your hip replacement	10
Discharge from hospital.....	14
Follow-up review at outpatient clinic	16
Risks and potential problems following hip replacement surgery	17

WELCOME

TO THE ULSTER INDEPENDENT CLINIC

If you are reading this information booklet then you are waiting for, or are considering having, a total hip replacement. This booklet will help you to understand what to expect from the time you are placed on your consultant's operation list through to your rehabilitation and follow-up reviews. Having an operation can be a worrying experience, but learning more about what to expect often helps to overcome the fear of the unknown.

It is extremely important that you and the person who will be helping to look after you following your discharge read this booklet.

THE PATIENT JOURNEY

Consultation with surgeon & added to theatre operation list - provisional date for surgery given

Phone call from UIC with date for pre-operative assessment

Written confirmation of date & time of admission and special instructions

Attendance at Pre-operative Assessment Clinic Invitation to Pre-op physiotherapy hip class

Preparation at Home

Admission for Operation

Hip Replacement

Discharge from Hospital

Follow-up Review at Outpatient Clinic

Attendance at Pre-operative Assessment Clinic

Before coming in for your operation, you will be seen at a pre-operative assessment clinic to decide on your fitness for surgery. Blood tests and other investigations such as an ECG (heart tracing) may be required in order to make the surgery as safe as possible. There will also be standard infectious diseases screening tests if you have attended another hospital within the past year.

It is important to bring a repeat prescription list or medications in original packaging to both pre-op assessment and on admission. If you are taking any medication(s) which need to be stopped (eg warfarin, clopidogrel or blood pressure medication) before the operation, then your anaesthetist will give you this instruction in writing during your appointment.

Please bring details of your medical/surgical history to this appointment.

During admission you will also be asked if you are happy to have your details included on the **National Joint Registry**. The National Joint Registry collects information on total hip and knee replacement operations from hospitals in England, Wales and Northern Ireland.

If you would like additional information on the National Joint Registry, please visit the website on www.njrcentre.org.uk.

Preparation at Home

There are lots of things that you and the person helping you can do to prepare in the days leading up to your operation. A useful check-list is provided below:

1. Read ALL of the information provided to you (including this booklet).
2. Adhere to any instructions you have been given about STOPPING CERTAIN MEDICATIONS.
3. Adhere to any instructions about FASTING before your admission.
4. Arrange with a family member/friend to support you at home for the first two weeks after discharge. Further guidance is available by contacting our Hospital Liaison Sister and in discussion at pre-operative assessment clinic.
5. Prepare your house for your recovery (food shopping, freezing meals etc to use upon your discharge).
6. Organise your own transport to and from the hospital.

Prepare what you need for your hospital stay:

7. Your regular medication, in original packaging, and repeat prescription list or a print out from your GP.
8. Personal toiletries, pyjamas/nightdress, dressing gown, slippers (not new and neat fit), loose comfortable clothes for discharge.

If you have been told by the anaesthetist or at pre-operative assessment about **stopping certain tablets**, make sure you do this, otherwise **your surgery will not go ahead**. Likewise, should your GP **change any of your medications** between the time of your pre-operative assessment clinic and your surgery, it is important that **either you or your doctor let us know** (contact your consultant's secretary).

Similarly, **if you have become unwell or develop an infection** of any description (e.g. a chest infection) **or need antibiotics for any reason before your admission please contact us as your surgery may need to be postponed until the infection has cleared**.

Please bring your medication and an up to date medication list into hospital with you. If you usually take warfarin, please bring in any records or booklets you have which give details of usual doses. These will be returned to you on your discharge from hospital.

You should also inform us if you have any leg ulcers, wounds, open sores or discharging spots as this may prevent your operation from proceeding.

Fasting

You should fast as indicated on your admission letter. This means no food, no sweets, no chewing gum, no fruit juices should be eaten or drunk. **Only** water may be taken **up to 2 hours before surgery**.

If you eat or drink after these times your surgery will be cancelled. It is very important that you tell us if you have eaten or drunk anything after the cut-off time.

Adopting a healthy lifestyle

It is important that you make good lifestyle choices both before and after your operation. If you smoke cigarettes, cigars or a pipe, you should stop smoking for as long as possible before your operation and for 6 weeks after your operation to reduce the risk of complications, particularly infections.

Body Mass Index (BMI) is the most commonly used method to classify adult weight. It is defined as weight in kilograms divided by the square of height in metres (kg/m²). A BMI of greater than 25 is overweight. A BMI of greater than 30 is classed as obese. A high BMI can be associated with increased risks around the time of the operation. If you are overweight, try to develop healthier eating habits. Maintaining a healthy weight will have long term health benefits to you.

Keep as active as possible before your operation and try to achieve the recommended minimum of 20 minutes of exercise on a daily basis.

For more information on these subjects and helpful hints and tips please visit the websites below:

- nidirect.gov.uk/information-and-services/health-and-wellbeing
- nichs.org.uk/your-health/how-to-keep-healthy/
- belfasttrust.hscni.net/HealthyLiving.htm

Admission for your Operation

You will either be admitted on the morning of your operation or on the afternoon of the day before. If you have not previously signed a consent for surgery form, then your surgeon will discuss the operation beforehand with you and you will both sign the consent form together. If you have already signed your consent form before being admitted, then your surgeon will check it with you before you go for your operation.

Following this, a mark will then be drawn on your leg. This is an arrow that points to the hip that you are going to have replaced. This is an important part of the process that we use to make sure that you have the correct operation on the correct side.

You will be admitted by the nurse on the ward. You will also be seen by your consultant surgeon and the anaesthetist who will talk you through the anaesthetic process and the management of pain following your operation. The anaesthetist may prescribe a 'pre-med' (pre-medication) prior to your operation. This is the name for drugs that can be given before an anaesthetic. These may include drugs to prevent sickness, to reduce acid in the stomach, to help you relax or to reduce pain.

Your Hip Replacement

You will be in theatre and the recovery ward for approximately 6 hours in total:

- Half an hour is spent in the anaesthetic room
- One to one and a half hours is for the operation itself
- Five hours is spent in the recovery ward after your surgery

Your anaesthetic

In most cases a spinal anaesthetic is used. This makes you numb from the waist down so that you feel nothing during your operation and for a few hours afterwards. The anaesthetist will also give you sedation to make you relaxed and sleepy during the operation. You will usually hear nothing during the operation but if you do, you will be relaxed and this should not upset you. Also, a screen is used so that you will not see any of the operation taking place.

In certain circumstances, it may be necessary to do the operation under general anaesthetic. This is rare and will be decided by the anaesthetist and discussed with you prior to surgery.

On the rare occasion a tube will be inserted into your bladder (urinary catheter). This is necessary to allow medical staff to monitor your kidney function. This is normally removed the day after your surgery or when you start to walk.

Type of hip replacement

A hip replacement is typically composed of a stem which is inserted into the femur, a cup which is implanted into the pelvis (acetabulum) and the head. The head and the cup are known as the bearing surfaces, which take the load of your joint.

The exact type of hip replacement used may or may not require cement, and the choice is based upon surgeon preference and patient factors (see examples below). All of the implants we use are tried and trusted with good medium to long-term results as proven by the National Joint Registry (NJR – see page 5).

Occasionally different implants have to be used because of problems arising during surgery.

After your surgery

After your surgery you will be transferred to the recovery ward in theatre for a period of monitoring, as detailed below, before you return to the main ward.

- Blood pressure monitored every 15-30 minutes
- Oxygen may be administered with a facemask
- IV fluids will be given
- Flotrons (diagram to the right) will be attached to your lower legs to promote circulation
- Pressure dressing will be on your hip



If you have had a spinal anaesthetic your legs will be checked for movement and sensation which usually return to normal within a few hours.

In the unlikely event you have a urinary catheter inserted, this will be removed as soon as possible after your surgery.

The day after your operation you will have routine blood tests performed and a post-operative X-ray of your new hip replacement will be taken.

It is **extremely important that you begin to mobilise as soon as possible** and you will be seen by the physiotherapy team who will help you start walking with crutches or a frame and give you exercises to practise.

Post-operative pain

After surgery, you may feel discomfort or pain as the anaesthetic wears off. You will be offered pain relief if and when you need it. Our nursing staff will evaluate your pain levels throughout your admission and can help you understand how best to manage your pain before you go home. Our pharmacist or a nurse will also meet with you and discuss your medications for discharge.

Discharge from Hospital

Normally you will be able to go home as early as 1-2 days after your surgery. The best place for you to recover is at home with support from your family or the person helping you.

It is vital that you have a post-operative X-ray of your new hip replacement while in hospital.

Pain relief at discharge

You will be given a short supply of pain relief and laxatives to take home with you which should last 4-5 days. You will receive a discharge letter for your GP which you should leave into the surgery as soon as possible after discharge. Your GP may decide to prescribe further pain relief.

If you were taking anti-inflammatory pain relief (such as naproxen or ibuprofen) you should discuss with the surgeon or a doctor in the hospital whether to restart it.

Removal of skin clips

Some surgeons use skin clips to close your hip replacement wound. If these items have been used in your care, arrangements will be made for their removal. Typically, this procedure occurs approximately ten days following your operation, either by a district nurse, at your GP surgery, or alternatively, at the UIC Outpatients Treatment Room.

Prevention of clot formation

You may need to wear compression stockings during and after the operation. This will help to prevent blood clots forming in the veins in your legs (deep vein thrombosis). You may need to have an injection of an anti-clotting medicine (or tablets) as well as or instead of wearing compression stockings.

If a daily injection is required you will be taught how to inject this yourself. The Clinic Pharmacy will supply you with the necessary injections and guidance on discharge.

It is also important to **mobilise regularly** with short walks throughout the day and adapt a healthy diet and fluid intake. Rotating your ankle and wiggling your toes to encourage blood circulation when resting also helps reduce the risk of clot formations.

Return to driving: You *cannot drive* until you are no longer using crutches or a frame, which could be up to **6 weeks** or longer. You may need to contact your insurers to make sure you are covered to drive following surgery.

Return to hobbies/sports: Most people can return to active hobbies such as swimming, gardening or golf 3 months after their operation.

Air travel: Each airline has its own regulations about flying after surgery. Check with your airline before you fly. Although no fixed guidelines exist, *we do not advise travelling on a long-haul flight (more than 3 hours) for the first 3 months after your surgery.*

Follow-up Review at Outpatient Clinic

You will usually be seen by your surgeon around 6 – 12 weeks following your surgery. This is to check on your progress and ensure there are no problems with your new hip.

It is extremely important that you have been practising the exercises given to you by the physiotherapists while you were in hospital.

Once your surgeon is satisfied with your progress you will normally then be reviewed one year after your operation, at which time an up-to-date X-ray will be taken to ensure the hip is functioning as it should be.

Risks and Potential Problems Following Hip Replacement Surgery

This section describes the potential risks of having a hip replacement and other problems you may have as a result of having the surgery.

It is important that you read and understand this section before proceeding with the operation.

Pain

A very small number of patients experience more pain than they had experienced beforehand, while some patients experience no change in their pain levels. This is often because they develop one of the complications mentioned later in this section. Occasionally however, patients have pain afterwards, for which we can find no cause.

Infection

After having a hip replacement, serious infections around the hip are rare (less than 1%). However, if you smoke, are obese or have diabetes, your risk of getting an infection is increased.

Reducing your risk of infection

Smoking increases the risk of infection and you should stop smoking from the time of pre-operative assessment until at least 6 weeks after your operation. You will not be allowed to smoke whilst in hospital. Also, if you are morbidly obese, which is a BMI of over 40, your risk of getting an infection is increased. If you are diabetic it is important that your blood sugar is well controlled as poor control can increase the chance of wound infection and other complications.

Alcohol or other drug abuse can also increase the risk of wound problems. Other risk factors include having skin conditions, for example psoriasis.

Always follow wound care advice given by staff before discharge.

When to suspect a possible wound infection

It is normal to have a degree of bruising and swelling in the leg after the operation.

However, the following are signs of potential infection:

1. If your wound starts to leak fluid having been dry or continues to leak fluid beyond 7 days after your operation.
2. If part or the entire wound becomes swollen, red, sore to touch or starts to open.
3. If you get a sudden increase in pain around your hip and feel shivery and/or unwell.
4. An increase in body temperature.
5. Occasionally pain that fails to settle following surgery or pain that develops some time later can be caused by infection and in these cases the wound can remain normal. This is one cause of pain that can occur up to one year after surgery.

If you suspect you have an infection, especially in the first few weeks after surgery, you should contact The Ulster Independent Clinic for advice; alternatively, contact your GP urgently or attend Accident and Emergency.

In the event of an infection

Most infections occur in the first 7-10 days after the operation. Your surgeon will see you and may prescribe oral antibiotics with follow up wound checks and treatment at UIC or G.P Surgery. On the rare occasion that a serious infection is suspected during this first 6 weeks, you will be readmitted to hospital and go back to theatre to have the wound opened and washed out.

After the operation (washout) you will need to take antibiotics for approximately 3 months and may have to stay in hospital for up to 6 weeks. This is so that the antibiotics can be given through a drip into your veins. After this 6-week period, antibiotics can be taken by mouth at home for a further 6 weeks.

In the long term, if we are unable to clear the infection the new hip will have to be removed. This means having a "2 stage" revision operation. In the first operation your new hip joint is removed and then after a number of weeks or months in the second operation a new hip joint is put back in again. This finally clears the infection in 9 out of 10 cases.

Reducing your risk of getting a clot

Getting out of bed as soon as possible after your operation is now considered very important. At the moment there are no drugs that have been proven to reduce the small risk of death from a clot in the lung (pulmonary embolus). A balance has to be struck between thinning the blood to avoid a blood clot and avoiding too much bleeding from the operation.

You should also:

1. Take frequent deep breaths to make sure you fill your lungs properly.
2. Move both arms and legs. You are allowed to move both knees up and down in the bed. This will not harm your new hip and will help to ease the pain.
3. Carry out foot and ankle exercises every hour.
4. Take short walks regularly as able.

You may be given blood thinning medication to help prevent clots from forming after the operation.

Blood Loss

A minority of patients will require a blood transfusion because of blood loss caused by the operation.

Fractures

Fractures (breakage of the femur or the pelvis) can sometimes happen during the operation. It can happen putting in the cup or the stem and may require a different type of implant in order to fix it.

When to suspect a clot in your leg (Deep Vein Thrombosis)

It is quite common to get small clots in the calf of either leg following surgery. The majority DO NOT require any treatment. Clots above the knee are considered to be more serious but are much rarer. If you have a sudden increase in pain and swelling in your leg then a scan of your leg may be required. Treatment may be necessary in the form of medication to thin your blood. This is to help prevent the clot travelling to your lung. Advice regarding warning signs of DVT is given below:

Contact UIC Hospital Liaison Sister* if you have:

- Pain or tenderness in your leg
- Swelling or warmth in your leg
- Red or discoloured skin on your leg
- Veins that stick out

*Tel: 028 9066 1212 Ext. 2400

When to suspect a clot in your lung (pulmonary embolus)

In a small number of patients, clots can travel to the lung (PE - pulmonary emboli) and can cause death. The risk of dying from a clot in the lung nowadays is very small and is fewer than 1 per 1000 cases.


Following discharge from hospital, if you have any of the following symptoms, call 999 or go to A&E immediately:

- Shortness of breath
- Coughing up blood
- Sudden chest pain
- Painful breathing

Dislocation

Your new hip can become dislocated (come out of joint) but this happens in fewer than 1 in 100 patients.

To reduce the risk of dislocation, take the following precautions for **6 weeks post surgery**.



DO NOT bend at the hip more than 90 degrees. This means do not bend forward to pick up objects. Get help or use a 'helping hand' to put on socks and shoes, or to wash from your knees down.



This also means **DO NOT** sit in low chairs or sofas - your knee(s) must never be higher than your hip.



DO NOT twist your operated leg inwards when walking or lying. This means do not swivel on the spot to turn when walking - step around with small steps.

In bed, when lying on your back you should keep a pillow between your legs to prevent the operated leg rolling inwards.



DO NOT cross your legs.

This means when you are sitting, **DO NOT** sit with your ankles crossed, or with one leg over the other.

If your hip becomes dislocated you will usually:

1. Experience severe pain.
2. Notice your foot on the operated side is turned in but occasionally it can be turned out.
3. Notice the leg on the operated side is shorter.

If you think that your hip has dislocated, then it is best to go to your nearest Accident and Emergency Department. This means going by emergency ambulance because if your hip is dislocated it would be too painful to travel by car.

Nerve Damage

The most common nerve problem after surgery is a “foot drop” and this occurs in fewer than 1 in 200 patients. This happens when the sciatic nerve, which supplies power and the feeling to your lower leg and foot, becomes stretched at the time of the operation. If this happens, it means that when you are walking your foot tends to drop and you are at risk of tripping easily. To help with this, you will have to wear a splint on your foot until the nerve recovers. Approximately 4 out of 5 patients make a good recovery from this, but it can take up to 18 months.

The femoral nerve which supplies power and feeling to the front of your thigh and knee can be damaged, but this is even less common.

Leg Length

During surgery we try and ensure that we restore the correct length in your new hip. Sometimes, for various reasons, your legs may not be the same length after surgery. Often before the operation your painful leg can feel shorter, whereas afterwards your operated leg may feel longer.

Delirium

Delirium means confusion and can affect up to 11% of patients following surgery. It usually resolves completely over several days but for some patients it can persist long-term and result in a loss of independent living.

Other Potential Problems

Nausea or feeling sick

Loss of appetite

Constipation

Hiccoughs (hiccups)

Chest infection

Kidney or
bladder infection

Atrial fibrillation / irregular
heartbeat

Low mood / depression

Tiredness

Trochanteric bursitis
(inflammation of the
fat pad around the hip)

How serious is this operation?

Having your hip replaced is a major operation. Your consultant will discuss with you pre-operatively common and rare complications which may arise. However, with any surgery in the Ulster Independent Clinic, all the staff involved in your care aim to provide as positive an experience as possible for you and your family.

We hope that this information booklet has been useful in providing insight into the hip replacement process. You will, of course, have the opportunity to discuss any aspect of the surgery with your surgeon and anaesthetist before proceeding.

If you would like any additional information about The Ulster Independent Clinic, please visit our website or call us:

Visit: www.ulsterindependentclinic.com

Call: 028 9066 1212



245 Stranmillis Road,
Belfast, BT9 5JH

T: 028 9066 1212

www.ulsterindependentclinic.com

Issued March 2019 / Revised August 2025 / UIC913