




BRGSF-HIS and CD3ε epitope humanized mice: Translatable preclinical mouse models for assessment of T-cell engagers-induced CRS

Perrine Martin-Jeantet¹; Florence Renart-Depontieu¹; Ludovic Bourre²; Gaëlle Martin¹; Angela Pappalardo¹; Dean O. Campbell²; Astrid Doerner²; Yacine Cherifi¹; Fabiane Sônego¹; Kader Thiam¹
¹genOway, Lyon, 69007, France; ²Crown Bioscience, 16550 West Bernardo Drive, Building 5, Suite 525, San Diego, CA, USA


Background: T cell engagers show high efficacy in B cells malignances. High risk of immune-related adverse events, including cytokine release syndrome (CRS), is reported in patients due to on-target off-site effects of T cell engagers. Thus, reliable and translational mouse models are required to predict potential safety issues and investigate their rescue. Here we describe two preclinical models mimicking to some extent CRS induction upon activation with anti-CD3 antibodies.

BRGSF-HIS mice


Model features

- BRGSF (Balb/C Rag2^{-/-} IL2Rγ^{-/-}, SIRPα^{NOD} and Flt3^{-/-}):
 - Highly immunodeficient with reduced murine myeloid cells
 - Normal life span (no anemia, no weight loss, normal fur texture & integrity, normal activity & posture)
 - Upon hCD34⁺ cells injection:
 - Lymphoid and Myeloid compartment development
 - Boost by hFlt3L enhances human myeloid cell development and accumulation
- 
- Humanization level in blood (12 weeks/QC)
 - Increase over time of T cells proportions
 - Flt3L treatment boosts human myeloid cells proportions

OKT3-induced CRS is enhanced by myeloid compartment and can be mitigated by Tocilizumab




Crosstalk between immune system & non-immune cells



	BRGSF-HIS	hCD3ε KI
Pros	- Human Immune response - Versatility as TAA and immune targets are human	Fully functional crosstalk among tumor cells, immune cells and stroma
Cons	Partial crosstalk among immune cells and stroma	- Mouse Immune response - Requires TAA humanization


BLINCYTO® (Blinatumomab) induces cytokines release



CD3ε epitope humanized mice

Model features


- CD3ε N-terminal epitope knock-in mice (hCD3ε mice) expressing the humanized epitope of anti-CD3 clone SP34:
 - Preserves the overall mouse CD3 complex functionality
 - Preserves the homeostasis of the immune system




CD3ε humanization does not alter T cells functions

Ex vivo T cell activation


T cells isolation from spleen activated with anti-CD3 clones 2C11 or SP34 (10μg/ml) for 72h. n=7 - mean +/- SEM



T-B cells cooperation




T cell engager-induced anti-tumor response



Bispecific Ab T-cell engager induces cytokine production

Ex vivo co-culture of Hepa1-6 + splenocytes for 24h in presence of:
BsAb1: anti-CD3 derived from SP34; TAA expressed on Hepa1-6 cell
BsAb2: negative control - anti-CD3 derived from SP34; TAA not expressed on Hepa1-6 cell



Conclusion: BRGSF-HIS and hCD3ε models are complementary to assess T cell engager-induced CRS in addition to efficacy.

www.genOway.com