



Sustained delivery of low-dose anti-CTLA-4 by genetically engineered encapsulated cells drives tumor response and prolongs survival in a colorectal cancer model



Julien Grogg^{a,b,d*}, Emily Charrier^{a,b,d}, Remi Vernet^{a,b}, Muriel Urwyler^{a,b}, Olivier Von Rohr^{a,b}, Valentin Saingier^{a,b}, Fabien Courtout^{a,b}, Aurélien Lathuilière^c, Adrien Engel^{a,b,d} & Nicolas Mach^{a,b}

^a Department of Oncology, Geneva University Hospitals and Medical School, 1211 Geneva, Switzerland; ^b Centre for Translational Research in Onco-Hematology, Oncology Division, Geneva University Hospital and University of Geneva, 1211 Geneva, Switzerland; ^c Department of Rehabilitation and Geriatrics, University of Geneva, 1211 Geneva, Switzerland; ^d MaxiVAX SA, 1202 Geneva, Switzerland.


ABSTRACT

Systemic therapy with CTLA-4 blocking antibody (aCTLA4) restores endogenous antitumor immunity and induces remarkable long-term clinical benefits in patients with melanoma. Yet immune-related side effects remain a major hurdle to extend its label to many more types of cancer. Intra- and peritumoral administration of aCTLA4 has recently emerged to optimize its dose/efficacy ratio while preventing its on-target, off-tumor systemic toxicities. Sustained delivery of low-dose aCTLA4 by genetically engineered encapsulated cells (MVX-3) could offer a promising option for cancer treatment addressing the shortcomings of systemic therapy.

MATERIALS AND METHODS



RESULTS



A. aCTLA4 secretion rate from MVX-3 before and after 7 days in vivo. B. aCTLA4 was detected in the serum at day 17 in both the MVX-3 and the ipilimumab (10mg/kg) treated conditions. C. Mean tumor volume per group over time. D. Average tumor growth rate from day 1 after tumor engraftment until end of the study determined by the method described by Hather et al. (2014). E. Proportion of Intratumoral CD8+ cells, Tregs (Foxp3+) and Human CTLA4+ cells at day 17. F. Kaplan-Meier curves showing the probability of survival over time. G. Response rate and toxic death rate per group. Results are expressed as mean $\pm$ standard deviation (SD). Statistical significance was assessed by Student's t test for comparison between two groups or by Log-rank (Mantel-Cox) for the Kaplan-Meier curve. Statistical significance between groups is presented as follows: *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001. Data analyses were performed using the software package GraphPad Prism 9 (GraphPad Software).

CONCLUSIONS

- Peritumoral administration of MVX-3 induced durable complete tumor rejection (2/7) and tumor growth control (4/7) when administered at doses 1'000 times lower than i.p. ipilimumab, whereas rapid tumor growth without any tumor rejection were observed in negative control mice.
- I.p. ipilimumab induced durable complete tumor rejection (9/12), while treatment related toxicities upon dosing led to premature mice termination (3/12).
- MVX-3 was found as equally effective as i.p. ipilimumab in decreasing the proportion of CTLA4+ helper and regulatory T cells in the tumor at Day 7 post treatment.
- Survival was also improved by MVX-3 compared to control.

These findings suggest that a sustained, controlled delivery of low-dose aCTLA4 by genetically engineered encapsulated cells could achieve similar therapeutic benefit as the systemic therapy, without the commonly associated severe toxicities. The safety and biological efficacy profile of MVX-3 encourage further preclinical and clinical explorations.



Copies of this poster obtained through QR codes are for personal use only and may not be reproduced without written permission of the authors.

Corresponding author e-mail: jgrogg@maxivax.ch

Funding Source: Funded by Innosuisse (CTI), Coromandel, Philanthropy Settlement, and MaxiVAX SA.

Disclosure: N. Mach is a founder and minority stockholder at MaxiVAX SA. All other authors have declared no conflicts of interest.