

Unity Fetal Risk™ Screen: Carrier Screen Frequencies

Disease	Gene	NM#	Ethnicity	Carrier Frequency Before Testing	Detection Rate	Carrier Risk After Testing
Alpha-Thalassemia¹ "Silent" group indicates a single alpha-globin gene deletion (aa/a-). "Other" group includes other HBA1/HBA2 variants such as double deletions (aa/-- or a-/a-) and non-deletion variants, including Hb Constant Spring.	HBA1/ HBA2	NM_000558.5 NM_000517.6	African American	Silent: 1 in 3 Other: 1 in 28	>95%	Silent: 1 in 60 Other: 1 in 560
			Asian*	Silent: 1 in 13 Other: 1 in 41	>95%	Silent: 1 in 260 Other: 1 in 820
			Northern European	Silent: 1 in 51 Other: <1 in 1,000	>95%	Silent: 1 in 1,020 Other: 1 in 20,000
			Hispanic	Silent: 1 in 17 Other: 1 in 398	>95%	Silent: 1 in 340 Other: 1 in 7,960
			General Population	Silent: 1 in 16 Other: 1 in 112	>95%	Silent: 1 in 320 Other: 1 in 2,240
Cystic Fibrosis²	CFTR	NM_000492.4	African American	1 in 84	>99%	<1 in 8,400
			Ashkenazi Jewish	1 in 29	>99%	<1 in 2,900
			Asian*	1 in 118	>99%	<1 in 11,800
			Northern European	1 in 25	>99%	<1 in 2,500
			Hispanic	1 in 59	>99%	<1 in 5,900
			General Population	1 in 38	>99%	<1 in 3,800
Sickle Cell Disease, Beta-Thalassemia, Hemoglobinopathies¹	HBB	NM_000518.5	African American	1 in 8	>99%	<1 in 800
			Ashkenazi Jewish	1 in 32	>99%	<1 in 3,200
			Asian*	1 in 22	>99%	<1 in 2,200
			Northern European	1 in 164	>99%	<1 in 16,400
			Hispanic	1 in 55	>99%	<1 in 5,000
			General Population	1 in 33	>99%	<1 in 3,300
Spinal Muscular Atrophy³	SMN1	NM_000344.4	African American	1 in 72	>90.3%	<1 in 800
			Ashkenazi Jewish	1 in 67	>92.8%	<1 in 3,200
			Asian*	1 in 59	>93.6%	<1 in 2,200
			Northern European	1 in 47	>95%	<1 in 16,400
			Hispanic	1 in 68	>92.6%	<1 in 5,000
			General Population	1 in 54	>91.2%	<1 in 3,300
Fragile X Syndrome⁴	FMR1	NM_002024.6	African American	1 in 267	>99%	<1 in 26,700
			Ashkenazi Jewish	1 in 102	>99%	<1 in 10,000
			Asian*	1 in 419	>99%	<1 in 41,900
			Northern European	1 in 170	>99%	<1 in 17,000
			Hispanic	1 in 253	>99%	<1 in 25,300
			General Population	1 in 201	>99%	<1 in 20,100

Unity Fetal Risk™ Screen: Carrier Screen Frequencies

Disease	Gene	NM#	Ethnicity	Carrier Frequency Before Testing	Detection Rate	Carrier Risk After Testing
Canavan Disease ¹	ASPA	NM_000049.4	African American	1 in 2,126	>99%	1 in 21,2600
			Ashkenazi Jewish	1 in 44	>99%	1 in 4,400
			Southeast Asian	1 in 1,470	>99%	< 1 in 50,000
			Northern European	1 in 428	>99%	1 in 42,800
			Hispanic	1 in 1,153	>99%	<1 in 50,000
			General Population	1 in 439	>99%	1 in 43,900
DMD-Associated Dystrophinopathies ^{1*}	DMD	NM_004006.3	General Population	1 in 717	>99%	<1 in 50,000
Familial Dysautonomia ¹	ELP1	NM_003640.5	African American	1 in 1,063	>99%	<1 in 50,000
			Ashkenazi Jewish	1 in 35	>99%	1 in 3,500
			East Asian	1 in 1,285	>99%	<1 in 50,000
			Southeast Asian	1 in 4,411	>99%	<1 in 50,000
			Northern European	1 in 567	>99%	<1 in 50,000
			Hispanic	1 in 824	>99%	1 in 8,2400
Medium-Chain Acyl-CoA Dehydrogenase Deficiency ^{5,6,7}	ACADM	NM_000016.6	African American	1 in 217	>99%	1 in 21,700
			Ashkenazi Jewish	1 in 134	>99%	1 in 13,400
			East Asian	1 in 335	>99%	1 in 33,500
			Southeast Asian	1 in 272	>99%	1 in 27,200
			Northern European	1 in 62	>99%	1 in 6,200
			Hispanic	1 in 92	>99%	1 in 9,200
Phenylalanine Hydroxylase Deficiency ^{5,8,9,10}	PAH	NM_000277.3	African American	1 in 145	>99%	1 in 14,500
			Ashkenazi Jewish	1 in 17	>99%	1 in 1,700
			East Asian	1 in 50	>96%	1 in 1,563
			Northern European	1 in 50	>99%	1 in 5,000
			Hispanic	1 in 69	>99%	1 in 16,200
			Turkish	1 in 26	>99%	1 in 2,600
PMM2-Congenital Disorder of Glycosylation ^{5,11,12,13}	PMM2	NM_000303.3	African American	1 in 210	>99%	1 in 21,000
			Ashkenazi Jewish	1 in 61	>99%	1 in 6,100
			East Asian	1 in 235	>99%	1 in 23,500
			Southeast Asian	1 in 309	>99%	1 in 30,900
			Northern European	1 in 72	>99%	1 in 7,200
			Hispanic	1 in 101	>99%	1 in 13,500
			General Population	1 in 70	>99%	1 in 7,000

Unity Fetal Risk™ Screen: Carrier Screen Frequencies

Disease	Gene	NM#	Ethnicity	Carrier Frequency Before Testing	Detection Rate	Carrier Risk After Testing
Smith-Lemli-Opitz Syndrome¹⁴	<i>DHCR7</i>	NM_001360.3	African American	1 in 183	>99%	1 in 18,300
			Ashkenazi Jewish	1 in 43	>99%	1 in 4,300
			Southeast Asian	1 in 417	>99%	<1 in 50,000
			Northern European	1 in 51	>99%	1 in 5,100
			Hispanic	1 in 167	>99%	1 in 16,700
			General Population	1 in 71	>99%	1 in 7,100
Tay Sachs Disease^{1,15,16}	<i>HEXA</i>	NM_000520.6	African American	1 in 257	>99%	1 in 25,700
			Ashkenazi Jewish	1 in 28	>99%	1 in 2,800
			East Asian	1 in 857	>99%	<1 in 50,000
			Southeast Asian	1 in 630	>99%	1 in 63,000
			Northern European	1 in 191	>99%	1 in 19,100
			Hispanic	1 in 316	>99%	1 in 31,600
			French Canadian	1 in 53	>99%	1 in 5,300
			Irish	1 in 50	>99%	1 in 5,000
			General Population	1 in 193	>99%	1 in 19,300

* For X-linked diseases, female carrier frequencies are presented.

The reported detection rates are derived from literature-based estimates of the fraction of disease alleles, adjusted by frequency, that are not detectable by this methodology.

References

- Westemeyer, M., Saucier, J., Wallace, J., Prins, S. A., Shetty, A., Malhotra, M., Demko, Z. P., Eng, C. M., Weckstein, L., Boostanfar, R., Rabinowitz, M., Benn, P., Keen-Kim, D., & Billings, P. (2020). Clinical experience with carrier screening in a general population: support for a comprehensive pan-ethnic approach. *Genetics in medicine : official journal of the American College of Medical Genetics*, 22(8), 1320–1328. <https://doi.org/10.1038/s41436-020-0807-4>
- Rohlf, E. M., Zhou, Z., Heim, R. A., Nagan, N., Rosenblum, L. S., Flynn, K., Scholl, T., Akmaev, V. R., Sirko-Osadsa, D. A., Allitto, B. A., & Sugarman, E. A. (2011). Cystic fibrosis carrier testing in an ethnically diverse US population. *Clinical chemistry*, 57(6), 841–848. <https://doi.org/10.1373/clinchem.2010.159285>
- Sugarman, E., Nagan, N., Zhu, H. et al. Pan-ethnic carrier screening and prenatal diagnosis for spinal muscular atrophy: clinical laboratory analysis of >72 400 specimens. *Eur J Hum Genet* 20, 27–32 (2012). <https://doi.org/10.1038/ejhg.2011.134>
- Owens, K. M., Dohany, L., Holland, C., DaRe, J., Mann, T., Settler, C., & Longman, R. E. (2018). FMR1 premutation frequency in a large, ethnically diverse population referred for carrier testing. *American journal of medical genetics. Part A*, 176(6), 1304–1308. <https://doi.org/10.1002/ajmg.a.38692>
- gnomAD. (2025). gnomAD v4.1.0 [Database]. Broad Institute of MIT and Harvard. <https://gnomad.broadinstitute.org/>
- Grosse, S. D., Khoury, M. J., Greene, C. L., Crider, K. S., & Pollitt, R. J. (2006). The epidemiology of medium chain acyl-CoA dehydrogenase deficiency: An update. *Genetics in Medicine*, 8(4), 205–212
- Therrell, B. L., Jr., Lloyd-Puryear, M. A., Camp, K. M., & Mann, M. Y. (2014). Inborn errors of metabolism identified via newborn screening: Ten-year incidence data and costs of nutritional interventions for research agenda planning. *Molecular Genetics and Metabolism*, 113(1–2), 14–26.
- Scriver, C. R., & Kaufman, S. (2001). The hyperphenylalaninemias: Phenylalanine hydroxylase deficiency. In C. R. Scriver, A. L. Beaudet, W. S. Sly, D. Valle, B. Childs, K. W. Kinzler, & B. Vogelstein (Eds.), *The metabolic and molecular bases of inherited disease* (8th ed., pp. 1667–1724). McGraw-Hill.
- Ozalp, I., Coşkun, T., Tokatli, A., Kalkanoğlu, H. S., Dursun, A., Tokol, S., Köksal, G., Özgüç, M., & Köse, R. (2001). Newborn PKU screening in Turkey: At present and organization for future. *The Turkish Journal of Pediatrics*, 43(2), 97–101.
- Hillert, A., Anikster, Y., Belanger-Quintana, A., Burlina, A., Burton, B. K., Carducci, C., Chiesa, A. E., Christodoulou, J., Đorđević, M., Desviat, L. R., Eliyahu, A., Evers, R. A. F., Fajkusova, L., Feillet, F., Bonfim-Freitas, P. E., Giżewska, M., Gundorova, P., Karall, D., Kneller, K., ... Blau, N. (2020). The genetic landscape and epidemiology of phenylketonuria. *American Journal of Human Genetics*, 107(2), 234–250. <https://doi.org/10.1016/j.ajhg.2020.06.006>
- Shi, L., Webb, B. D., Birch, A. H., Elkhoury, L., McCarthy, J., Cai, X., Oishi, K., Mehta, L., Diaz, G. A., Edelman, L., & Kornreich, R. (2017). Comprehensive population screening in the Ashkenazi Jewish population for recurrent disease-causing variants. *Clinical Genetics*, 91(4), 599–604
- Matthijs, G., Schollen, E., Bjursell, C., Erlandson, A., Freeze, H., Imtiaz, F., Kjaergaard, S., Martinsson, T., Schwartz, M., Seta, N., Vuillaumier-Barrot, S., Westphal, V., & Winchester, B. (2000). Mutations in PMM2 that cause congenital disorders of glycosylation, type Ia (CDG-Ia). *Human Mutation*, 16(5), 386–394.
- Jaeken, J., & Matthijs, G. (2001). Congenital disorders of glycosylation. *Annual Review of Genomics and Human Genetics*, 2, 129–151. <https://doi.org/10.1146/annurev.genom.2.1.129>
- Lazarin, G. A., Haque, I. S., Evans, E. A., & Goldberg, J. D. (2017). Smith-Lemli-Opitz syndrome carrier frequency and estimates of in utero mortality rates. *Prenatal diagnosis*, 37(4), 350–355. <https://doi.org/10.1002/pd.5018>
- Triggs-Raine, B., Richard, M., Wasel, N., Prence, E. M., & Natowicz, M. R. (1995). Mutational analyses of Tay-Sachs disease: Studies on Tay-Sachs carriers of French Canadian background living in New England. *American Journal of Human Genetics*, 56(4), 870–879.
- National Tay-Sachs & Allied Diseases Association. (n.d.). Tay-Sachs disease. https://tay-sachs.org/taysachs_disease.php