**VantageAg** 

Ingredients: Phosphoric Acid and Sulfuric Acid



## Phosphate

Better for your crops.

Better for the environment.

NET CONTENTS

**20**L

GENERAL INFORMATION: Vantage Ag - Phosphate is a unique formulation for use when plant available phosphate is deficient or low, based on soil and/or tissue samples. It should be used as part of a comprehensive nutrient plan for plant growth, development, yield and quality.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. ALWAYS READ LABEL BEFORE USING.

DIRECTIONS FOR USE: This highly bio-available formulation is designed for foliar applications but can also be applied as a band in the soil or injected through drip or micro-sprinklers. It will work in all soil types and conditions. This formulation can be used on a wide range of crops. Apply Vantage Ag - Phosphate using recommended rates and directions with your phosphate management plans.

APPLICATION RATES: Foliar applied: Apply 250 – 870 mL of product in a minimum of 95 L of water per hectare (equivalent to 100–350 mL in 10 gallons of water per acre). Repeat as needed. Soil applied: Apply 310 – 1090 mL of product per hectare (equivalent to 125–440 mL per acre) in furrow. Do not apply to seed.

COMPATIBLITY: This product will mix with other products, but always conduct a jar test first. Order of mixing; fill sprayer tank to 1/2 full, add pesticide, continue to fill tank and agitate. When tank is 3/4 full, add Phosphate to the tank last.

CAUTION: CORROSIVE. Do not eat, drink or smoke while using this product. Wear protective gloves and clothing and eye, face protection

and a NIOSH approved N95 respirator. Wash hands thoroughly with soap and water after handling. Avoid release into aquatic environments and do not pour down drains.

STORAGE: CORROSIVE. Protect from freezing, product will become inactive. Store in a cool dry place.

FIRST AID MEASURES: If on skin: Remove contaminated clothing. Wash skin thoroughly with soap and water. Wash clothing before reuse. If in eyes: Remove contact lenses if present and easy to do so and flush with running water for at least 15 minutes while holding eyelids open. Get medical attention. Ingestion: Drink 1-2 glasses of water and induce vomiting. Never induce vomiting or give anything by mouth to an unconscious person. Call physician immediately. Inhalation: Remove to fresh air. If breathing is difficult, administer oxygen.

WARRANTY DISCLAIMER AND NOTICE: The directions for use of this product are believed to be adequate and should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Vantage Ag, the manufacturer or seller. The manufacturer or sellers make no warranties, guarantees, conditions or representations of any kind to buyer or user. Tests have not been carried out under all possible conditions and the seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

In no event shall Vantage Ag, the manufacturer or seller be liable for consequential, special or indirect damages resulting from the use, handling, application, storage or disposal of this product or for damages in the nature of penalties and the buyer and user waive any right they may have to such damages.

A VANTAGE AG
Unit #1
6285 - 205th Street
Langley, BC V2Y 1N7
1-778-919-3370

