




I'm not a robot   
reCAPTCHA

**I am not a robot!**

### Eus procedure code.




The text is about the removal of redundant code and references to studies on endoscopic ultrasound-guided drainage of pancreatic fluid collections. If there's a valid reason to keep it, feel free to put it back! The studies listed are: Sj End of HubSpot Embed Code -> Transgastric drainage of pancreatic pseudocysts and abscesses has been a topic of interest in recent studies. A review of the literature reveals several approaches, including endoscopic ultrasound-guided drainage using therapeutic echo endoscopes (Giovannini et al., 2001) and lumen-apposing stents (Gornals et al., 2013). Additionally, transenteric drainage with fully covered self-expanding metallic stents has also been explored (Talreja et al., 2008). In some cases, a combination of endoscopic and percutaneous drainage may be necessary for optimal results. For instance, Ross et al. (2010) described combined endoscopic and percutaneous drainage of organized pancreatic necrosis. It's worth noting that transgastric drainage is not always the most effective approach, as evidenced by Varadarajulu et al.'s (2011) description of a multiple transluminal gateway technique for EUS-guided drainage of symptomatic walled-off pancreatic necrosis. In any case, when searching for codes or reference materials, it's essential to ensure that the method being described is accurately represented. A code such as 48999 or 48520 may be relevant, but it's crucial to verify the specifics of the procedure before applying them in practice. References: Giovannini M, Pesenti C, Rolland AL, et al. Endoscopic ultrasound-guided drainage of pancreatic pseudocysts or pancreatic abscesses using a therapeutic echo endoscope. *Endoscopy*. 2001;33:473-7. Gornals JB, De la Serna-Higuera C, Sanchez-Yague A, et al. Endosonography-guided drainage of pancreatic fluid collections with a novel lumen-apposing stent. *Surg Endosc*. 2013;27:14-28-34. Ross A, Gluck M, Irani S, et al. Combined endoscopic and percutaneous drainage of organized pancreatic necrosis. *Gastrointest Endosc*. 2010;71:79-84. Talreja JP, Shami VM, Ku J, et al. Transenteric drainage of pancreatic fluid collections with fully covered self-expanding metallic stents (with video). *Gastrointest Endosc*.

If there's a valid reason to keep it, feel free to put it back! The studies listed are: Sj End of HubSpot Embed Code -> Transgastric drainage of pancreatic pseudocysts and abscesses has been a topic of interest in recent studies. A review of the literature reveals several approaches, including endoscopic ultrasound-guided drainage using therapeutic echo endoscopes (Giovannini et al., 2001) and lumen-apposing stents (Gornals et al., 2013).




Eus guided cystogastrostomy. Eus guided cystogastrostomy cpt. Eus cystogastrostomy cpt.

The text is about the removal of redundant code and references to studies on endoscopic ultrasound-guided drainage of pancreatic fluid collections. If there's a valid reason to keep it, feel free to put it back! The studies listed are: Sj End of HubSpot Embed Code -> Transgastric drainage of pancreatic pseudocysts and abscesses has been a topic of interest in recent studies. A review of the literature reveals several approaches, including endoscopic ultrasound-guided drainage using therapeutic echo endoscopes (Giovannini et al., 2001) and lumen-apposing stents (Gornals et al., 2013). Additionally, transenteric drainage with fully covered self-expanding metallic stents has also been explored (Talreja et al., 2008). In some cases, a combination of endoscopic and percutaneous drainage may be necessary for optimal results. For instance, Ross et al.



If there's a valid reason to keep it, feel free to put it back! The studies listed are: SJ End of HubSpot Embed Code -> Transgastric drainage of pancreatic pseudocysts and abscesses has been a topic of interest in recent studies. A review of the literature reveals several approaches, including endoscopic ultrasound-guided drainage using therapeutic echo endoscopes (Giovannini et al., 2001) and lumen-apposing stents (Gornals et al., 2013). Additionally, transenteric drainage with fully covered self-expanding metallic stents has also been explored (Talreja et al., 2008). In some cases, a combination of endoscopic and percutaneous drainage may be necessary for optimal results. For instance, Ross et al. (2010) described combined endoscopic and percutaneous drainage of organized pancreatic necrosis. It's worth noting that transgastric drainage is not always the most effective approach, as evidenced by Varadarajulu et al.'s (2011) description of a multiple transluminal gateway technique for EUS-guided drainage of symptomatic walled-off pancreatic necrosis. In any case, when searching for codes or reference materials, it's essential to ensure that the method being described is accurately represented. A code such as 48999 or 48520 may be relevant, but it's crucial to verify the specifics of the procedure before applying them in practice. References: Giovannini M, Pesenti C, Rolland AL, et al. Endoscopic ultrasound-guided drainage of pancreatic pseudocysts or pancreatic abscesses using a therapeutic echo endoscope. *Endoscopy*. 2001;33:473-7. Gornals JB, De la Serna-Higuera C, Sánchez-Yague A, et al. Endosonography-guided drainage of pancreatic fluid collections with a novel lumen-apposing stent. *Surg Endosc*. 2013;27:14-28-34. Ross A, Gluck M, Irani S, et al. Combined endoscopic and percutaneous drainage of organized pancreatic necrosis. *Gastrointest Endosc*. 2010;71:79-84. Talreja JP, Shami VM, Ku J, et al. Transenteric drainage of pancreatic-fluid collections with fully covered self-expanding metallic stents (with video). *Gastrointest Endosc*. 2008;68:1,199-203. Varadarajulu S, Phadnis MA, Christein JD, et al. Multiple transluminal gateway technique for EUS-guided drainage of symptomatic walled-off pancreatic necrosis. *Gastrointest Endosc*. 2011;74:74-80.



For instance, Ross et al. (2010) described combined endoscopic and percutaneous drainage of organized pancreatic necrosis. It's worth noting that transgastric drainage is not always the most effective approach, as evidenced by Varadarajulu et al.'s (2011) description of a multiple transluminal gateway technique for EUS-guided drainage of symptomatic walled-off pancreatic necrosis. In any case, when searching for codes or reference materials, it's essential to ensure that the method being described is accurately represented. A code such as 48999 or 48520 may be relevant, but it's crucial to verify the specifics of the procedure before applying them in practice. References: Giovannini M, Pesenti C, Rolland AL, et al. Endoscopic ultrasound-guided drainage of pancreatic pseudocysts or pancreatic abscesses using a therapeutic echo endoscope. *Endoscopy*. 2001;33:473-7. Gornals JB, De la Serna-Higuera C, Sánchez-Yague A, et al. Endosonography-guided drainage of pancreatic fluid collections with a novel lumen-apposing stent. *Surg Endosc*. 2013;27:14-28-34. Ross A, Gluck M, Irani S, et al. Combined endoscopic and percutaneous drainage of organized pancreatic necrosis. *Gastrointest Endosc*. 2010;71:79-84. Talreja JP, Shami VM, Ku J, et al. Transenteric drainage of pancreatic-fluid collections with fully covered self-expanding metallic stents (with video). *Gastrointest Endosc*. 2008;68:1,199-203. Varadarajulu S, Phadnis MA, Christein JD, et al. Multiple transluminal gateway technique for EUS-guided drainage of symptomatic walled-off pancreatic necrosis. *Gastrointest Endosc*. 2011;74:74-80.