

TEST NAME: Cardio-Check

3702 Cardio Check - Serum

Methodology: Enzymatic, Immunoturbidimetric, Chemiluminescent and Turbidimetric/Immunoturbidimetric

Cardio Check

Lipids

Cholesterol

Reference
Range

LDL Cholesterol	168	<= 99 mg/dL
HDL Cholesterol	64	>= 40 mg/dL
Triglycerides	74	<= 149 mg/dL
Total Cholesterol	247	150-199 mg/dL

Cholesterol Ratios

Triglycerides/HDL*	1.2	<= 2.0
Total Cholesterol/HDL*	3.86	<= 5.99

Lipoproteins

Reference
Range

Apo A-1	165	95-186 mg/dL
Apo B	127	49-173 mg/dL
Apo B / Apo A-1*	0.77	<= 0.95
Lp(a)	67	<= 29 mg/dL

**CARDIOVASCULAR
RISK**

Independent Risk Factors

Cardiometabolic Markers

Reference
Range

hs-CRP	2.19	<= 0.99 mg/L
Homocysteine*	9.2	5.2-11.4 µmol/L
Insulin *	2.4	1.9-23.0 µIU/mL

Sex Hormone Markers

Reference
Range

Testosterone, Total	5.52	1.75-7.81 ng/mL
Sex Hormone Binding Globulin	32.1	13.3-89.5 nmol/L


TEST NAME: Cardio-Check

Commentary

The performance characteristics of all assays have been verified by Genova Diagnostics Inc. All Assays have been cleared by the U.S. Food and Drug Administration.

Commentary is provided to the practitioner for educational purposes and should not be interpreted as diagnostic or as treatment recommendations. Diagnosis and treatment decisions are the practitioner's responsibility.

The **Reference Range** is a statistical interval representing 95% or 2 Standard Deviations (2 S.D.) of the reference range population. One Standard Deviation (1 S.D.) is a statistical interval representing ~68% of the reference population. Values between 1 and 2 S.D. are not necessarily abnormal. Clinical Correlation is suggested.



*Additional Interpretive Information:

Triglycerides/HDL Ratio: The expected range is based on a thorough analysis of the peer-reviewed literature regarding the relationship between Trg/HDL ratio and cardiometabolic risk. The increased risk associated with higher values are highlighted in *Diabetes Metab Syndr* 2019; 13(1):382-388. More information can be found in the Cardio Check support guide and the literature listed below:

<https://pubmed.ncbi.nlm.nih.gov/30641729/>

Total Cholesterol/HDL Ratio: The expected range is based on a thorough analysis of the peer-reviewed literature regarding the relationship between TC/HDL ratio and cardiovascular disease risk. The increased risk associated with higher values are highlighted in *J Am Col Card* 2010; 55(1):35-41. More information can be found in the Cardio Check support guide and the literature listed below.

<https://pubmed.ncbi.nlm.nih.gov/20117361/>

Apo B/ Apo A-1: The expected range is based on a thorough analysis of the peer-reviewed literature regarding the relationship between the Apo B/ Apo A-1 ratio and cardiovascular disease risk. The increase risk associated with higher values are highlighted in *Annals of Internal Medicine* 2007; 146:640-8. More information can be found in the Cardio Check support guide and literature listed below:

<https://pubmed.ncbi.nlm.nih.gov/17470832/>

Homocysteine: The reference range for the biomarker Homocysteine is based on the sex-specific 5th to 95th percentile values for men and women (20 to 39 years of age) in the NHANES nutritionally replete cohort. *Annals of Internal Medicine* 1999; 141 (331-338).

Insulin: Fasting insulin levels over 8.57 µU/ml were shown to be predictors of metabolic syndrome development.