Preventing and detecting cancers

UK born after 1960 will be diagnosed with some form of cancer during their lifetime. Even more concerning is that cancer rates among under 50s in developed countries have risen sharply in the past 30 years, prompting epidemiologists to describe this as an 'epidemic'. While 38% of cancers are preventable, including

Research suggests that 1 in 2 people in the

lung, bowel, breast and melanoma skin cancer; others are not. This makes early detection one of the most powerful tools for improving outcomes.

At Preventicum, we are committed to helping our clients reduce risk factors and detect potential cancers as early as possible, when treatment outcomes are most effective.



new cases a year



1 in 2 will be diagnosed with cancer in their lifetime

cancer risk factors An individuals' risk for developing cancer depends on

Understanding

a combination of age, genetics and lifestyle factors including:

- for 5% of cancer cases Overexposure to UV radiation, occupational
 - exposure to substances, exposure to specific

Smoking - the leading cause of cancer


Being overweight or obese accounts

- infections, alcohol and eating too little fibre account for 3-4% of cancer cases **Exposure to ionising radiation and** eating processed meat account for 1-2% of cancer cases
 - Air pollution and a sedentary lifestyle account for 1% or less of cancer cases

ORAL CONTRACEPTIVES

Number of preventable cancer

cases and associated risk factors:



countries, highlighting the importance of health assessments from a younger age. Analysis of the G20 countries found that cancer rates among 25-29 year olds increased by

Rising cancer incidence

22% between 1990 and 2019, with colorectal cancer up by 70% in those aged 15-39. Cancer is now the leading cause of death for young adults in upper-middle income countries. A study published in BMJ Oncology found that

global, early-onset cancer cases rose by nearly

in younger adults

Recent data identified a significant rise in cancer diagnoses among people under 50 in developed

80% since 1990 and cancer-related deaths increased by 27%. Although there is no single explanation for this trend, it reinforces the value of early and regular health screening.



increase in colorectal cancer

increase in cancer

rates in the 25-29

year old age range



Lifestyle choices increase chance of developing cancer



as well as heart disease, stroke and many other conditions. We use advanced imaging, detailed consultations and extensive laboratory testing to provide a detailed picture of your current health and future risk.

Early detection remains one of our most powerful tools against cancer. At Preventicum, our world class Optimal and Ultimate Assessments detect the earliest signs of and risk factors for at least 15 cancers,

MRI brain Ultrasound thyroid

LiverMultiScan MRI abdomen MRI abdomen Ultrasound abdomen Ultrasound abdomen MRI abdomen



*For men over 45 or a family history of prostate cancer

Prostate cancer At Preventicum, our Optimal+ and Ultimate+ Prostate cancer is the most common cancer in men in the UK, accounting for 27% of all Assessments are for men over 45 or a new cancer cases. 12,000 men die from family history of prostate cancer. prostate cancer in the UK annually, making it the second most common cause of cancer These assessments include: death in men, though mortality rates have Consultation with your Preventicum Doctor fallen by 10% which may be related to PSA blood testing advancements in early detection and treatment. **# Ultrasound volume scan** Key risk factors include age, family history Biparametric MRI scan of the prostate, and ethnicity. Rates are lower among men of providing detailed, high-definition images Asian ethnicity and higher in men of Black to detect abnormalities that could be ethnicity. Family history is also important and associated with prostate disease or cancer

population and up to sevenfold in men under 65. There is no association between BRCA1 gene and prostate cancer.

52,300 new cases a year

nd


Bowel cancer

Bowel cancer is the fourth most common cancer. and the second leading cause of cancer death in the UK. While rates have remained stable overall, incidence in younger age groups has increased over the past 30 years. Mortality has fallen by 11% in the last decade, largely due to earlier detection.

Faecal Immunochemical

most common cause of

cancer deaths in males





most common most common

cause of cancer

BOWEL CANCER IS THE

cause of death

FIT is a sensitive and specific screening tool that detects hidden blood in stool, helping to identify those who may benefit

from colonoscopy. At Preventicum, our assessments include a FIT for clients over 40 or when clinically indicated. If a positive result is detected, we can arrange prompt referrals to our trusted network of colorectal specialists.

Test (FIT)


For clients who may benefit from further investigation, we offer a dedicated screening colonoscopy pathway with Dr Sean Preston, leading Gastroenterologist and Director of Endoscopy at The Princess Grace Hospital.

Cancers such as leukaemia and lymphoma can be difficult to detect due to the nonspecific nature of presenting symptoms. Our Doctors combine a detailed medical history, thorough physical examination, extensive blood tests and advanced imaging to identify lymph node or spleen changes, significantly

improving the detection of these cancers.


Testicular cancer rates have increased by 27% since the early 1990s, with the highest

Haematological



ultrasound, considered the gold standard for detecting abnormalities, allowing us to view small or hidden lumps that may not be felt during a physical examination.

Other cancers Our detailed MRI and ultrasound imaging of



Our assessments include a dedicated thyroid ultrasound to identify any potentially concerning lesions. For clients who may benefit from further

the abdomen and pelvis allow us to assess the health of organs such as the pancreas, ovaries, uterus, liver, kidneys and bladder, detecting early changes before symptoms arise. Cervical screening can be performed during your Preventicum assessment and detects Human Papilloma Virus (HPV), which is the most significant risk factor for cervical cancer. Ideally, testing should be carried out mid-cycle for the most accurate results. If this timing does not

align with your Preventicum assessment, please

talk to your GP about cervical screening.

Thyroid




Looking ahead

There is growing interest in Multi Cancer Early Detection (MCED) tests, which aim to identify multiple cancers from a single blood sample. While research remains in its early stages, Preventicum continues to closely monitor and appraise the latest developments in this field.

We remain committed to integrating the

Preventicum





References: Cancer Research UK | Institute for Health Metrics and Evaluation (IHME) Zhao J, Xu L, Sun J, et al Global trends in incidence, death, burden and risk factors of early-onset cancer from 1990 to 2019 BMJ Oncology 2023;2:e000049. doi: 10.1136/bmjonc-2023-000049