

Finished size of Leaflet is approximately 93.5mm wide x 128.5mm deep (that is the folded leaflet size - opened out it's 187mm wide x 128.5mm deep) – THIS LEAFLET IS AT 100%

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MITE MASTER 290 INSECTICIDE

ACTIVE CONSTITUENT: 290 g/L OMETHOATE
(an anticholinesterase compound)

GROUP 1B HERBICIDE

For the control of redlegged earth mite, blue oat mite and lucerne flea on pastures, cereals, oilseed and legume crops, and bluegreen aphid and cowpea aphid on pasture legumes, lucerne, faba beans and vetch.

This Booklet is part of the Label.
READ THOROUGHLY BEFORE USING

APVMA APPROVAL No: 61682 / 0107



GROW CHOICE PTY LTD
113 Fitzroy Street
Tamworth NSW 2340
ACN 069 839 961
Tel: (02) 6766 3979

DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Pastures, Cereals, Oilseed and legume crops	Redlegged earth mite (Not Qld) Blue oat mite (Qld, NSW only)	100 mL/ha	AUTUMN/WINTER: Spray 2-5 weeks after opening rains when mites have hatched to prevent seedling damage or loss. Mite Master is rain fast within 1 hour of spraying. Residual control is less on 1-2 true leaf plants. Mite Master should be mixed with minimum tillage or direct drill sprays to kill existing mite populations.
			SPRING: Spray mite infested pastures and legumes to prevent damage, yield loss and to reduce infestation pressure in the following autumn. Mite Master may be added to pasture herbicides.
	Lucerne flea (Not Qld)		Spray on appearance and before damage occurs. If second hatching occurs a follow up spray may be required.
Pasture Mite (Byobba mite) (NSW only)		120 mL/ha	Spray on appearance and before damage occurs.
Pasture legumes, Lucerne, Faba bean, Vetch	Bluegreen aphid	100 or 200 mL/ha	Spray when aphids start to build up. Use the higher rate when faster knockdown is required.
	Cowpea aphid	200 mL/ha	Spray when aphids start to build up.
Pasture	Spotted clover aphid	100 mL/ha	Apply at first sign of aphid activity.
Barrier spraying	Redlegged earth mite (NSW, WA only)	300 mL/ha	To prevent invasion, spray bare earth outside crop as well as weeds along the fence line using a offset jet. Note: Rain will reduce the efficacy of the bare earth treatment but leave an effective barrier on the weeds.
Poppy	Redlegged earth mite (Tas only)	100 mL/ha	Spray on appearance and before damage occurs.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION

GENERAL INSTRUCTIONS

Insecticide Resistance Warning

GROUP 1B HERBICIDE

For insecticide resistance management, Mite Master 290 Insecticide is a Group 1B insecticide.

Some naturally occurring insect biotypes resistant to Mite Master 290 Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. The effectiveness of Mite Master 290 Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Grow Choice Pty Ltd accepts no liability for any losses that may result from the failure of Mite Master to control resistant insects.

Mite Master 290 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Grow Choice representative or local agricultural department agronomist.

Mixing

Mix the required quantity of Mite Master in sufficient water to give even coverage. Boom sprayers and aircraft are satisfactory for spraying Mite Master.

Compatibility

Compatible with most commonly used herbicides from direct drilling and minimum tillage to pasture topping. Ensure that Mite Master is thoroughly mixed in the spray tank before adding a herbicide. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Grow Choice Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Grow Choice cannot be responsible for the behaviour of such mixtures.

PROTECTION OF LIVESTOCK

DANGEROUS TO BEES. DO NOT SPRAY ANY PLANTS IN FLOWER WHILE BEES ARE FORAGING.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Product and spray are poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Obtain an emergency supply of atropine tablets 0.6mg. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield and impervious footwear. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water.

If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131 126) or a Doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

Material Safety Data Sheet

For further information refer to the Material Safety Data Sheet which is available from the supplier.

NOTICE TO BUYER

Grow Choice Pty. Ltd. will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.