

# Deccox®-L

Type B Medicated Feed

## Product Description

Deccox®-L (decoquinate) Type B Medicated Feed prevents coccidiosis in cattle and calves, contains 4,542 g (0.5%) of decoquinate per ton of premix, and can be incorporated into complete feed or mixed into milk replacer. It is intended for the manufacture of Type B and Type C medicated feeds only.

## Indications for Use

- For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in ruminating and non-ruminating calves, including veal calves.

## FDA Status

- Category I drug and does not require a feedmill license for manufacture of medicated feeds.

## Withdrawal Periods & Residue Warnings

- No withdrawal period is required when used according to labeling.
- Do not feed to cows producing milk for human consumption.

## Packaging

- 50-pound, multiwall paper bag with protective barrier ply.

## Storage

- Store in a cool, dry place. Keep bag tightly closed.

## Key Points

- Synthetic coccidiostat developed for coccidiosis prevention in calves.
- Inhibits mitochondrial respiration in coccidia protozoa, halting development before clinical signs appear.
- Prevention of coccidiosis minimizes gut damage, bloody diarrhea, weight loss, immune suppression and secondary infections like bovine respiratory disease.
- Low toxicity and minimal risk of overdose.
- It contains lactose providing the ability to be mixed into a milk replacer formulation.
- Can be mixed into daily ration or into milk replacer formulation.

See reverse side for complete product use directions, claims and additional information on this product.

## Beef and Dairy Information: Deccox®-L — Type B Medicated Feed

<b>Active Drug Ingredient</b>	Decoquinate 0.5% (4,542 g/ton)		
<b>For Manufacture of Type B and Type C Medicated Feeds for Cattle</b>			
<b>Indications for Use</b>	For the prevention of coccidiosis in ruminating and non-ruminating calves (including veal calves) and cattle caused by <i>Escherichia coli</i> and <i>E. zuernii</i> .		
<b>Mixing Directions</b>	Mix 5.7 to 40 lbs. DECCOX-L with non-medicated feed ingredients to manufacture one ton of complete feed or milk replacer containing 12.9 to 90.8 grams decoquinate as in the examples below.		
	Pounds Deccox-L per ton of Type C	Pounds non-medicated feed per ton of Type C	Type C Decoquinate concentration (g/ton)
	5.7	1,994.3	12.9
	20	1,980	45.4
	40	1,960	90.8
<b>WITHDRAWAL PERIODS AND RESIDUE WARNINGS</b> - No withdrawal period is required when used according to labeling. Do not feed to cows producing milk for human consumption.			
<b>Caution</b> - Do not use in feeds containing bentonite.			

**Storage:** Store in a cool, dry place. Keep bag tightly closed.

Restricted Drug (California) — Use only as directed.

Approved by FDA under NADA # 039-417

**To learn more about Deccox-L  
talk with a Phibro expert at**

800-677-4623 or visit [www.pahc.com](http://www.pahc.com)