





## HEROIN BLACK TAR IDENTIFICATION




### Substance Test








#### Read Before Use



As a precaution, gloves and eye protection should be worn when handling drug substances and using test units. Do not puncture any part of unit. Do not remove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.

Test Preparation	1	Remove the probe from test unit.	Sample suspected substance.		
			2	Touch or Swipe or Dig	
			Touch probe to the suspected substance.	Swipe either side of the probe across the suspected substance.	Firmly press probe on the tablet while scrubbing in a circular motion until the tip is completely covered with substance.
					

Test Preparation	3	Secure probe into test unit. Press firmly.	Identification	4	From both sides of the kit, squeeze on the circle of <b>Chamber 1</b> until the capsule breaks.	5	Gently shake test unit for 2-3 seconds.
							
		<b>STOP</b> Do not continue unless probe is secured properly in place.					

Results		The appearance of a purple color indicates the presence of an opiate such as <b>Heroin, Morphine</b> or <b>Codeine</b> .
		If no purple color develops, an opiate, such as Heroin, Morphine or Codeine, is <b>not</b> present.
		<b>Proceed to Dilution (steps 6-9), regardless of whether or not a color appeared.</b>

Dilution	6	From both sides of the kit, squeeze on the circle of <b>Chamber 2</b> until the capsule breaks.	7	Gently shake test unit for 2-3 seconds.
				
				<b>Wait 5 minutes.</b> The test window will warm up. This is normal.

Dilution	8	From both sides of the kit, squeeze on the circle of <b>Chamber 3</b> until the capsule breaks.	9	Gently shake test unit for 2-3 seconds.
				
				The test window will warm up. This is normal.



#### Disposal

After Dilution, the test kit is safe to dispose of in the trash.

01-19