














Read Before Use

As a precaution, gloves and eye protection should be worn when handling drug substances and using test units. Do not puncture any part of unit. Do not remove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.

Test Preparation	1	Remove the probe from test unit.	2 Sample suspected substance.		
			Touch or	Swipe or	Dig
		Touch probe to the suspected substance.		Swipe either side of the probe across the suspected substance.	Firmly press probe on the tablet while scrubbing in a circular motion until the tip is completely covered with substance.
					

Test Preparation	3	Secure probe into test unit. Press firmly.	Identification	4	From both sides of the kit, squeeze on the circle of Chamber 1 until the capsule breaks.	5	Gently shake test unit for 20 seconds.
							
		STOP Do not continue unless probe is secured properly in place.					

Results		The appearance of a red-purple (wine) color indicates the presence of Flunitrazepam .
		The appearance of a yellow color indicates the possible presence of Nitrazepam , Clonazepam or Nimetazepam .
		The appearance of a green color indicates the possible presence of Lorazepam or Temazepam .
		If none of the above colors develop, the above listed substances are <u>not</u> present.



Disposal

The test kit is safe to dispose of in the trash.

01-19