




### Read Before Use

As a precaution, gloves and eye protection should be worn when handling drug substances and using test units. Do not puncture any part of unit. Do not remove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.

### Test Preparation

**1** Remove the probe from test unit.



**2** Sample suspected substance.

Touch or Swipe


Touch probe to the suspected substance.



Swipe either side of the probe across the suspected substance.



**3** Secure probe into test unit. Press firmly.



STOP


Do not continue unless probe is secured properly in place.

### Identification

**4** From both sides of the kit, squeeze on the circle of **Chamber 1** until the capsule breaks.




**5** Gently shake test unit for 10 seconds.




### Identification

**6** From both sides of the kit, squeeze on the circle of **Chamber 2** until the capsule breaks.



**7** Gently shake test unit for 10 seconds.



### Results



The appearance of blue color particles adhering to the sampler tip or precipitating out of solution indicates the presence of **PCP** (Phencyclidine).

\* Note: The liquid will remain dark pink.



If no blue color develops, PCP is **not** present.



### Disposal

The test kit is safe to dispose of in the trash.

01-19