



# RuralVet

ANIMAL HEALTH SOLUTIONS

## Tick, Worm & Fly

2025/2026

Spring Summer Catalogue



# tick, worm + fly



## CATTLEGUARD POUR ON 15L

Active: Moxidectin - alternative to Cydectin pour on

Indications: For control of ticks, flies and worms

WHP: Nil Milk, Nil Meat

Dose: 10ml/100kg

**\$930**

**2 or more \$900**



## MECTEC LOW VOLUME POUR ON 10L

Active: Ivermectin

Indications: For control of ticks, flies and worms

WHP: Nil Milk, Meat 42 days

Dose: 5ml/100kg

**\$450**

**2 or more \$400**



## INCARCERATE POUR ON 20L

Active: Deltamethrin – alternative to Easy Dose pour on

Indications: For control of flies

WHP: Nil Milk, Nil Meat

Dose: 5ml/100kg

**\$650**

**2 or more \$600**



## NEOVEMOX LONG ACTING INJECTION 500ML

Active: Moxidectin – alternative to Moxistar LA & Cydectin LA

Indications: For extended control of ticks and worms in heifers

WHP: Milk 80 days, Meat 56 days

Dose: 1ml/100kg

**\$600**



## DECTOMAX V INJECTABLE

Active: Doramectin and Levamisole (New combined active formula)

Indications: For control of ticks and worms in heifers

WHP: Milk 60 days, Meat 35 days

Dose: 2ml/50kg

**500ml \$145**

**3L Victory Pk + Gun \$840**



## EXITRAZ WP CATTLE DIP / SPRAY 5x500GM

Active: Amitraz – alternative to Amitik WP Soluble

Indications: For control of ticks

WHP: Nil Milk, Nil Meat

Dose: 1 x 500g sachet / 1000L of water

**\$240**

# tick, worm + fly



## EXITRAZ EC CATTLE SPRAY 2L

Active: Amitraz – alternative to Amitik EC and Taktic EC

Indications: For control of ticks

WHP: Nil Milk, Nil Meat

Dose: Spray 400ml / 200L of water

**\$110**



## BACKRUB SELF-FEEDING

Indications: Self Feeding Backrub for control of flies.

Can be connected to ready to use backrub products such as Cattle Coat

**2M \$175**

**3M \$210**



## CATTLE-COAT RTU ORGANIC BACKRUB 20L

Active: Essential Oils and Organic Extracts

Indications: For control of flies by backrub

WHP: Nil Milk, Nil Meat

Dose: Ready to Use Backrub Oil

**\$270**



## ACTI-TRACE MINERAL INJECTION 500ML

Active: Zinc, Copper, Manganese, Selenium - alternative to Multimin

Indications: Boosts essential trace minerals in high demand periods

WHP: Nil Milk, Nil Meat

Dose: Up to 1yr: 1ml/50kg, 1-2yrs: 1ml/75kg,

Over 2yrs 1ml/100kg



**\$350**



## VITAMIN ADE 500ML

Active: Vitamin A, D3 & E

Indications: For treatment and prevention of Vitamin A, D & E deficiency

WHP: Nil Milk, Nil Meat

Dose: Cattle 5ml, Calves 1ml first week after birth

**\$184**



## COBALIFE VITAMIN B12 +/- Se INJECTION 500ML

Active: Vitamin B12 with or without Selenium

Indications: For treatment and prevention of VB12 & Selenium deficiency

WHP: Nil Milk, Nil Meat

Dose: Cattle 4-6ml pre calving,

Calves 2-3ml from 2mnths

**B12 \$70**

**B12 + Se \$83**

# tick, worm + fly



## CYLENCE ULTRA EAR TAG 20PK

Active: Piperonyl Butoxide and Betacyfluthrin

Class: Synthetic Pyrethroid (SP)

Indications: For control of flies for up to 4mths

WHP: Nil Milk, Nil Meat

Dose: 2 tags per animal (1 tag per ear)

**\$80**



## TRIFECTA 10L

Active: Levamisol & Oxfendazole & Abamectin

Indications: Triple active oral treatment for control of intestinal worms

WHP: Milk Do not use less than 28 days before calving, Meat 21 days

Dose: 1ml per 10kg liveweight

**\$837**



## COXIDOX 50

Active: Toltrazuril – alternative to Baycox & Coxi-stop

Indications: Oral treatment for control of coccidiosis in calves

WHP: Milk do not use in lactating cows, Meat 56 days

Dose: 3ml per 10kg liveweight once

**LOW PRICE**  
**\$195**



## MECTEC PLUS 500ML INJECTABLE

Active: Ivermectin & Clorsulon - alternative to Ivomec Plus & Baymec Gold

Indications: Ivermectin sensitive intestinal worms and adult liver fluke

WHP: Nil Milk, Meat 28 days

Dose: 1ml/50kg

**\$170**



## NILZAN LV ORAL DRENCH - 2L & 5L

Active: Levamisole and Oxyclozanide

Indications: Levamisole sensitive intestinal worms and adult liver fluke

WHP: Nil Milk, Meat 14 days

Dose: 1ml/10kg

**2L \$406**

**5L \$834**



## EXIFLUKE 240 ORAL DRENCH - 2.5L & 10L

Active: Triclabendazole

Indications: For control of all stages of liver fluke

WHP: Do not use less than 21 days before calving,

Meat 14 days

Dose: 2.5ml/50kg

**2.5L \$364**

**10L \$930**