

CR134 – microhelix, tRNA acceptor-stem mimic for flexizyme acylation (22 nucleotides)

GGCUCUGUUCGCAGAGCCGCCA

References -

1) Sequence source: Goto, Y., Katoh, T., & Suga, H. (2011). Flexizymes for genetic code reprogramming. In *Nature Protocols* (Vol. 6, Issue 6, pp. 779-790). Springer Science and Business Media LLC. <https://doi.org/10.1038/nprot.2011.331>

2) PDB structure file: 3CUN



3) Xiao, H., Murakami, H., Suga, H., & Ferre-D'Amare, A. R. (2008). Structural basis of specific tRNA aminoacylation by a small in vitro selected ribozyme. In *Nature* (Vol. 454, Issue 7202, pp. 358-361). Springer Science and Business Media LLC. <https://doi.org/10.1038/nature07033>

4) Murakami, H., Ohta, A., Ashigai, H., & Suga, H. (2006). A highly flexible tRNA acylation method for non-natural polypeptide synthesis. In *Nature Methods* (Vol. 3, Issue 5, pp. 357-359). Springer Science and Business Media LLC. <https://doi.org/10.1038/nmeth877>

5) Zhang, J., & Ferre-D'Amare, A. R. (2014). A Flexible, Scalable Method for Preparation of Homogeneous Aminoacylated tRNAs. In *Methods in Enzymology* (pp. 105-113). Elsevier. <https://doi.org/10.1016/b978-0-12-801122-5.00005-2>