

# Are You Struggling From Chronic Low Back Pain?


For lumbar spinal stenosis patients, the minimally invasive *mild*<sup>®</sup> Procedure may provide lasting pain relief\* and increased mobility<sup>1</sup>.

\*as demonstrated in a 2-year follow up study


# Lumbar Spinal Stenosis (LSS)



# Lumbar Spinal Stenosis is typically a Degenerative Condition



**HEALTHY  
SPINE**




**AGING SPINE WITH LSS**

Aging and natural wear and tear on the spine can cause conditions that put pressure on the spinal nerves<sup>2</sup>

# Lumbar Spinal Stenosis Impacts a Large Population

**~20%**  
of people over  
60 have LSS<sup>3</sup>



**2M**   
U.S. patients in  
active treatment<sup>4</sup>

# Common Symptoms of Lumbar Spinal Stenosis




LSS patients may experience  
**PAIN, NUMBNESS,**  
tingling, or heaviness in back,  
legs, or buttocks when  
standing/walking



Symptoms may be  
**RELIEVED**  
by sitting, bending forward,  
or sleeping curled in the  
fetal position

# A Major Root Cause of Lumbar Spinal Stenosis




**Up to 85% of spinal canal narrowing** is caused by thickened ligament called the ligamentum flavum<sup>5</sup>  
**This excess ligament causes pressure on the nerves in the lower back.<sup>2</sup>**

# Minimally Invasive Lumbar Decompression

*mild*<sup>®</sup>

# Minimally Invasive Outpatient Procedure

## *mild* Addresses a Major Root Cause of LSS



Incision size of 5.1 mm<sup>8</sup>  
No implants left behind, *only a bandage*



Outpatient procedure



Safety profile similar to an ESI<sup>6</sup>




Can be performed using local anesthetic and light sedation



Patients are typically discharged within 24 hours<sup>7</sup>

# National Medicare Coverage



# Proven Outcomes



# Lasting Relief with *mild*

A similar safety profile to an ESI<sup>6</sup>, with statistically superior relief.<sup>1</sup>

5-Year Durability<sup>9</sup>  
**88%** of patients avoided  
surgical decompression  
for at least 5 years



# Significant Functional Improvement after *mild*

Cleveland Clinic Study, 1-Year Study<sup>10</sup>



Mean **Standing Time**

**600%**

Improvement



# Significant Functional Improvement after *mild*

Cleveland Clinic Study, 1-Year Study<sup>10</sup>



Mean **Walking Distance**

Over

**1,500%**

Improvement



# High Patient Satisfaction with *mild*

Studies show that *mild* continues to improve patient functionality over time, including increased mobility and reduction of pain<sup>1</sup>

## Patient Satisfaction Rate

**76.8%** MiDAS ENCORE  
2-Year Study<sup>1</sup>



# References

1. Staats PS, Chafin TB, Golovac S, Kim CK, Li S, Richardson WB, Vallejo R, Wahezi SE, Washabaugh EP 3rd, Benyamin RM; MiDAS ENCORE Investigators. Long-Term Safety and Efficacy of Minimally Invasive Lumbar Decompression Procedure for the Treatment of Lumbar Spinal Stenosis With Neurogenic Claudication: 2-Year Results of MiDAS ENCORE. *Reg Anesth Pain Med*. 2018 Oct;43(7):789-794. doi: 10.1097/AAP.0000000000000868. PMID: 30199512; PMCID: PMC6319572. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6319572/>
2. Wu L, Munakomi S, Cruz R. Lumbar Spinal Stenosis. [Updated 2024 Jan 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK531493/>.
3. Kalichman L, Cole R, Kim DH, et al. Spinal stenosis prevalence & association with symptoms: The Framingham Study. *Spine J*. 2009;9(7):545-550. doi:10.1016/j.spinee.2009.03.005.
4. All-payer claims analysis
5. Hansson T, Suzuki N, Hebelka H, Gaulitz A. The narrowing of the lumbar spinal canal during loaded MRI: the effects of the disc and ligamentum flavum. *Eur Spine J*. 2009;18(5):679-686. doi:10.1007/s00586-009-0919-7
6. Benyamin RM, Staats PS, MiDAS ENCORE Investigators. mild® is an effective treatment for lumbar spinal stenosis with neurogenic claudication: MiDAS ENCORE Randomized Controlled Trial. *Pain Physician*. 2016;19(4):229-242.
7. Deer T, Kim C, Wahezi SE, Qu H, Sayed D; MOTION Study Investigators. Objective Real-World Outcomes of Patients Suffering from Painful Neurogenic Claudication Treated with the mild® Procedure: Interim 6-Month Report of a Randomized Controlled Trial. *J Pain Res*. 2021 Jun 10;14:1687-1697.
8. Stryker Data on File
9. Mekhail, N., Costandi, S., Nageeb, G., Ekladios, C., & Saied, O. (2021). The durability of minimally invasive lumbar decompression procedure in patients with symptomatic lumbar spinal stenosis: Long-term follow-up. *Pain practice : the official journal of World Institute of Pain*, 10.1111/papr.13020. Advance online publication. <https://doi.org/10.1111/papr.13020>
10. Mekhail N, Costandi S, Abraham B, Samuel SW. Functional and patient-reported outcomes in symptomatic lumbar spinal stenosis following percutaneous decompression. *Pain Pract*. 2012;12(6):417-425. doi:10.1111/j.1533-2500.2012.00565.x.

# Disclaimer

Only your doctor can make the medical judgment on which products and treatments are right for our own individual condition. Your physician will explain all the possible complications of the procedure, as well as side effects. Individual results vary and not all patients will receive the same post-procedure activity level.

The information presented is for educational purposes only. Stryker is not dispensing medical advice. The information presented is intended to demonstrate a Stryker product. A surgeon must always refer to the package insert, product label and/or instructions for use, including the instructions for cleaning and sterilization (if applicable), before using any Stryker product. Stryker or its affiliated entities own, use, or have applied for the following trademarks or service marks: mild and Stryker. All other trademarks are trademarks of their respective owners or holders.

The absence of a product, feature, or service name, or logo from this list does not constitute a waiver of Stryker's trademark or other intellectual property rights concerning that name or logo.