

AUDIT

Piccinelli (1997) Efficacy of the alcohol use disorders identification test as a screening tool for hazardous alcohol intake and related disorders in primary care: a validity study
 PMCID: PMC2125904 <https://doi.org/10.1136/bmj.314.7078.420>

For all questions, one drink equals...



How often do you have a drink containing alcohol?

- Never
- Monthly or less
- 2 to 4 times a month
- 2 to 3 times a week
- 4 or more times a week

How many drinks containing alcohol do you have on a typical day when you are drinking?

- 1 or 2
- 3 or 4
- 5 or 6
- 7, 8, or 9
- 10 or more

How often do you have six or more drinks on one occasion?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

How often during the last year have you found that you were not able to stop drinking once you had started?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

How often during the last year have you failed to do what was normally expected from you because of drinking?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

How often during the last year have you had a feeling of guilt or remorse after drinking?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

How often during the last year have you been unable to remember what happened the night before because you had been drinking?

- Never
- Less than monthly
- Monthly
- Weekly
- Daily or almost daily

Have you or someone else been injured as a result of your drinking?

- No
- Yes, but not in the last year
- Yes, during the last year

Has a relative, friend, doctor, or another health worker been concerned about your drinking or suggested you cut down?

- No
- Yes, but not in the last year
- Yes, during the last year

Total score _____