

Spinal Anaesthetic

A05 Lite - Expires end of January 2026



This fact sheet is for general information about this procedure only. It is not intended to be used as medical advice or to replace advice that your relevant healthcare professional would give you. If you have a particular medical problem, please consult a healthcare professional.

© EIDO Systems International Limited. The operation and treatment information in this document is published under license from EIDO Systems International and is protected by copyright laws. Other than for your personal, non-commercial use, you may not copy, print out, download or otherwise reproduce any of the information.

eidohealthcare.com



What is a spinal anaesthetic?

A spinal anaesthetic (or spinal) involves injecting local anaesthetics and other painkillers into the subarachnoid space (an area filled with fluid near your spinal cord). This numbs your nerves to give pain relief in certain areas of your body. A spinal can be used either on its own while you are awake, or together with sedation or a general anaesthetic. A spinal can also be used after a procedure to give effective pain relief.

How does a spinal work?

The subarachnoid space is the bag of fluid that surrounds your spinal cord and the nerves that come out from it. Local anaesthetics and other painkillers are injected using a fine needle into this space.

How is a spinal given?

To insert the needle, your anaesthetist will ask you to either sit up or lie on your side.

Your anaesthetist will insert the needle and when they are certain that it is in the right position they will inject anaesthetic through it.

It should not be painful, although it can be uncomfortable.

The time that the spinal lasts for varies but is usually 1 to 3 hours. Your anaesthetist will put enough anaesthetic through the needle to make sure that it lasts longer than the expected length of the procedure.

What complications can happen?

- The spinal not working properly
- Low blood pressure
- Headache
- Shivering after your procedure
- Allergic reaction to the equipment, materials or medication
- Itching
- Difficulty passing urine
- Backache
- Cardiovascular collapse (where your heart stops)

- Unexpected high block, if the local anaesthetic spreads beyond the intended area
- Infection around your spine
- Nerve damage
- Various other more serious problems have been reported with spinals, including spinal abscess and blood clots
- Paralysis or death

A complication may happen after you have had a spinal.

- Pus, redness, tenderness or pain
- A high temperature
- Feeling unwell
- Discomfort when in a bright room or sunlight
- Neck stiffness
- Difficulty moving or feeling your legs
- Difficulty passing urine
- Bowel incontinence

What happens after the procedure?

The effect of the spinal will wear off after a few hours.

Summary

A spinal anaesthetic can be used for most people, usually giving a safe and effective form of pain relief both during and after a procedure.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewer

Rakesh Kapila (MBChB, FRCA)

Illustrator

Medical Illustration Copyright © Nucleus Medical Art. All rights reserved. www.nucleusinc.com