

Certificate of Analysis

KAT2A (GCN5L2)

Human histone acetyltransferase KAT2A

(Recombinant enzyme expressed in *E.coli*)

Item # EPI024

Lot # 139716

Product Description: Recombinant human KAT2A (GCN5L2), amino acids 497-662, expressed in *E.coli*. Purified using immobilised metal affinity chromatography.

MW = 19.3kDa.

Tag cleaved by thrombin protease.

Aliases: PCAF-b

Formulation: 1mg/ml of enzyme in 5mM Tris/HCl pH7.4, 250mM NaCl, 50% Glycerol. Frozen solution.


Storage and Stability: Stable for 1 year at -70°C from date of shipment. For maximum recovery of product, centrifuge original vial prior to removing the cap.

Handling Recommendations: Rapidly thaw the vial under cold water and immediately place on ice. Aliquot unused material into pre-chilled microcentrifuge tubes and store at -70°C.


**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR ANIMALS**

Quality Control Testing

HAT Assay: 0.125-4ng of this lot of enzyme transferred acetyl groups from [³H] Acetyl CoA to histone H3 full length in the assay described on page two. Assay background was subtracted from the actual counts to yield the results shown below.



MS: Size was confirmed by mass spectrometry using a Q-TOF.



SDS-PAGE and Coomassie Stain: Purity was assessed by SDS-PAGE and Coomassie blue staining using 4µg of KAT2A (GCN5L2).

Certificate of Analysis

KAT2A (GCN5L2) Assay Protocol

Stock Solutions:

1. **Reaction buffer:** 50mM Hepes/Tris pH7.4, 0.1mM EDTA/Tris, 0.01% Triton.
2. **KAT2A (GCN5L2):** Dilute with reaction buffer. Use 0.125-4ng per assay point.
3. **Histone H3 full length:** Dilute with reaction buffer to 2500nM.
4. **[3H] Acetyl CoA:** Dilute with reaction buffer to 150nM.
5. **Filtration Buffer :** 33mM Citric acid pH2.2

Assay Procedure (96 well plate format):

1. Add 5µl of 10% DMSO per assay to each well.
2. Add 25µl of [3H] Acetyl CoA.
3. Add 10µl (**0.125-4ng**) **KAT2A (GCN5L2)**.
4. Add 10µl of Histone H3 full length.
5. Incubate for 10 minutes at 22°C.
6. Stop the reaction by filtration on a GF/B Filter. Wash 3 times with filtration buffer.
7. Dry and add scintillation cocktail.
8. Read in a scintillation counter. Compare the signal of enzyme samples with that of a background sample that contains all assay components except the substrate Histone H3 full length.

Certificate of Analysis

KAT2A (GCN5L2) Sequence Information

Protein Human KAT2A (GCN5L2)
Tags *tag cleaved by thrombin protease*
Accession number GenBank NP_066564.2

Recombinant KAT2A (GCN5L2) amino acid sequence:

1 GSGIIEFHVI GNSLTPKANR RVLLWLVGLQ NVFSHQLPRM PKEYIARLVF
51 DPKHKTLALI KDGRVIGGIC FRMFPTQGFT EIVFCAVTSN EQVKGYGTHL
101 MNHLKEYHIK HNILYFLTYA DEYAIGYFCK QGFSKDIKVP KSRYLGYIKD
151 YEGATLMECE LNPRIPTY

Reviewed and approved by site quality representative.

Unless otherwise stated in our catalogue or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

© 2014 **Eurofins Pharma Discovery Services UK Limited** is an independent member of Eurofins Discovery Services.