

Certificate of Analysis

sirtuin 1

Human NAD-dependent protein deacetylase sirtuin-1, isoform a
(Recombinant protein expressed in *E.coli*)

Item # EPI059

Lot # 212439

Product Description: N-terminal 6His-tagged, recombinant, amino acids 193-741, human sirtuin 1, expressed in *E.coli*. Purified using immobilised metal affinity chromatography.

MW = 63.8kDa.

Alias: SIRT1

Formulation: 0.78mg/ml of protein in 25mM Hepes/NaOH pH 7.4, 125mM NaCl, 50% glycerol. Frozen solution.


Storage and Stability: On receipt of material store at -70°C. Unopened reagent is stable for a minimum of 1 year from date of shipment when stored at recommended storage temperature. Avoid repeat freeze/thaw cycles. For maximum recovery of product, centrifuge original vial prior to removing the cap.

Handling Recommendations: Rapidly thaw the vial under cold water and immediately place on ice. Aliquot unused material into pre-chilled micro-centrifuge tubes and immediately snap-freeze the vials in liquid nitrogen prior to re-storage at -70°C.


**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR ANIMALS**

Quality Control Testing

Enzymatic Assay: 62.5-1000ng of this lot of enzyme deacetylated 200µM Fluorogenic HDAC substrate in the assay described on page two. Assay background was subtracted from the actual counts to yield the results shown below.



MS: Size was confirmed by mass spectrometry using a Q-TOF.



Certificate of Analysis

Sirtuin 1 Assay Protocol

Stock Solutions:

1. **Reaction buffer:** 50mM Tris/HCl 8, 137mM NaCl, 2.7mM KCl, 1mM MgCl₂, 0.2% BSA.
2. **Sirtuin 1:** Dilute with reaction buffer. Use 62.5-1000ng per assay point.
3. **Mix β -NAD / Fluorogenic HDAC Substrate:** Dilute with reaction buffer to 1mM and 400 μ M respectively.

Assay Procedure (384 well plate format):

1. Add 2 μ l of H₂O per assay to each well.
2. Add 8 μ l (**62.5-1000ng**) **sirtuin 1**.
3. Incubate for 5 minutes at 22°C.
4. Add 10 μ l of Fluorogenic HDAC substrate/ β -NAD.
5. Incubate for 20 minutes at 22°C.
6. Add 20 μ l of HDAC Developer.
7. Incubate for 15 minutes at 22°C.
8. Read at 460nm after an excitation flash at 355nm. Compare the signal of enzyme samples with that of a background sample that contains all assay components except the enzyme sirtuin 1.

Certificate of Analysis

Sirtuin 1 Sequence Information

Protein Human sirtuin 1

Tags N-terminal 6His

Accession number GenBank NP_036370.2

Recombinant Sirtuin 1 amino acid sequence:

```
1 MHHHHHSSG VDLGTENLYF QSMIGTDPRT ILKDLLPETI PPPELDDMTL
51 WQIVINILSE PPKRKKRKDI NTIEDAVKLL QECKKIIVLT GAGVSVSCGI
101 PDFRSRDGIY ARLAVDFPDL PDPQAMFDIE YFRKDPRPFF KFAKEIYPGQ
151 FQPSLCHKFI ALSDEKGLL RNYTQNIDTL EQVAGIQRII QCHGSFATAS
201 CLICKYKVC EAVRGDIFNQ VVPRCPRCPA DEPLAIMKPE IVFFGENLPE
251 QFHRAMKYDK DEVDLLIVIG SSLKVRPVAL IPSSIPHEVP QILINREPLP
301 HLHFDVELLG DCDVIINELC HRLGGEYAKL CCNPVKLSEI TEKPPRTQKE
351 LAYLSELPPT PLHVEDSSS PERTSPDSS VIVTLDDQAA KSNDDLVDSE
401 SKGCMEEKPQ EVQTSRNVES IAEQMPNPD LKNVGSSTGEK NERTSVAGTV
451 RKCWPNRVAK EQISRRLDGN QYLFLPPNRY IFHGAEVYSD SEDDLSSSS
501 CGSNSDSGTC QSPSLEEPME DESEIEEFYN GLEDEPDVPE RAGGAGFGTD
551 GDDQEAINEA ISVKQEVTDN N
```

Reviewed and approved by site quality representative.

Unless otherwise stated in our catalogue or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.