



associates in
F·A·M·I·L·Y
DENTISTRY

AFTER DENTAL IMPLANT SURGERY

BLEEDING: Slight bleeding after surgery is not unusual. Your saliva may be tinged with blood for up to twenty-four (24) hours. In some situations, gauze will be placed over the site after surgery, leaving in place over the surgery site for at least two (2) hours maintaining firm biting pressure. Do not forcefully spit or irritate the surgery site. This may cause further bleeding. If bleeding persists, it may be controlled with pressure (moisten a clean gauze and place directly over the surgery site. Apply moderate biting pressure for approximately ½ hour. Repeat if necessary).

PAIN: It is not uncommon to have discomfort and pain for the first few days after surgery. Take pain medications as needed. The first day, it is recommended to take medication even with minimal pain as it is easier to prevent pain than to decrease it. Pain medications are most effective when taken before local anesthesia diminishes and normal sensation returns. Do not take pain medication on an empty stomach.

SWELLING: After surgery, swelling around the mouth, cheeks, eyes, and side of the face is very common. This is the body's normal reaction to surgery and eventual repair. Swelling usually will not become apparent until the day following surgery and will not reach its maximum until 2-3 days post-operatively. However, the swelling may be minimized by the immediate use of ice packs. Apply an ice pack to the outside of your face over the surgery site for twenty (20) minutes on, twenty (20) minutes off for the first twenty-four (24) hours, moist heat can be applied to the area as ice will increase swelling. Sleeping with your head elevated above the level of the heart for the first 2-3 post operative nights tends to lessen swelling.

BRUISING: Bruising is also very common after surgery. Bruising may be present over the surgery site and various areas of the face and neck. The development of black, blue, green, or yellow discoloration is due to blood spreading beneath the tissues. This is a normal post-operative occurrence, which may occur 2-3 days post-operatively and last up to 14 days. Moist heat applied to the area may speed up the removal of the discoloration.

NAUSEA: Avoid taking any medications on an empty stomach. Take all medications with a small amount of food. If you feel nauseated, drink clear liquids such as ginger ale, 7UP, apple juice, broth or try eating toast and crackers.

DIET: After surgery, drink lots of clear liquids and eat soft, nutritious foods. Avoid acidic foods such as tomatoes, orange juice, and citrus fruits until the surgery site has healed (2weeks). Do not drink through a straw for at least 48 hours. If you had surgery on only one side of your mouth, favor the other side while chewing for the first few days.

ORAL HYGIENE: Hygiene should not be neglected as an accumulation of food and debris may promote inflammation and/or infection. You may rinse and brush your teeth starting the day after surgery. Brush your teeth as usual, do not brush the surgical area for 4-5 days. Chlorhexidine mouthrinse can also be used as a mouthrinse with non-vigorous rinsing. A prescription for Chlorhexidine will be provided if necessary.

ACTIVITY: Limit strenuous activity for the first 2-3 days after surgery. This will reduce bleeding and minimize swelling. Strenuous work or exercise may promote bleeding and increase swelling which may be damaging to the surgery site and delay healing.

FEVER: After surgery, it is normal for the body temperature to be slightly elevated for 24 hours. Please remember to drink sufficient amounts of clear liquids to keep your body hydrated. Contact the office if there is a high temperature or if a low-grade temperature persists after the first 24 hours.

SMOKING/ALCOHOL: Smoking and alcohol use will delay healing and predispose you to infection which can result in the failure of the implant or bone graft. This will result in possible surgical intervention, delay in treatment, and possible loss of implant/bone graft. **Do not smoke, vape, use chewing tobacco or alcohol until the surgical site is completely healed (~ 2 weeks).**

SUTURES: Sutures may be placed in the area of surgery to minimize post-operative bleeding and to help healing. Sometimes they may become dislodged, this is no cause for alarm. In most cases these sutures are dissolvable and should disappear on their own in approximately 7-14 days. If they are still present, our office will remove them on your post-op appointment.

MEDICATIONS: It is very important that you take all your prescribed medications from your physician as prescribed. DO NOT DISCONTINUE ANY MEDICATIONS (especially blood thinners) UNLESS INSTRUCTED BY YOUR PHYSICIAN. Additionally, take all medications that our office prescribes for you. Each medication has a specific role in the post-operative period of your surgery, which will decrease the possibility of infection, swelling, and pain. Discontinue the antibiotic medication if a rash develops or if an upset stomach persists. **Report this immediately to our office.** **Women please note:** if you are currently taking birth control pills, be aware that taking antibiotics will decrease the effectiveness of the birth control pills and alternative methods should be utilized.

WEARING YOUR PROSTHESIS: Follow the specific doctor's instructions on the use of partial dentures, flippers, or full dentures after surgery. Excessive pressure on the surgical site may lead to complications or failure of the dental implants or bone grafts.

NUMBNESS: Be cautious until the anesthesia wears off as you may not have feeling in the soft tissues, and you may unknowingly injure yourself. After surgery you will have prolonged numbness in the surgical area. Care should be exercised in eating or drinking any hot fluids as you may cause injuries to the soft tissues. Ideally, eat and drink on the opposite side until the numbness subsides. If numbness persists after 24 hours, please contact our office.

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will assist you, but if you have questions about your progress, please call the office at [231-946-9644](tel:231-946-9644)