

Patient Information		
Patient Name:		Order Date:
DOB:	Ht:	Wt:
Allergies:		
Diagnosis:		ICD-10 Code:
Patient Status: ☐ New to Therapy ☐ Continuing Therapy		Next Treatment Date:
Lab Orders		Pre-Medication Orders
☐ CBC w/diff and Platelets ☐ CMP ☐ Immunoglobulin (IgG) level ☐ Pregnancy Test ☐ Draw labs during infusion ☐ Other: _____ Frequency: ☐ Baseline ☐ Every 3-6 months ☐ Other:		☐ Methylprednisolone 125mg IVP over 5 min; Administer 30 min prior to infusion Diphenhydramine PO 25 mg ☐ or 50 mg ☐ PO 30-60 min prior; may repeat x1 in 4 hours Diphenhydramine IVP 25 mg ☐ or 50 mg ☐ IVP 30-60 min prior; may repeat x1 in 4 hours Cetirizine ☐ 10 mg PO or ☐ 10 mg IVP 30 minutes prior ☐ Famotidine 20 mg PO 30 minutes prior ☐ Dexamethasone 20 mg PO 30 minutes prior Acetaminophen ☐ 650 mg or ☐ 1000 mg PO 30 minutes prior; may repeat x1 in 4 hours
Other Orders & Notes		
☐ IV to PO premedication conversion per nurse discretion after tolerance of loading doses		
Treatment Order		
☐ Imaavy (nipocalimab-aahu)	Injection (IV)	☐ Initial dose: 30 mg/kg IV over ≥30 minutes (one time), Total calculated initial dose: _____mg ☐ Maintenance dose: 15 mg/kg IV over ≥15 minutes every 2 weeks x 1 year Total calculated maintenance dose: _____mg
Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN		
Nursing		
Skilled RN to establish and maintain vascular access as needed, draw ordered labs, conduct patient assessments, and educate on infusion services, medication administration, self-monitoring and safety Imaavy Infusion Guidance: • Administer via IV infusion only; dilute per PI. • Initial dose: infuse over ≥30 minutes. • Subsequent/maintenance doses: infuse over ≥15 minutes every 2 weeks. • Monitor during infusion and for 30 minutes post-infusion for hypersensitivity or infusion-related reactions.		
Adverse Event (AE) Protocol		
Epinephrine (based on patient weight) EpiPen® Auto-Injector 0.3mg (1:1000) Inject IM or SubQ to patients who weigh ≥ 30 kg (66 lbs); may repeat in 3-5 minutes x1 if necessary. EpiPen Jr® Auto-Injector 0.15mg (1:2000) Inject IM or SubQ for patients who weigh 15 - 30 kg (33-66lbs); may repeat in 3-5 minutes x1 if necessary. Diphenhydramine 50mg (1 mL) If not previously administered, give 50mg slow IVP, or administer IM if no IV access; May repeat x1 after 10 minutes, if necessary. Solumedrol 125 mg IVP Sodium Chloride 0.9% 500mL Infuse IV at a rate of 30mL/hr. Zofran 4mg ODT or IVP for nausea Pepcid 20mg IV for urticaria/rash Acetaminophen 325-650 mg PO x1 PRN pain if not already administered OR ibuprofen 200-400mg PO prn pain		
Provider Information		
By signing this form and utilizing our services, you are authorizing Nira Medical, and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient.		
Office Contact Name:		Email Address:
Prescriber Name:		Date:
Prescriber NPI:		Phone: Fax:
Prescriber Signature:		