

THIAMETHOXAM	GROUP	4A	INSECTICIDE
LAMBDA-CYHALOTHRIN	GROUP	3A	INSECTICIDE

# Tent Insecticide CS

For indoor and outdoor use.

For Use In, On, and Around Buildings and Structures for the Control of Listed Pests, on Non-Residential Lawns and Turf, Ornamental Trees and Shrubs Around Non-Residential, Institutional, Commercial, and Industrial Buildings.

## ACTIVE INGREDIENT:

Thiamethoxam (CAS No. 153719-23-4) ..... WT. BY % 11.6%

Lambda-cyhalothrin\* (CAS No. 91465-08-6) ..... 3.5%

OTHER INGREDIENTS: ..... 84.9%

TOTAL: ..... 100.0%

\* Microencapsulated

Contains 1.04 lbs. thiamethoxam and 0.32 lb. lambda-cyhalothrin in a CS formulation.

In the State of New York, this product may only be applied for indoor uses.

## KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for First Aid, Precautionary Statements, Directions for Use, and Storage and Disposal.

EPA Reg. No. 83529-357

Manufactured For:

**Sharda USA LLC** 

7217 Lancaster Pike, Suite A  
Hockessin, Delaware 19707

EPA Est. No. **AG** 72159-GA-001; **MA** 83411-MN-001;

**MC** 89332-GA-001; **SC** 39578-TX-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

**Net Contents: 1 Quart (0.25 Gal.)**

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.	

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### **CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### **Personal Protection Equipment (PPE)**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to wildlife, highly toxic to aquatic invertebrates, and extremely toxic to fish. Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, **DO NOT** apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. When making applications, care should be used to avoid household pets, particularly fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming plants or weeds. **DO NOT** apply this product or allow it to drift to blooming plants or weeds if bees are foraging.

**NON-TARGET ORGANISM ADVISORY STATEMENT** - This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

#### Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. (See manual at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>.)

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

#### Groundwater Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### Spray Drift Advisory

**DO NOT** allow this product to drift.

## PROTECTION OF POLLINATORS



**APPLICATION RESTRICTIONS** EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon  in the **DIRECTIONS FOR USE** for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

**This product can kill bees and other insect pollinators.**

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org>

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: [www.aapco.org](http://www.aapco.org)

Pesticide incidents should also be reported to the National Pesticide Information Center at: [www.npic.orst.edu](http://www.npic.orst.edu) or direct to EPA at: [beekill@epa.gov](mailto:beekill@epa.gov).

## PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** use this product in or on electrical equipment due to the possibility of shock hazard from the water used in mixing this product.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### USE RESTRICTIONS



- **DO NOT** apply this product while bees are foraging.
  - **DO NOT** apply to plants that are flowering.
  - Only apply after all flower petals have fallen off.
- **DO NOT** apply this product, by any application method, to linden, basswood, or *Tilia* species.
  - **DO NOT** use this product with oil.
  - **DO NOT** apply this product in any room being used as a living, eating, sleeping, or recovery area by patients, the elderly, or infirm when they are in the room.
  - **DO NOT** apply to classrooms, nurseries, or daycare facilities when in use.
  - **DO NOT** apply to institutions (including libraries, sports facilities, etc.) when occupants are present in the immediate treatment area.
  - **DO NOT** apply to carpets and hard surfaces as a pin stream or coarse spray treatment.
  - **DO NOT** allow animals and people access to treated areas until the spray deposit has dried.
  - **DO NOT** apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
  - **DO NOT** treat surfaces likely to be contacted by food. **DO NOT** apply when food-handling establishments are in operation or foods are exposed. Food must be removed from area being treated. **DO NOT** apply directly to food or allow applications to contaminate food.
  - Use in all indoor-permitted sites, including food-handling establishments, must be restricted to areas that eliminate exposure to food-handling surfaces and areas that are not easily accessible to occupants.
  - **DO NOT** spray less than the labeled rate when treating for ants, fleas, flies, mosquitoes, ticks, bed bugs, scorpions, wasps, hornets, yellow jackets, carpenter bees or termites.
  - **DO NOT** allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are located in/around structures being treated.
  - **DO NOT** use water-based sprays of **Tent Insecticide CS** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.
  - **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
  - During application, **DO NOT** allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters.
  - **DO NOT** apply as broadcast spray application using a mechanically pressurized handgun in industrial and commercial areas (e.g., tires, rail, yards, junk yards).
  - For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

**All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:**

1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 2 ft. up 3 ft. out from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a surface band not to exceed one inch in width;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

**DO NOT** water treated area to the point of runoff.

**DO NOT** apply or irrigate to the point of runoff.

**DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

**DO NOT** apply the product into fish pools, ponds, streams, or lakes. **DO NOT** apply directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitats can occur. **DO NOT** allow the product to enter any drain during or after application. **DO NOT** allow to enter indoor or outdoor drains.



Follow proper disposal procedures on this label.

**Crack-and-Crevise Treatments:** Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and crack-and-crevices, applications must be made in a such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

#### **USE PRECAUTIONS**

Shake well before using.

Let treated surfaces dry before allowing humans and pets to contact surfaces.

Surface Staining: **Tent Insecticide CS** has not stained or caused damage to painted or varnished surfaces, plastics, fabrics, tile, siding or other surfaces where water applied alone causes no damage. However, treat a small area and allow it to dry to determine whether staining will occur. To avoid contamination from previous treatments, thoroughly wash out sprayer, hoses and screens with water and detergent before using **Tent Insecticide CS**.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
LAMBDA-CYHALOTHRIN	GROUP	3A	INSECTICIDE

### INSECT RESISTANCE MANAGEMENT

For resistance management, **Tent Insecticide CS** contains lambda-cyhalothrin (a Group 3A insecticide, pyrethroids, pyrethrins) and thiamethoxam (a Group 4A insecticide, neonicotinoids). Any insect population may contain individuals naturally resistant to **Tent Insecticide CS** and other Group 3A/4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Tent Insecticide CS** or other Group 3A/4A insecticides within a growing season, or among growing seasons, with different Groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode-of-action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Sharda USA LLC at <https://shardausa.com/>.

## STRUCTURAL AND PERIMETER PEST CONTROL

### Use Sites

Use **Tent Insecticide CS** as a perimeter, crack-and-crevice, and/or spot treatment in and around structures or modes of transportation. Permitted areas of use include the following buildings, structures, and modes of transport:

- aircraft (cargo and other non-cabin areas only)
- boiler rooms
- buses
- closets
- correctional facilities
- dumpsters
- factories
- fencing
- food manufacturing, processing, and serving establishments, such as bakeries, breweries, canneries, granaries, grain mills, etc.
- furniture
- garages
- garbage rooms
- greenhouses (non-commercial)
- hospitals
- hotels and motels
- industrial buildings
- laboratories
- landscape vegetation
- locker rooms
- machine rooms
- mausoleums
- mop closets
- mulch
- offices
- pet kennels
- porches
- railcars
- restaurants
- storage rooms
- schools
- stores
- trailers
- trees
- trucks
- utility passages
- vessels, vestibules
- warehouses
- wineries

Use in all indoor permitted sites, including food-handling establishments, must be restricted to areas that eliminate exposure to food-handling surfaces and areas that are not easily accessible to occupants.

For indoor applications, retreat at 21-day intervals or as necessary to maintain control. Refer to the outdoor use restrictions in the **Outdoor Pests** section.

### MIXING INSTRUCTIONS

**Tent Insecticide CS** is intended for dilution with water for application using hand-held or power-operated application equipment as a coarse spray for crack-and-crevice, spot, and surface treatments. Application equipment that delivers low-volume treatments may also be used. A surface treatment of **Tent Insecticide CS** may be applied by using a paintbrush or other porous applicator attached to a handle. Fill applicator tank with the desired volume of water and add **Tent Insecticide CS**. Close and agitate before use in order to ensure proper mixing. Re-agitate applicator tank before use if application is interrupted. Mix only amount of treatment volume as required.

Removable chemical extraction probes (also known as “stingers”) used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

### Tank Mixing

**Tent Insecticide CS** may be tank-mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, conduct a small volume mixing test with the other products. If mixed with sanitizers, add **Tent Insecticide CS** to the tank first. If other chemicals are added to the applicator tank, add **Tent Insecticide CS** last. Fill tank to desired volume and re-agitate while making applications.



It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

### **Foam Applications**

**Tent Insecticide CS** may be converted to foam and the foam used to treat structural voids to control or prevent pests including ants, wasps, wood-destroying insects, or other arthropods harboring in walls, under slabs, or in other void areas. Mix end-use dilution with manufacturer's recommended volume of foaming agent. For applications inside wall voids, expansion ratios of 15:1 or drier are suggested.

### **Fixed In-Wall Delivery Systems**

Closed, in-wall insecticidal delivery systems such as permanently installed piping or flexible tubing may also be used to deliver diluted product to inaccessible areas. Generally, about one ounce of finished liquid product is needed per 35 - 40 ft. of tubing (based on 1/8" diameter tubes). For these systems, use 0.065% - 0.13% dilution rates as listed in the **General Household Pests** table. Prepare the finished product at the appropriate dilution, then inject into system as recommended by delivery system manufacturer.

### **Spot Treatments**

Spot treatments may be made by spraying, misting, painting, or otherwise applying to surfaces covering an area no larger than 2 sq. ft. For areas limited to spot or crack-and-crevice treatments, multiple spot treatments can be made to surfaces as long as 20% of the total target area is not exceeded.

Spot treatments must not exceed 2 sq. ft. in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

### **GENERAL HOUSEHOLD PESTS (INDOOR AND OUTDOOR)**

Use the rate specified to control the listed pests. For pests for which a rate range is given, use the lowest rate for low pest pressure or scheduled maintenance control programs and the highest rate for heavy pest pressure, pest-conducive environmental conditions (such as high temperatures or moisture), or when longer residual activity is desired.

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions**

<b>PEST CONTROLLED</b>		<b>CONCENTRATION OF COMBINED ACTIVE INGREDIENTS IN FINAL DILUTION</b>
Ants <sup>1</sup> Beetles (including Multicolored Asian Lady Beetle) Boxelder Bugs Carpenter Bees Carpet Beetles Centipedes Cigarette Beetles Clover Mites Cluster Flies Cockroaches Confused Flour Beetles Crickets Earwigs Firebrats Fleas (indoors)	Flies Grasshoppers Lesser Grain Borers Millipedes Mosquitoes Red Flour Beetles Rice Weevils Saw-toothed Grain Beetles Silverfish Sowbugs Spiders Springtails Stink Bugs (including Brown Marmorated Stink Bug and Kudzu Stink Bug) Ticks	0.065% - 0.13% Mix 0.55 fl. oz. (16 ml.) to 1.1 fl. oz. (32 ml.) of <b>Tent Insecticide CS</b> per gallon of water.
Bed Bugs Elm seed bug <sup>2</sup> Fleas (outdoors)	Scorpions Spotted Lanternfly <sup>2</sup> Wasps, Hornets and Yellow Jackets	0.13% Mix 1.1 fl. oz. (32 ml.) of <b>Tent Insecticide CS</b> per gallon of water.

<sup>1</sup> For treatment of fire ant mounds, see **Applications to Turfgrass** section of this label.

<sup>2</sup> Not Registered for use on Spotted Lanternfly and Elm seed bug by California.

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

**INDOOR PESTS**

**Ants (Including Carpenter Ants and Nuisance Ants):** Apply a 0.065% - 0.13% dilution as a spot, crack-and-crevice, and/or void treatment to ant trails around doors and windows and other places where ants may be found. Apply as a residual surface treatment to active areas such as baseboards; corners; around pipes; in and behind cabinets; behind and under refrigerators, sinks, furnaces and stoves; and in cracks and crevices. For best results, locate and treat nests.

**Bed Bugs:** Efficacy of this product is not affected by heat treatments, so applications can be made before and after heat treatments in integrated bed bug elimination programs.

Apply a 0.13% dilution as a spot, crack-and-crevice, or void treatment to harborage areas. Example harborages include voids behind attached non-upholstered headboards, bed frames (wood and metal only), baseboards, moldings, doorframes, window frames, and shelving; behind mirrors and picture frames; and around edges of carpet or other floor coverings.

Apply a 0.13% dilution with pin-stream, coarse spray or low pressure equipment to cracks and joints of bed frames (wood and metal only), non-upholstered headboards and wood slats and metal coils of box springs. Apply only to areas where prolonged contact by humans or pets will not occur. **DO NOT** use on bed linens, mattresses, or other areas where human contact is common. Applications can be repeated at 7-day intervals as needed to achieve control.

To help remove bed bugs, clean and vacuum infested areas, such as floor surfaces, furniture and mattresses before treatment. Dispose of vacuum bag off-site; with bag-less vacuums, remove all bed bug eggs, nymphs and adults from the canister or capsule off-site.

**Crawling Insects (Beetles, Clover Mites, Cockroaches, Crickets, Earwigs, Firebrats, Silverfish, and Spiders):** Apply a 0.065% - 0.13% dilution as a spot, crack-and-crevice, and/or void treatment to areas where these pests hide, such as baseboards; corners; storage areas; closets; around water pipes, doors, and windows; attics and eaves; cabinets; behind and under refrigerators or other household items such as furniture, sinks, furnaces, and stoves; and to the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices.

**Flying Insects (Flies, Mosquitoes, and Wasps):** Tent Insecticide CS may be applied as a spot or crack-and-crevice treatment directly into cracks and crevices or other non-exposure areas as a low-pressure spray using equipment capable of delivering a pin stream of insecticide inside residential buildings (in attics, crawl spaces, and other non-living spaces), as well as in and around carports, garages, and storage sheds; also see **Outdoor Pests** for specified rates. **DO NOT** make surface applications to living spaces. For wasps, hornets, and yellow jackets, see section on **Control of Yellow Jackets, Hornets, and Wasps**.

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

<b>INDOOR PESTS (continued)</b>
<p><b>Occasional Invaders (Boxelder Bugs, Centipedes, Millipedes, Multicolored Asian Lady Beetles, Pillbugs, Scorpions, Sowbugs, Spotted Lanternfly* and Stink Bugs):</b> Apply a 0.065% - 0.13% dilution as a spot, crack-and-crevice, and/or void treatment around doors and windows, baseboards and other places where these pests may be found or where they may enter premises.</p> <p><b>Pantry Pests (i.e., Carpet beetle, Cigarette beetle, Confused flour beetle, Lesser grain borer, Red flour beetle, Rice weevil, and Saw-toothed grain beetle):</b> Apply a 0.065% - 0.13% dilution as a spot or crack-and-crevice treatment to cupboards, shelving, and storage areas. Remove all utensils, uncovered foodstuffs (or any having original package opened), and shelf paper before making application. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff, or other items. Any foodstuff accidentally contaminated with treatment solution should be destroyed.</p> <p><b>Application within Food-Handling Establishments (places other than private residences in which exposed food is held, processed, prepared, or served, including, but not limited to, areas for receiving, storing, packing (canning, bottling, wrapping, boxing), and preparing foods; edible waste storage; enclosed processing systems (mills, dairies, edible oils, syrups); and serving areas):</b> Use as a crack-and-crevice, spot, or void treatment in and around both food and non-food areas. Apply a 0.065% - 0.13% dilution in small amounts directly into cracks and crevices, using equipment capable of delivering a pin stream of insecticide; in points between different elements of construction; between equipment and floor; in openings leading to voids; and in hollow spaces in walls, equipment and bases. Clean food-contact surfaces and equipment with an effective cleaning compound and rinse with potable water before using.</p> <p>Limit spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments should not exceed 2 sq. ft. Take extreme care that the product is not introduced into the air.</p> <p><b>DO NOT</b> contaminate food and food-processing surfaces. <b>DO NOT</b> treat surfaces likely to be contacted by food. <b>DO NOT</b> apply when food-handling establishments are in operation or foods are exposed. Food must be removed from area being treated. <b>DO NOT</b> apply directly to food or allow applications to contaminate food.</p> <p><b>Application within Food-Serving Areas (facilities where foods are served, such as dining rooms)</b> Apply a 0.065% - 0.13% dilution as a crack-and-crevice, spot, or void treatment to selected surfaces such as under elements of construction and into cracks and crevices. <b>DO NOT</b> treat surfaces likely to be contacted by food. <b>DO NOT</b> apply when facility is in operation or foods are exposed. Food must be covered or removed from area being treated. <b>DO NOT</b> apply directly to food or allow applications to contaminate food.</p> <p>Application of this product in the food areas and/or food-serving areas of food-handling establishments, other than as a spot, crack-and-crevice, or void treatment, is not permitted.</p> <p>Not Registered for use on Spotted Lanternfly and Elm seed bug by California.</p>

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

**OUTDOOR PESTS**

Apply as a crack-and-crevice, spot, or perimeter barrier treatment. To control **ants, carpenter bees, centipedes, cockroaches, crickets, fleas, flies, millipedes, mosquitoes, pill bugs, scorpions, sowbugs, spiders, stink bugs, ticks, wasps**, and other similar perimeter arthropod pests, apply **Tent Insecticide CS** at the rate specified for that particular pest (**Table 1**) with either hand or power application equipment as a residual treatment around buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages, refuse dumps, and other similar areas where these insect pests are active. Apply a treatment volume of 1 - 10 gals./1,000 sq. ft. (approximately 44 - 440 gals./A) unless specified otherwise.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- Applications to soil or vegetation around structures
- Applications to lawns, turf, and other vegetation
- Applications to building foundations, up to a maximum height of 3 feet above grade
- Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure
- Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width
- Applications made through the use of a coarse, low-pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests



**DO NOT** apply to or allow this product to contact plants in bloom or apply if bees are foraging in or adjacent to the treatment area.

**DO NOT** apply this product while bees are foraging.

**DO NOT** apply to plants that are flowering.

Only apply after all flower petals have fallen off.

**Ants:** Apply a 0.065% - 0.13% dilution of **Tent Insecticide CS** as a spot, crack-and-crevice, pin-stream, or perimeter-banded treatment to ant trails and potential entry points such as around doors or windows. As a banded application, apply according to instructions for perimeter treatments below.

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

**OUTDOOR PESTS (continued)**

**Tent Insecticide CS** contains thiamethoxam, a plant-systemic insecticide that can reduce populations of plant-feeding insects, such as aphids, mealybugs, and whiteflies. Control of these insects can reduce the amount of honeydew, a food source for ants foraging in and around structures. To reduce populations of honeydew-producing insects, apply **Tent Insecticide CS** at up to 0.13% as a foliar spray or directed treatment to soil at the base of the landscape plants. When targeting landscape ornamental pests, see directions for use in the **Application to Ornamentals** section.

**Fleas and Ticks:** To control nuisance fleas and ticks (e.g., dog ticks), apply to kennels, yards, runs, and other areas where pets may frequent. For best coverage to control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent. Apply a dilution of 0.065% - 0.13% to control ticks. To control fleas, apply only at the highest concentration of 0.13%. Apply a treatment volume of 2 - 10 gals./1,000 sq. ft. to thoroughly wet the treated surface; use a higher volume when treating landscaping materials, such as mulch or stone, or under dense vegetation in order to obtain the coverage needed for best control.

**DO NOT** apply to pasture or cropland, and **DO NOT** allow animals and people access to treated areas until the spray deposit has dried. Applications can be made on a monthly basis, beginning in the spring, and can continue until frost to control both larvae and adult ticks. **DO NOT** apply directly to pets.

**House Flies and Cluster Flies:** Apply a 0.065% - 0.13% dilution as a spray or low-volume mist to outdoor surfaces of structures, patios, fences, or other areas where flies are likely to collect or rest, or to areas of structures that serve as entry points (doors, windows, under eaves).

Applications under eaves may be made as a surface spray.

For cluster flies, use spot treatments around entry points or fly resting areas of structural surfaces, paying particular attention to south-facing surfaces. In winter and spring, when flies become active and are emerging, interior crack-and-crevice, spot, and void treatments can help reduce the infestation in areas such as attics, crawl spaces, or indoor non-living spaces.

**Mosquitoes:** For adult mosquito control, apply a 0.065% - 0.13% dilution as a structural perimeter spray or spot treatment to landscape plantings and turf, around buildings and foundations, under decks, or to other areas that are likely to serve as adult mosquito resting areas. Yards or other frequented areas enclosed by landscaping can benefit from a banded treatment to reduce invading adult mosquitoes by the treatment of perimeter vegetation. Apply **Tent Insecticide CS** at specified rates at 2 - 5 gallons of water per 1,000 sq. ft. (87 - 218 gals./A). Low-volume mist blowers may be used as well. Application to vegetation away from structures may require additional state certifications, e.g., in turf or ornamental categories. Consult your state regulatory agency for requirements.

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

**OUTDOOR PESTS (continued)**

**Occasional Invaders and Aggregating Pests (Beetles, Boxelder bugs, Centipedes, Clover mites, Crickets, Earwigs, Grasshoppers, Scorpions, Silverfish, Spiders, Springtails, Sowbugs, Spotted Lanternfly\* and Stink bugs):** Apply a 0.065% - 0.13% dilution in a spray volume of 1 - 10 gals./1,000 sq. ft. to outdoor structural surfaces where insects are found and around likely entry points such as doors, windows, and utility penetrations. Banded applications as described in the **Outdoor Pests - Perimeter Treatments around Structures** section (below) can provide additional protection against these pests entering the structure.

For Spotted Lanternfly\* aggregations and egg masses on or near structures, or other similar surfaces, use spot treatments around entry points or on resting areas of structural surfaces. See directions for perimeter treatments around structures when targeting outdoor pests.

It is recommended that egg masses on trees, cement blocks, rocks, and any other hard surface be scraped and removed. Spray Spotted Lanternfly\* nymphs, adults and eggs directly whenever possible.

**Control of Yellow Jackets, Hornets, and Wasps (Localized Treatment):** Use caution when treating nests of stinging insects as **Tent Insecticide CS** does not provide instant knockdown. Protective equipment for the applicator may be required to reduce the risk of insect stings during application. The use of an extension wand is recommended when applying the liquid or foam to nests to reduce the risk of being stung. For best results, treat nests in the evening when most insects will be present.

For control of ground-nesting yellow jacket wasps, make an application of 0.13% liquid or finished foam into the below-ground nest. Apply a sufficient amount of liquid or foam to fill the nest and coat the opening to the nest. The use of an extension wand is recommended when applying to reduce the risk of being stung.

For application to paper-covered wasp nests (e.g., bald-faced hornets), apply a 0.13% dilution of **Tent Insecticide CS** as a finished foam or as a spray into the opening of the nest. Apply a sufficient amount of foam to fill the nest without allowing runoff. As a spray, apply product to the entire outside surface of the nest as well as the nest opening. The use of an extension wand is recommended when applying the foam to reduce the risk of being stung.

For visible wasp nests, spray the outside of the nest with a 0.13% dilution. Apply spray to the entire surface of the nest. For wasp nests within voids such as under soffits and behind rain gutters, make an application of 0.13% liquid or foam into the nesting area. Apply a sufficient amount of foam to fill the nesting void.

*(continued)*

**Table 1: General Household Pests (Indoor and Outdoor),  
Application Instructions, and Pest-Specific Directions (continued)**

**OUTDOOR PESTS (continued)**



**DO NOT** allow this product to contact plants in bloom if bees are foraging in or adjacent to the treatment area.

**DO NOT** apply this product while bees are foraging.

**DO NOT** apply to plants that are flowering.

Only apply after all flower petals have fallen off.

**Outdoor Pests – Perimeter Treatments Around Structures:** Applying a continuous band of insecticide around a building foundation and around windows, doors, service line entrances, eaves, vents, and other areas can greatly reduce the potential for entry by crawling pests. For best results, before application, remove debris and leaf litter from next to the foundation; cut back vegetation and branches that touch the foundation, walls, or roof; and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation.

Apply **Tent Insecticide CS** in a volume of 1 - 10 gallons per 1,000 sq. ft. or 44 - 440 gals./A around the structure (according to state regulations governing commercial pest control and the previously stated restrictions on outdoor applications) and upwards along the foundation to 3 feet and spot treatments around windows, doors, and roof overhangs. The amount of **Tent Insecticide CS** to be diluted in water will depend upon the pest species, the infestation level, and the service interval desired (see **Table 1** for application rates). Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active.

Agitate occasionally during application.

Not Registered for use on Spotted Lanternfly and Elm seed bug by California.



**Table 2: Dilutions for Structural Perimeter Applications**

APPLICATION RATE (AMOUNT OF TENT INSECTICIDE CS PER 1,000 SQ. FT.)	APPLICATION VOLUME (GALLONS OF SOLUTION PER 1,000 SQ. FT.)	AMOUNT OF TENT INSECTICIDE CS TO DILUTE IN WATER ACCORDING TO SPRAY TANK VOLUMES		
		10 gallons	25 gallons	50 gallons
0.55 fl. oz. (16 ml.)	1	5.5 fl. oz. (163 ml.)	13.8 fl. oz. (408 ml.)	27.5 fl. oz. (813 ml.)
	2	2.8 fl. oz. (83 ml.)	6.9 fl. oz. (204 ml.)	13.8 fl. oz. (408 ml.)
	5	1.1 fl. oz. (33 ml.)	2.8 fl. oz. (83 ml.)	5.5 fl. oz. (163 ml.)
	10	0.55 fl. oz. (16 ml.)	1.4 fl. oz. (41 ml.)	2.8 fl. oz. (83 ml.)
1.1 fl. oz. (32 ml.)	1	11.0 fl. oz. (325 ml.)	27.5 fl. oz. (813 ml.)	55 fl. oz. (1,627 ml.)
	2	5.5 fl. oz. (163 ml.)	13.8 fl. oz. (408 ml.)	27.5 fl. oz. (813 ml.)
	5	2.2 fl. oz. (65 ml.)	5.5 fl. oz. (163 ml.)	11.0 fl. oz. (325 ml.)
	10	1.1 fl. oz. (33 ml.)	2.8 fl. oz. (83 ml.)	5.5 fl. oz. (163 ml.)

#### **Control of Wood-Destroying Insects**

Use **Tent Insecticide CS** to control the following wood-destroying insects. For specific application instructions, refer to label text.

**Table 3: Wood-Destroying Pests and Pest-Specific Use Directions**

PESTS	APPLICATION METHOD	CONCENTRATION OF COMBINED ACTIVE INGREDIENTS IN FINAL DILUTION	COMMENTS
Carpenter Ants <sup>1</sup> Carpenter Bees Drywood Termites Wood-Destroying Beetles and Borers	<i>Remedial Control:</i> Liquid or foam - Gallery treatment <i>Preventive treatment:</i> Wood Surface Spray	0.13% Mix 1.1 fl. oz. (32 ml.) <b>Tent Insecticide CS</b> per gallon of water.	Re-treat as directed.
<sup>1</sup> Carpenter ants may be controlled at a lower rate of 0.065%; rates of 0.065% to 0.13% may be used per instructions under <b>Control of Carpenter Ants</b> below.			
<p><b>Control of Drywood Termites:</b> When used as specified in this label, <b>Tent Insecticide CS</b> provides effective remedial control of localized infestations of drywood termites, including species of <i>Incisitermes</i>, <i>Cryptotermes</i>, and <i>Marginitermes</i>. Knowledge of the biology and behavior of the drywood termite species involved, the locations, and the extent of the infestation(s) will help to ensure successful control.</p> <p>Treatment requirements for drywood termite control may vary due to state and local regulations. For advice concerning current drywood termite control regulations under local conditions, consult your State structural pest control regulatory agency.</p> <p><b>Directions for Remedial Control of Drywood Termites in Infested Wood:</b> To control drywood termites in localized areas of infested wood in structures, apply 0.13% <b>Tent Insecticide CS</b> as a liquid or foam to voids and galleries in damaged wood, in spaces between wooden structural members, or between wood and foundations. Locate galleries by using visual signs (e.g., fresh fecal pellets or blistered wood), the presence of live termites, mechanical sounding techniques (e.g., tapping on the wood surface and listening for changes in sound to indicate changes in wood density), listening devices, motion detection devices or other technologies that help pinpoint drywood termite activity.</p> <p><b>Wood Injection Method:</b> Drill small diameter holes of appropriate size for the injection tip or use a self-puncturing tip positioned to intersect termite galleries within infested wood. Drywood termite emergence or pellet kick-out holes connect directly to galleries and are indicators of potential sites to drill and inject <b>Tent Insecticide CS</b>. Care should be taken to avoid electrical wiring, plumbing, etc. when drilling and injecting. <b>DO NOT</b> drill or puncture completely through wood. Spacing of the holes will depend on the distribution of insect activity and galleries. Injection holes may be clustered in areas with insect activity as indicated by damage, live insects, or other indicators previously described. Injection holes on opposite sides of large (4" x 10" or larger) structural beams may be necessary to effectively penetrate galleries.</p>			

(continued)

**Table 3: Wood-Destroying Pests and Pest-Specific Use Directions (continued)**

Apply up to 50 ml. (1.7 fl. oz.) of 0.13% dilution at each injection hole. If using **Tent Insecticide CS** as a foam preparation, inject a sufficient amount to fill galleries without allowing runoff.

Re-apply if insect activity within treated areas is detected four or more weeks following treatment. For best results in treating galleries, inject **Tent Insecticide CS** into new injection holes positioned between previous injection sites.

**Wood Surface Treatment:** Surface applications may also be used to supplement spot treatment for drywood termites as a means to prevent re-infestation by swarming adult termites. Apply a 0.13% dilution as a surface spray or mist to exposed wood surfaces in areas not used as living spaces, such as attics, crawl spaces, unfinished basements, or structural voids. Apply treatment until surface is thoroughly wetted.

**Control of Carpenter Ants:** Apply a 0.065% - 0.13% dilution to ant trails around doors and windows and other places where ants may be found. Indoors, apply as a residual surface treatment to active areas such as baseboards; corners; around pipes; in and behind cabinets; behind and under refrigerators, sinks, furnaces, and stoves; and in cracks and crevices. For best results, locate and treat nests. Outdoors, apply as a crack-and-crevice, spot, or banded treatment; also see **Outdoor Pests**.

Inside voids where ants or nests occur, drill (or use existing opening) and inject spray or foam into the cavity. Inside decking materials or other structural materials, poles, or trees infested with carpenter ants, drill and inject spray or foam into cavities. Apply up to 50 ml. (1.7 fl. oz.) of 0.065% - 0.13% dilution at each injection hole. If using **Tent Insecticide CS** as a foam preparation, inject a sufficient amount to cover the area without allowing runoff.

**Control of Carpenter Bees (localized treatment):**

**Gallery Treatment:** Apply a 0.13% dilution of **Tent Insecticide CS** into carpenter bee galleries as a spray, mist, or foam application. As a liquid, apply up to 50 ml. (1.7 fl. oz.) of finished solution of **Tent Insecticide CS** at each injection hole. If using **Tent Insecticide CS** as a foam preparation, inject a sufficient amount to fill galleries without allowing runoff.

**Wood Surface Treatment:** Apply coarse spray as a spot treatment to thoroughly wet wood surfaces where bees have been previously active or to provide protection against further damage. Apply early in the spring to prevent bees from invading wood. When bees have infested wood, surface applications can help control embedded larvae and bees that emerge from the wood. Applications can be made on a monthly basis to maintain protection of treated areas.

*(continued)*

**Table 3: Wood-Destroying Pests and Pest-Specific Use Directions (continued)**

**Control of Wood-Destroying Beetles and Borers:** For control of wood-destroying beetles or borers, including old house borers, powderpost beetles, false powderpost beetles, death-watch beetles, and ambrosia or bark beetles. Apply **Tent Insecticide CS** diluted to 0.13% as a spray, mist, or foam to galleries, in structural voids, and/or as wood surface treatments.

For old house borers, or other beetles that form large galleries, treat the gallery system by drilling and injecting product with sufficient volume to cover the galleries.

For beetles or borers that **DO NOT** form galleries which can be readily injected (e.g., powderpost beetles), apply product as a liquid, mist, or foam to exposed wooden surfaces in non-living space areas in crawlspaces, basements, and attics; to structural voids; to spaces between wood elements of a structure; or at joints between wood and foundations.

Apply as a coarse liquid spray with low pressure (<25 psi), as a foam or mist application, or by brushing diluted product onto the surface. Applications should be made in sufficient volumes to coat the target surface, but less than the volume that creates runoff. Re-treat as needed to maintain protection. Allow treated surfaces to dry before contacting them.

#### **Non-Residential Lawn, Turfgrass, and Ornamentals**

**Tent Insecticide CS** contains two insecticides, thiamethoxam and lambda-cyhalothrin. Thiamethoxam and lambda-cyhalothrin provide long-lasting, broad-spectrum contact and systemic control of turf and ornamental plant insects.

**Tent Insecticide CS** may be used for applications to maintain outdoor areas where turf and ornamentals are grown, such as non-residential landscapes around institutional, public, commercial, and industrial buildings; parks; recreational areas; and athletic fields. Consult your state extension office or pesticide regulatory officials for further information.



- **DO NOT** apply this product while bees are foraging.
- **DO NOT** apply to plants that are flowering.
- Only apply after all flower petals have fallen off.

#### **Non-Residential Lawn, Turfgrass, and Ornamentals Use Restrictions**

- **DO NOT** apply this product to lawns/turfgrass or allow it to drift to blooming plants or weeds if bees are foraging in or adjacent to the treatment area. Time application during periods when pollinating insects are not present, such as early morning or late evening. **DO NOT** apply to plants/ornamentals that are flowering. Only apply after all flower petals have fallen off.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or commercial sod production.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product to edible crops.
- **DO NOT** apply this product by aerial application.
- **DO NOT** water the treated area to the point of runoff.
- **DO NOT** make applications during rain.
- **DO NOT** apply more than 32 fl. oz. of **Tent Insecticide CS** per acre per calendar year.

### **Mixing Instructions (Non-Residential Turf and Ornamental Dilutions)**

**Tent Insecticide CS** may be mixed with water and used in all types of standard application equipment. Fill applicator tank with 1/2 to 3/4 volume of water and add **Tent Insecticide CS**. Adjust water pH with a buffering agent if necessary to achieve a pH of 5 - 7. Slowly add **Tent Insecticide CS** to applicator tank water with maximum agitation. Fill the tank to the appropriate volume as determined by the listed rates and the size of the tank and continue to agitate while making applications. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only the amount of treatment volume required.

### **Tank-Mixing (Non-Residential Turf and Ornamental Dilutions)**

**Tent Insecticide CS** may be tank-mixed with other currently-registered pesticides unless expressly prohibited by the product label. To ensure compatibility, conduct a small-volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Tent Insecticide CS** last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

### **Application To Non-Residential Turfgrass**

**Sites of Application** - **Tent Insecticide CS** may be used for applications to turfgrasses around non-residential, institutional, public, commercial and industrial buildings; parks; recreational areas; and athletic fields.

**Broadcast Applications to Non-Residential Turfgrass** - Apply **Tent Insecticide CS** as a broadcast application to turfgrass for control of targeted pests. Use sufficient water volume (2 - 10 gals./1,000 sq. ft.) to uniformly distribute **Tent Insecticide CS** over the area being treated and to adequately move the active ingredient into the canopy and thatch layers. **Tent Insecticide CS** is not phytotoxic to any major turfgrass species. **Tent Insecticide CS** can be applied to established turf or at seeding or during seed germination.

**Application Timing** - Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may impact the performance of **Tent Insecticide CS** against white grubs. **DO NOT** mow the turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into the turfgrass or soil. **DO NOT** apply when turfgrass is waterlogged or when soils are saturated with water (i.e., will not accept irrigation).

**DO NOT** apply more than 32 fluid ounces of **Tent Insecticide CS** per acre per calendar year.

**DO NOT** allow this product to contact plants in bloom.

**KEEP CHILDREN AND PETS OFF TREATED AREAS FOLLOWING THE APPLICATION UNTIL SPRAY HAS DRIED.**

**Table 4: Turf Pests, Application Instructions, and Pest-Specific Use Directions**

PEST		DOSAGE OF TENT INSECTICIDE CS
Ants	Sod webworms	<p>Make a single application of 0.6 fl. oz. (19 ml.) per 1,000 sq. ft. (or 28 fl. oz. per acre.) Or Make two applications of 0.3 fl. oz. (10 ml.) per 1,000 sq. ft. (or 14 fl. oz. per acre) on a 3- to 5-week interval.</p>
Armyworms	White grubs <sup>3</sup>	
Black turfgrass atenius (adults)	Aphodius	
Bluegrass billbug (adults and larvae)	Asiatic Garden Beetle	
Chinch bugs <sup>1</sup>	Black Turfgrass Ateenius	
Crickets	European Chafer	
Cutworms	Green June Beetle	
Fire Ants	Japanese Beetle	
Fleas	May or June Beetle	
Hyperodes weevil (adults and larvae)	Northern Masked Chafer	
Japanese beetles (adult)	Oriental Beetle	
Mole Cricket (young adults and nymphs) <sup>2</sup>	Southern Masked Chafer	
Chiggers		<p>Make three applications of 0.2 fl. oz. (7 ml.) per 1,000 sq. ft. (or 10.5 fl. oz. per acre) on a 3- to 5-week interval.</p>
Grasshoppers		

<sup>1</sup> **Chinch Bug** - Apply when insects are first observed. Repeat applications on a 3- to 5-week interval to obtain control.

<sup>2</sup> **Mole Crickets** - Apply from first egg hatch to peak egg hatch. If the soil is dry, irrigate after application to move insecticides into the soil.

<sup>3</sup> **White grubs** - Begin applications up to 45 days before the historical peak of adult flight to second instar grub of the species being targeted. For optimum control, treat from peak flight to peak egg hatch. Irrigate to move **Tent Insecticide CS** into the root zone. Full-season control of white grubs will be obtained with either a single 28 fl. oz. application or two 14 fl. oz. applications.

*(continued)*

**Table 4: Turf Pests, Application Instructions, and Pest-Specific Use Directions (continued)**

Apply to turf prior to the establishment of high insect pressure or significant turf damage.

**Fire Ant Mound Treatment** - Treat individual mounds with a drench application using a watering can. Use 0.15 fluid ounces of **Tent Insecticide CS** per 2.5 gallons of water. Thoroughly soak each mound and a 3 ft. diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. For best results, apply in early morning or late evening hours. Applications can be made on a monthly basis to maintain protection of treated areas.

**Surface Insect Control (including armyworm, cutworms, and fleas)** - Apply **Tent Insecticide CS** at specified rates in 2 - 5 gallons of water per 1,000 sq. ft. (87 - 218 gals./A). The use of a spreader-sticker may be useful if high rainfall amounts are forecast; otherwise, the addition of adjuvants is not necessary under normal conditions for surface insect control in turf.

Delay watering or mowing for 12 - 24 hours after application for optimum control of surface-feeding insect pests.

**Thatch-Inhabiting Insect Control (including chinch bugs)** - Apply **Tent Insecticide CS** at specified rates in 2 - 10 gallons of water per 1,000 sq. ft. (87 - 440 gals./A). The use of a nonionic wetting agent, penetrant, or similar adjuvant is recommended. Irrigate after application with up to 1/2 inch of water to move the **Tent Insecticide CS** into the thatch layer. If irrigation is not available, then use high water application rates volumes (>4 gals./1,000 sq. ft. or 174 gals./A) for optimum results.

For specific information about developmental stages of the target pest and optimal timing of applications, consult with your State Cooperative Extension Service.

### **Application to Ornamentals**

Apply to ornamentals in non-residential landscaped areas and landscaped areas around institutional, public, commercial and industrial buildings; parks; recreational areas; and athletic fields (including trees, shrubs, flowers, evergreens, foliage plants, and groundcovers).

**DO NOT** apply more than 32 fluid ounces (0.26 lb. a.i./A thiamethoxam and 0.08 lb. a.i./A lambda cyhalothrin) of **Tent Insecticide CS** per acre per calendar year.

**DO NOT** apply when the wind speed is greater than 15 mph.



- **DO NOT** apply this product while bees are foraging.
- **DO NOT** apply to plants that are flowering.
- Only apply after all flower petals have fallen off.

**DO NOT** apply this product, by any application method, to linden, basswood, or *Tilia* species.

**Table 5: Pests of Ornamental Plants, Application Instructions,  
and Pest-Specific Use Directions**

<b>PESTS<sup>1</sup></b>		<b>DOSAGE OF TENT INSECTICIDE CS</b>
Adelgids (including Hemlock wooly adelgid)	Leaf gall-forming insects	4.2 - 13.9 fl. oz. per 100 gallons or 125 - 410 ml. per 100 gallons
Armyworms	Leafhoppers	
Azalea caterpillars	Leafminers (adults)	
Aphids	Leaf rollers	
Bagworms	Leaf skeletonizers	
Black vine weevils (adult)	Mealybugs	
Boxelder bugs	Midges	
Budworms	Oleander moth larvae	
California oakworms	Pine shoot beetles	
Cankerworms	Pinetip moths	
Crickets	Plant bugs	
Cutworms	Psyllids (including Asian citrus psyllid)	
Elm leaf beetles	Root weevils	
Fall webworms	Sawflies	
Flea beetles	Scale insects (crawlers)	
Grasshoppers	Spittlebugs	
Japanese beetles (adults)	Striped beetles	
June beetles (adults)	Thrips	
Lace bugs	Tip moths	
Leaf-feeding caterpillars	Tussock moth larvae	
Broadmites	Mealybugs	8.5 - 13.9 fl. oz. per 100 gallons or 250 - 410 ml. per 100 gallons
Brown soft scale	Pine needle scale (crawlers)	
California red scale (crawlers)	Whiteflies	
Clover mites		

*(continued)*



**Table 5: Pests of Ornamental Plants, Application Instructions,  
and Pest-Specific Use Directions (continued)**

PESTS <sup>1</sup>	DOSAGE OF TENT INSECTICIDE CS
Spotted Lanternfly <sup>2</sup>	13.9 fl. oz. per 100 gallons or 410 ml. per 100 gallons
<sup>1</sup> For nuisance pests other than plant-feeding pests (e.g., spiders, cockroaches, mosquitoes), refer to <b>Outdoor Pests</b> section for use instructions. <sup>2</sup> Not approved for use in California.	
<p>Start application to ornamentals prior to the establishment of high insect pest populations. Make reapplications as directed to keep pest populations under control, using higher rates within the listed rate range as pest pressure increases.</p> <p>Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatments to spot treatments.</p> <p>Good spray coverage is necessary to provide the most effective level of control. Spray uniformly to wet plant, but not to the point of runoff or drip. Addition of a spreader-sticker at specified rates may enhance the control of insects on certain species of ornamentals having waxy, hard-to-wet foliage.</p> <p>Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.</p> <p><b>NOTE:</b> While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution. Spray a selection of ornamental plants and observe them for 7 - 10 days prior to treating large areas if local use experience is unavailable. This is especially advisable if <b>Tent Insecticide CS</b> is being mixed with another product or ingredient besides water. See <b>Tank Mixing (Turf and Ornamental Dilutions)</b> instructions.</p> <p><b>Scale insects:</b> Thoroughly cover the plant with <b>Tent Insecticide CS</b> spray, including trunks, stems, twigs, and foliage for control of scale insects.</p> <p><b>Bagworm:</b> Apply <b>Tent Insecticide CS</b> when bagworm larvae begin to hatch. Spray directly on the larvae. Application is most effective when the larvae are young.</p> <p><b>Spotted Lanternfly*:</b> Successful control of this pest requires thorough coverage of plant trunks, stems, branches and foliage. Applications of <b>Tent Insecticide CS</b> made with hydraulic spray equipment will require higher water volumes, up to 200 gallons per acre.</p> <p>Not Registered for use on Spotted Lanternfly and Elm seed bug by California.</p>	

### **Ornamental, Non-Bearing Fruit and Nut Trees and Woody Shrubs**

**Tent Insecticide CS** contains thiamethoxam, which is a systemic insecticide that will be translocated upward into the plant from root uptake. **Tent Insecticide CS** must be placed in an area where roots can absorb the active ingredient. It may take 2 - 4 weeks for the active ingredient to move from the root zone to the area of the trees affected. If insect pests are already present and causing damage, consider making a foliar application and then follow up with a soil application. For best results, water in after application with enough water to move the insecticide into the root zone of the tree.

For trees and shrubs, use 0.2 fluid ounces (6 ml.) of **Tent Insecticide CS** per inch cumulative trunk diameter (DBH) of the plant(s) or per foot of shrub height (less than 6 ft. tall).

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal. **DO NOT** pour or dispose down the drain or sewer. Call your local solid waste agency for local disposal options.

**PESTICIDE STORAGE:** Keep this product in its tightly closed original container. **DO NOT** store in areas where temperatures are below 32 degrees F. Store in a cool, dry (preferably locked) area away from other pesticides, fertilizer, food, or feed that is inaccessible to children and animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**Rigid HDPE Plastic Less Than or Equal to 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities.

**Rigid HDPE Plastic Greater Than 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

**Rigid HDPE Plastic Bulk and Mini-Bulk HDPE Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**



### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
LAMBDA-CYHALOTHRIN	GROUP	3A	INSECTICIDE

# Tent Insecticide CS

For indoor and outdoor use.

For Use In, On, and Around Buildings and Structures for the Control of Listed Pests, on Non-Residential Lawns and Turf, Ornamental Trees and Shrubs Around Non-Residential, Institutional, Commercial, and Industrial Buildings.

<b>ACTIVE INGREDIENT:</b>	<b>WT. BY %</b>
Thiamethoxam (CAS No. 153719-23-4) .....	11.6%
Lambda-cyhalothrin* (CAS No. 91465-08-6) .....	3.5%
<b>OTHER INGREDIENTS:</b> .....	84.9%
<b>TOTAL:</b> .....	<b>100.0%</b>

\* Microencapsulated

Contains 1.04 lbs. thiamethoxam and 0.32 lb. lambda-cyhalothrin in a CS formulation.

In the State of New York, this product may only be applied for indoor uses.

## KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for First Aid, Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

**Sharda USA LLC** 

7217 Lancaster Pike, Suite A  
Hockessin, Delaware 19707

EPA Reg. No. 83529-357

EPA Est. No. **AG** 72159-GA-001; **MA** 83411-MN-001;

**MC** 89332-GA-001; **SC** 39578-TX-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

**Net Contents: 1 Quart (0.25 Gal.)**