

Nile 70% EC

For Use on Vegetables, Fruits, Nuts, Melons, Outdoor Ornamental Plants, and Agronomic Crops.

For Foliar Control of Fungal Disease and Insect Pests.

Can be Used Up to Day of Harvest.

ACTIVE INGREDIENT:

Clarified Hydrophobic Extract of Neem Oil* **WT. BY %** 70.0%

OTHER INGREDIENTS: **30.0%**

TOTAL: **100.0%**

*This product contains 5.46 lbs. of clarified hydrophobic extract of neem oil per US gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-102-83529

EPA Est. No. **AG** 72159-GA-001; **MA** 83411-MN-001;

MC 89332-GA-001; **SC** 39578-TX-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

This container will treat up to a half-acre when used at the highest application rate (only for containers less than 5 gallons).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, including barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is hazardous to fish and aquatic invertebrates.

For Terrestrial Uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. **DO NOT** apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, including barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinylchloride (PVC) or Viton
- Shoes plus socks

PRODUCT INFORMATION

Thorough coverage is necessary to provide good disease and mite control. **Nile 70% EC** prevents fungal attack of plant tissue and contact activity to mite pressure on the plant. Avoid excessive runoff of spray material for maximum control. For best results, maintain constant agitation in spray equipment. When used in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale. As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn. Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

Restrictions:

- For optimal performance, **DO NOT** mix with cold water (less than 45°F).
- **DO NOT** apply when weather conditions favor drift or when the likelihood of runoff is high.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply to known sensitive plant species, including impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
- **DO NOT** apply to wilted or otherwise stressed plants, or to newly transplanted materials or crops prior to root establishment.
- **DO NOT** use this product after bloom on table grapes or following bunch closure on wine grapes.
- **DO NOT** use this product after pit hardening on stone fruit.

Tank Mixing

Nile 70% EC has been found to be compatible with most commonly used fungicides, insecticides and fertilizers. Check physical compatibility first with other products before mixing by using a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. After thoroughly mixing, let mixture stand for at least 5 minutes. Growers must tank mix combinations for phytotoxicity on a sample of plants prior to use. Due to the wide variation in climatic conditions, cultural practices, and other factors, the User assumes full responsibility for any crop damage or other liability resulting from the use of **Nile 70% EC** in a tank mix combination. Avoid tank mixes with captan, chinomethionat, chlorothalonil, and sulfur, or other chemically similar products as unpredictable results or leaf burn may occur.

Mixing Instructions

Add **Nile 70% EC** to one-half (1/2) full tank under agitation containing water of 45°F or greater before filling to desired level. If water temperature is below 45°F, premix **Nile 70% EC** at a 1:1 ratio with tepid water to ensure good emulsification. Then dilute to final volume. When combining with other products, including wettable powder insecticides or fungicides, add these items first when the tank is approximately 1/3 full. Ensure that there is good agitation while mixing for complete emulsification. Maintain agitation during spray application. **DO NOT** use if this oil does not form a uniform cloudy emulsion.

LANDSCAPE AND NURSERY USE INSTRUCTIONS

FUNGICIDE USE

Ornamental Disease Control

Nile 70% EC is an effective fungicide for the prevention and control of various fungal diseases including black spot on roses, powdery mildew, downy mildew, anthracnose, rust, leaf spot, botrytis, needle rust, scab, flower, twig, and tip blight, and alternaria.

Nile 70% EC will provide broad-spectrum control of many fungal diseases affecting ornamental flowering plants, trees, shrubs, and foliage when used at the rate of 1 gallon per 100 gallons of water. Apply **Nile 70% EC** at a rate of 1:100 when spraying as a preventative for outdoor applications. Under extreme outdoor conditions, apply **Nile 70% EC** at the 1:50 application rate (refer to the below **Management of Landscape and Ornamental Diseases** and/or **Management of Turf Diseases** tables for detailed use rates). This will provide the necessary control under the extreme outdoor conditions. When using **Nile 70% EC** as a preventative, make application on a 14-day schedule. If disease is present, use a 7-day application schedule to control the disease and prevent further damage from occurring.

Application Instructions

Apply until plant surface is uniformly wet with minimal runoff at 25 - 40 PSI with hand sprayer or 100 - 200 PSI with power sprayer as a fine spray to both leaf surfaces. Avoid excessive application.

Management of Landscape and Ornamental Diseases

Spray Target	Concentration	Spray Interval	Restrictions
Outdoor Flowering Plants	1:50 - 1:100 (2% - 1%)	7 - 14 days	DO NOT spray impatiens flowers.
Outdoor Woody Plants	1:50 - 1:100 (2% - 1%)	14 days	-

Management of Turf Diseases

Spray Target	Concentration	Spray Interval	Precautions
Snow Mold	1:100 - 1:50 (1% - 2%)	14 days	In Fall before first snow.
Dollar Spot	1:100 - 1:200 (1% - 0.5%)	7 - 14 days	At first sign of disease.
Brown Patch	1:100 - 1:200 (1% - 0.5%)	7 - 14 days	At first sign of disease.

INSECTICIDE/MITICIDE USE

For use to control whiteflies, mealybugs, aphids, leafhoppers, mites and scales on ornamentals, trees, and shrubs in and around commercial nurseries and other commercial structures.

Use **Nile 70% EC** at the rate of 1 - 2 gals. per 100 gals. of water. The low rate (1%) can be used when pest pressure is low. Use the high rate (2%) for high pest infestations or to control more troublesome pests, including mites and scales. Determine the best rate for your particular situation.

Make sprays on a 7- to 14-day interval depending on the severity of the pest problem. Use the 7-day schedule until the pest population is reduced. Then use a 14-day interval for control.

AGRICULTURAL USE INSTRUCTIONS

Nile 70% EC is a broad-spectrum fungicide of certain diseases and controls mites in citrus, deciduous fruits and nuts, vegetable crops, cereal grains, and other miscellaneous crops.

Apply **Nile 70% EC** by conventional, low-volume, and aerial application equipment.

Restrictions:

- **DO NOT** use **Nile 70% EC** after bloom on table grapes or following bunch closure on wine grapes. **DO NOT** use **Nile 70% EC** after pit hardening on stone fruit.
- Use care when applying near streams, ponds, lakes, or other bodies of water. **DO NOT** apply **Nile 70% EC** when weather conditions favor drift or when the likelihood of runoff is high.

Pest Control	Suppression	Diseases		
Aphids	Whiteflies	<i>Alternaria</i>	Greasy Spot	Molds
Mealybugs	Thrips	<i>Anthraco</i>	Leaf Spot	Scabs
Mites		Early Blight	Downy Mildew	Rusts
Rust Mites		Leaf Blight	Post-Bloom Fruit Drop	Shothole*
Spider Mites		<i>Botrytis</i>	Powdery Mildew	
Scales (Soft Scale)				

*Not Registered for Use By California.

For Diseases: Apply **Nile 70% EC** as a 1% solution in sufficient amounts of water to achieve complete coverage of foliage (see the **SPECIFIC CROP LISTINGS** for directions per crop groupings). When used as a foliar fungicide, begin applications when conditions are favorable for disease but before infection. Apply **Nile 70% EC** every 7 - 14 days until threat of disease is over. Use **Nile 70% EC** in rotation with other fungicides to increase the spectrum of control.

For Mites: For best results, apply **Nile 70% EC** in sufficient amounts of water to achieve complete coverage. **Nile 70% EC** is most effective when applied before mites or eggs are present in large numbers (see the **SPECIFIC CROP LISTINGS** for directions per crop groupings). Apply at a rate of 1% - 2% **Nile 70% EC**. Reapply **Nile 70% EC** every 7 - 21 days until pressure is over.

For Insects: For best results, apply **Nile 70% EC** at 1% - 2% in sufficient amounts of water to achieve complete coverage. **Nile 70% EC** is most effective when applied before insects or eggs are present in large numbers. Repeat application based on pest pressure and field scouting for best results. Use the dilution tables to calculate dilution rates.

Application Instructions

- **Ground Application:** Apply **Nile 70% EC** by ground application in a minimum of 25 gals. of water per acre.
- **Chemigation:** **DO NOT** apply this product through any type of irrigation system.
- **Aerial Application:** For those crops where aerial applications are indicated, apply a minimum of 5 gals. of water per acre unless otherwise specified. Avoid application under conditions when uniform coverage cannot be obtained or when excessive drift may occur. **DO NOT** apply directly to humans, animals, or aquatic animals.

Dilution Table for Low-Volume (Including Aerial) Applications (5 - 30 Gallons per Acre)

Gallons of Water	Nile 70% EC	DILUTION
5	1 Quart*	5%
10	1 Quart*	2.5%
15	1 Quart*	1.67%
20	1 Quart*	1.25%
25	1 Quart*	1%
30	1 Quart	0.83%

*For Citrus, Stone Fruit, Tropical Fruit, Pome Fruit, and Nuts, refer to specific application instructions.

Dilution Table for High-Volume Application (25 - 250 Gallons per Acre)*

Gallons of Water	Gals. of Nile 70% EC (0.5% Solution)	Gals. of Nile 70% EC (1% Solution)	Gals. of Nile 70% EC (1.5% Solution)	Gals. of Nile 70% EC (2% Solution)
25	Use 1% solution	0.25 (32 fl. oz.)	0.375 (48 fl. oz.)	0.5 (64 fl. oz.)
50	0.25 (32 fl. oz.)	0.5 (64 fl. oz.)	0.75 (96 fl. oz.)	1
75	0.375 (48 fl. oz.)	0.75 (96 fl. oz.)	1.125	1.5
100	0.5	1	1.5	2
125	0.625 (80 fl. oz.)	1.25	1.875	2.5
150	0.75	1.5	2.25	3
175	0.875 (112 fl. oz.)	1.75	2.625	3.5
200	1	2	3	4
225	1.125	2.25	3.375	4.5
250	1.25	2.5	3.75	5

*For Vegetable, Cereal Grains, and Other Miscellaneous Crops, refer to specific use directions.

SPECIFIC CROP LISTINGS

BULB, COLE, AND LEAFY VEGETABLES

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Arugula	Chicory	Collard Greens	Parsley
Broccoli	Cilantro	Mustard Greens	Purslane
Chinese Broccoli (Gai Lan)*	Collards	Rape Greens	Radicchio
Brussels Sprouts	Corn Salad*	Kale	Rapini*
Cabbage (Heads and Leaf)	Cress	Kohlrabi	Rhubarb
Chinese Cabbage (Bok Choy, Napa)	Watercress	Leek	Shallots
Chinese Mustard Cabbage (Gai Choy)*	Endive	Lettuce (Head and Leaf)	Spinach
Cauliflower	Fennel	Low-Bok*	Chinese Spinach (Amaranth, Tampala)
Celery	Garlic	Onions	Swiss Chard
Celtuce*	Greens*	Orach*	Turnip Tops*
Chervil			

*Not Registered for Use By California.

CITRUS, POME (DECIDUOUS FRUITS), STONE, AND TROPICAL FRUITS

CITRUS

Use **Nile 70% EC** on citrus to control diseases to include Anthracnose, Post-Bloom Fruit Drop, and Scab. Apply as a foliar application at the rate of 1% in 25 - 200 gals. of water for high-volume applications or a minimum of 0.5 gal. **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume applications. **DO NOT** apply more than 2.5 gals. of **Nile 70% EC** per acre, per application.

Reapply **Nile 70% EC** every 7 - 14 days until threat of disease is over.

For control of Alternaria and Greasy Spot, apply **Nile 70% EC** at a 1% solution in 25 - 125 gals. of water per acre in combination with copper (refer to EPA registered product label for specific use and application information) as a preventative disease spray tank mixture.

Use **Nile 70% EC** on citrus to control mites at a rate of 1% - 2% in 25 - 200 gals. of water for high-volume application or 0.5 gal. **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. As a dormant oil spray, use **Nile 70% EC** as a 2% solution or 1 quart **Nile 70% EC** minimum per acre. **DO NOT** apply more than 5 gals. of **Nile 70% EC** per acre, per application.

POME (DECIDUOUS FRUITS)**Fungicide Use**

Apply **Nile 70% EC** at a rate of 1% in 25 - 100 gals. of water per acre or 0.5 quart **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Apply no more than 2.5 gals. of **Nile 70% EC** per acre in solution per application. Repeat applications every 7 - 14 days.

Miticide Use

Use **Nile 70% EC** at a rate of 1% - 2% in 25 - 100 gals. of water per acre or 0.5 quart **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Apply no more than 2 gals. of **Nile 70% EC** per acre per solution. Apply at first sign of pests or pest damage. Repeat application as needed for adequate control every 7 - 21 days. **DO NOT** apply more than 5 gals. **Nile 70% EC** per acre, per application.

STONE AND TROPICAL FRUITS**Fungicide Use**

Apply **Nile 70% EC** at a rate of 1% in 25 - 200 gals. of water for high-volume applications or 0.5 gal. **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Reapply **Nile 70% EC** every 7 - 14 days until threat of disease is over.

Miticide Use

Use **Nile 70% EC** at first sign of pests or pest damage at the appropriate rate listed below as a foliar spray in sufficient water for thorough coverage. Repeat applications as needed for adequate control every 7 - 21 days. Apply a 1% - 2% solution of **Nile 70% EC** for high-volume application or a minimum of 1 quart **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. **DO NOT** apply more than 5 gals. of **Nile 70% EC** per acre, per application.

DO NOT use **Nile 70% EC** after pit hardening on stone fruit.

CROPS (including but not limited to)			
Apples	Coconuts*	Loquats (nispero)	Pears
Apricots	Crabapples	Lychee	Oriental Pears
Avocados	Dates	Mangoes	Pineapples
Bananas	Feijoa	Mandarins	Plantains
Calamondin	Figs	Satsuma Mandarins	Plums
Cherries	Grapefruits	Nectarines	Pomegranates
Barbados Cherries (Acerola)*	Guavas	Olives	Prunes
Sour Cherries	Jujubes*	Oranges	Pummelos
Sweet Cherries	Kiwifruit	Papayas	Quinces
Cherimoya	Kumquats	Passion Fruit	Tangelos
Citrus Citron	Lemons	Peaches	Tangerines
Citrus Hybrids	Limes		

*Not Registered for Use By California.

CUCURBITS

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Balsam Pears (Bitter Melons)	Crenshaw	Melons (Incl. Hybrids)	Pumpkins
Cantaloupes	Cucumbers	Honeydew Melons	Squashes
Casaba	Gherkins	Muskmelons	Watermelons (Incl. Hybrids)
Chinese Wax Gourds	Gourds	Persian Melons	Zucchini
Citron Melons	Honey Balls		

HERBS AND SPICES

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Anise	Chives	Marjoram	Sage
Balm	Coriander	Mint	Savory
Basil	Cumin	Nasturtium	Spearmint
Borage	Curry Leaf	Pennyroyal	Sweet Bay
Chamomile	Dandelion	Peppermint	Tarragon
Caraway	Dill	Rosemary	Thyme
Catnip	Fennel	Rue	Wintergreen
Celery	Marigold		

LEGUME AND FRUITING VEGETABLES

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Beans	Ground Cherries	Peanuts	Snow Peas
Broad Beans	Guar	Pepinos	Soybeans
Calabaza	Lentils	Peppers	Tomatillos
Chickpeas	Peas	Pigeon Peas	Tomatoes
Eggplants			

NUTS

Fungicide Use

Apply **Nile 70% EC** at a rate of 1% in 25 - 100 gals. of water per acre or 0.5 quart **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Apply no more than 2.5 gals. of **Nile 70% EC** per acre in solution per application. Repeat applications every 7 - 14 days.

Miticide Use

Use **Nile 70% EC** at a rate of 1% - 2% in 25 - 100 gals. of water per acre or 0.5 quart **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Apply no more than 2 gals. of **Nile 70% EC** per acre per solution. Apply at first sign of pests or pest damage. Repeat application as needed for adequate control every 7 - 21 days. **DO NOT** apply more than 5 gals. **Nile 70% EC** per acre, per application.

Walnuts: Avoid spraying after husk split.

CROPS (including but not limited to)			
Almonds	Cashews	Hickory Nuts	Pecans
Beech Nuts	Chestnuts	Japanese Horse Chestnuts	Pistachios
Brazil Nuts	Chinquapins	Macadamia Nuts	Walnuts
Butternuts	Filberts		

ROOT AND TUBER VEGETABLES

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Artichokes	Ginger	Sweet Potatoes	Turmeric
Beets	Ginseng	Radishes	Turnips
Cardone*	Horseradish	Rutabaga	Yams
Carrots	Parsnips	Taro (Dasheen)*	Yam Beans*
Cassava*	Potatoes		

*Not Registered for Use By California.

SMALL FRUITS AND BERRIES

Refer to specific application instructions found directly below the heading **AGRICULTURAL USE INSTRUCTIONS**.

CROPS (including but not limited to)			
Blackberries	Currants	Grapes	Raspberries (Black and Red)
Blueberries	Dewberries*	Huckleberries	Strawberries
Boysenberries	Elderberries	Loganberries	Youngberries
Cranberries	Gooseberries	Olallie Berries*	

*Not Registered for Use By California.

VEGETABLE, CEREAL GRAINS, AND OTHER MISCELLANEOUS CROPS

Fungicide Use

Apply **Nile 70% EC** at the rate of 0.5% - 1% in 25 - 100 gals. of water per acre or 2 pints **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume applications. Repeat applications every 7 - 14 days. Apply no more than 2 gals. of **Nile 70% EC** per acre per application. Larger plants will require higher amounts of water to obtain good coverage. For mushrooms, apply **Nile 70% EC** at a rate of 0.5% - 2% in 25 - 100 gals. of water per 8,000 square feet. Repeat applications every 7 - 14 days.

Miticide Use

Apply **Nile 70% EC** at a rate of 0.5% - 2% in 25 - 100 gals. of water per acre or 2 pints **Nile 70% EC** in a minimum of 5 gals. of water per acre for low-volume application. Apply no more than 2 gals. **Nile 70% EC** per acre in solution per application. Apply in a solution of sufficient volume to achieve complete coverage of foliage. Reapply **Nile 70% EC** every 7 - 21 days as needed for adequate control. Larger plants will require higher amounts of water to obtain good coverage.

CROPS (including but not limited to)			
Acerola (Barbados cherries)*	Coriander	Mushrooms	Sesame
Alfalfa	Corn	Nispero	Sorghum
Arracacha (Apio)*	Popcorn	Oats	Soursop (Guanabanas)*
Asparagus	Sweetcorn	Okra	Star Apples*
Barley	Cotton	Quenepa (kinep)*	Sugarcane
Breadfruit	Hops	Rice	Sunflowers
Buckwheat	Jicama*	Safflower	Tobacco
Canola	Jojoba	Seagrapes*	Wheat
Coffee			

*Not Registered for Use By California.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original container. Store in a dry place, away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. **DO NOT** store below 40°F (4°C).

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

Rigid Plastic Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.

Rigid Plastic Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

NOTES

NOTES

NOTES

Nile 70% EC

**For Use on Vegetables, Fruits, Nuts, Melons,
Outdoor Ornamental Plants, and Agronomic Crops.
For Foliar Control of Fungal Disease and Insect Pests.**

Can be Used Up to Day of Harvest.

ACTIVE INGREDIENT:	WT. BY %
Clarified Hydrophobic Extract of Neem Oil*	70.0%
OTHER INGREDIENTS:	30.0%
TOTAL:	100.0%

*This product contains 5.46 lbs. of clarified hydrophobic extract of neem oil per US gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

**See label booklet for complete Precautionary Statements
and Directions For Use.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-102-83529 EPA Est. No. **AG 72159-GA-001; **MA** 83411-MN-001; **MC** 89332-GA-001; **SD** 39578-TX-001; **TX** 07401-TX-001**

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

ENVIRONMENTAL HAZARDS

This product is hazardous to fish and aquatic invertebrates.

For Terrestrial Uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. **DO NOT** apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original container. Store in a dry place, away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. **DO NOT** store below 40°F (4°C).

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.

Rigid Plastic Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Net Contents: 2.5 Gals.* ☐ 265 Gals.

* Unless alternate checked

OPEN HERE