

Tifton

Systemic Turf and Ornamental Fungicide

For the Prevention and Control of Turf Diseases and the Diseases of Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Groundcovers, plus Deciduous and Evergreen Trees and Shrubs.

ACTIVE INGREDIENT:		WT. BY %
Thiophanate-Methyl: (dimethyl 4,4'-o-phenylenebis[3-thioallophanate])	2.08%
OTHER INGREDIENTS:	97.92%
TOTAL:	100.00%

Tifton is a granular formulation that contains 0.0208 lb. active ingredient per pound of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-331

EPA Est. No. **CS** 70815-GA-001; **SC** 39578-TX-001;

MA 83411-MN-001; **MC** 89332-GA-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, Applicators, and Other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils) are required for applicators using hand held equipment

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or areas where surface water is present or to intertidal areas below the mean high-water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticides.

Intended for use only by certified applicators or those under their immediate supervision.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: DO NOT enter or allow entry into treated areas until granules and dust have settled unless wearing personal protective equipment listed on this label.

PRODUCT INFORMATION

Tifton is a broad-spectrum fungicide for use on a wide variety of turf and ornamental diseases. **Tifton** may be applied using ground based, hand-operated (push) or mechanical (power) equipment; **DO NOT** apply this product by hand or with hand-held rotary spreader (e.g., bellygrinder).

Use Precautions:

- This product may only be used by individuals/firms licensed or registered by the State to apply ornamental or turf pest control products.
- Use only properly calibrated application equipment when applying this product.
- Thoroughly clean all application equipment after each use.
- For application schedules appropriate to your area, contact your local State Extension Service for recommendations.

Use Restrictions:

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** use on turf being grown for sale or other commercial use as sod.
- **DO NOT** graze animals on treated turf.
- **DO NOT** feed clippings to livestock or poultry.

Maximum Application Rates and Minimum Retreatment Intervals:

Tifton is a granular formulation that contains 0.0208 lb. active ingredient per pound of product.

- **Residential or Public Areas** -
 - **DO NOT** apply more than 131 lbs. of **Tifton** (2.72 lbs. a.i.) per acre per application. Or no more than 3 lbs. of **Tifton** (0.06 lb. a.i.) per 1,000 sq. ft. per application.
 - **DO NOT** apply more than 523 lbs. of **Tifton** (10.88 lbs. a.i.) per acre per year. Or no more than 12 lbs. of **Tifton** (0.25 lb. a.i.) per 1,000 sq. ft. per year.
 - **DO NOT** apply more than 4 applications per year.
 - **Minimum Retreatment Interval:** 14 days
- **Golf Course Aprons, Greens, and Tees** -
 - **DO NOT** apply more than 392 lbs. of **Tifton** (8.15 lbs. a.i.) per acre per application. Or no more than 9 lbs. of **Tifton** (0.19 lb. a.i.) per 1,000 sq. ft. per application.
 - **DO NOT** apply more than 1,048 lbs. of **Tifton** (21.8 lbs. a.i.) per acre per year. Or no more than 24 lbs. of **Tifton** (0.5 lb. a.i.) per 1,000 sq. ft. per year.
 - **Minimum Retreatment Interval:** 14 days
- **Golf Course Fairways (except Florida)** -
 - **DO NOT** apply more than 262 lbs. of **Tifton** (5.45 lbs. a.i.) per acre per application. Or no more than 6 lbs. of **Tifton** (0.12 lb. a.i.) per 1,000 sq. ft. per application.
 - **DO NOT** apply more than 262 lbs. of **Tifton** (5.45 lbs. a.i.) per acre per year. Or no more than 6 lbs. of **Tifton** (0.12 lb. a.i.) per 1,000 sq. ft. per year.
 - **Minimum Retreatment Interval:** 14 days
- **Golf Course Fairways (Florida only)** -
 - **DO NOT** apply more than 131 lbs. of **Tifton** (2.72 lbs. a.i.) per acre per application. Or no more than 3 lbs. of **Tifton** (0.06 lb. a.i.) per 1,000 sq. ft. per application.
 - **DO NOT** apply more than 131 lbs. of **Tifton** (2.72 lbs. a.i.) per acre per year. Or no more than 3 lbs. of **Tifton** (0.06 lb. a.i.) per 1,000 sq. ft. per year.
 - **DO NOT** apply more than 1 application per year.
 - **Minimum Retreatment Interval:** 14 days

RESISTANCE MANAGEMENT

In order to prevent the development of tolerant strains of fungi, **Tifton** should only be used with other fungicides that use different modes of action. Sharda USA LLC does NOT recommend using **Tifton** with products containing the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can contribute to the development of disease tolerance. **Tifton** can remain useful for disease control as long as specified precautions are followed. A tolerant strain of fungi may be present if a **Tifton** treatment is found to be ineffective. Promptly contact your State Agricultural Experiment Station or State Cooperative Extension Service for disease identification and advice on other suitable fungicide or disease control strategies.

COMMERCIAL TURF APPLICATIONS

Tifton may be used on all cool and warm season grasses including Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures in Commercial Turf, Residential Turf, Public Turf (Parks, Athletic Fields, Schools, Day Care Centers, etc.), and Golf Courses (Greens, Tees, Fairways, and Aprons).

When used in accordance with the instructions in this label, **Tifton** is not phytotoxic to any of the above-mentioned grasses.

Follow additional **Use Precautions** and **Use Restrictions** provided under **PRODUCT INFORMATION** section of this label.

Application Instructions

Apply under dry or moist leaf/soil conditions. For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 - 0.25 inch of water. Treating aprons when treating golf greens is advised.

Refer to the following table for specific rates and timing depending on the disease being treated. When treating under severe disease pressure, use the highest specified rate listed below.

Diseases	Disease Specific Instructions	Application Rate of Tifton (Lbs./1,000 Sq. Ft.)
Dollar Spot (<i>Moellerodiscus</i> , <i>Lanzia</i> , <i>Sclerotinia</i>) Large Brown Patch (<i>Rhizoctonia solani</i>) Ascochyta Leaf Blight (<i>Ascochyta</i> spp.) Copper Spot (<i>Gloeocercospora sorghi</i>) Red Thread (<i>Laetisaria fuciformis</i>) Fusarium Patch (<i>Fusarium nivale</i>) Zoysia Patch (<i>Rhizoctonia solani</i>)	For Prevention: Apply prior to the expected development of the disease using the lowest rate listed. For Treatment: Apply when disease first appears and repeat applications every 14-days. Note: May be rotated with chlorothalonil, iprodione, or mancozeb.	1.5 - 6 lbs. (0.03 - 0.12 lb. a.i.)
Pink Snow Mold (<i>Microdochium nivale</i>)	Make first application prior to cessation of turf growth activity in middle-late fall or early winter, or in spring during periods of thaw, or when disease symptoms first appear. Throughout the expected disease period, continue making applications at 14-day intervals. For best results, make the first application before the turf is completely dormant.	3 - 6 lbs. (0.06 - 0.12 lb. a.i.)
Anthraxnose, Foliar (<i>Colletotrichum graminicola</i>) Anthraxnose, Basal (<i>Colletotrichum graminicola</i>)	For Prevention: Once soil temperature reaches 60°F, make 2 applications at 14-day intervals. For Treatment: Apply at first sign of disease and continue applications at 14-day intervals. Note: May be rotated with chlorothalonil or triadimefon.	3 - 6 lbs. (0.06 - 0.12 lb. a.i.) 6 - 9 lbs. (0.12 - 0.19 lb. a.i.)
Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) Take-All-Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	Apply in mid-July or when disease symptoms first appear and repeat at 14-day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor.	
Cool Season Brown Patch (<i>Rhizoctonia cerealis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrae</i>) Spring Dead Spot (<i>Leptosphaeria korrae</i>)	For Prevention: Apply in the fall prior to cessation of turf growth. Make a second application in early spring when disease first appears or when soil temperatures reach 55°F - 60°F, whichever occurs first. For Treatment: Apply in early spring when disease first appears and repeat applications every 14-days.	
Coprinus Snow Mold (<i>Coprinus psychromorbidus</i>)	Make 2 treatments at 21-day intervals in late fall to early winter. Make the last application just prior to first permanent snow cover.	
Fusarium Blight (<i>Fusarium roseum</i> and <i>F. triticum</i>)	Apply when disease first appears. Make repeat applications at 14-day intervals.	
Gray Leaf Spot (Blast) (<i>Pyricularia grisea</i>)	Make a preventative application prior to the expected onset of disease development. Continue applications at 14-day intervals as needed.	
Leaf Spot (<i>Drechslera</i> spp.) Leaf, Crown, and Root Diseases (<i>Bipolaris</i> , <i>Curvularia</i> , and <i>Exserohilum</i>)	Apply when disease first appears and make applications at 14-day intervals as needed. Note: For severe cases, rotations with chlorothalonil, iprodione, or mancozeb are advised.	6 - 9 lbs. (0.12 - 0.19 lb. a.i.)
Stripe Smut (<i>Ustilago striiformis</i>)	For Prevention: Apply in spring and fall. For Treatment: Make 2 applications, the first when the disease first appears and the second 14-days later.	
Summer Patch (<i>Magnaporthe poae</i>)	For Prevention: Using a 21-day reapplication interval, make 3 applications starting in late April or early May. For Treatment: Apply at 14-day intervals when disease first appears. Rotations may be used as part of the 3-application program.	
Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	Apply in early June or when advised by your local Agricultural Extension Service. Reapply at 14-day intervals as needed. For season-long disease prevention, rotations and/or tank mix combinations may be used.	

COMMERCIAL HORTICULTURAL APPLICATIONS

Tifton is a broad-spectrum systemic fungicide for control of soil borne diseases affecting bulbs, corms, rhizomes, roots, crowns, and stems caused by *Botrytis*, *Cylindrocladium*, *Fusarium*, *Gliocladium*, *Myrothecium*, *Penicillium*, *Ramularia*, *Rhizoctonia*, *Sclerotinia*, and *Thielaviopsis*.

Tifton may be used on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites.

Note: Because **Tifton** does not control *Pythium* or *Phytophthora*, rotating with metalaxyl, mephenoxam, etridiazole, propamocarb, mono, and dipotassium salts of phosphorous acid or fosetyl-AI is required to control *Pythium* and *Phytophthora*.

Restrictions:

- **DO NOT** apply to home orchards/fruit trees after fruit set.
- **DO NOT** use fruit or nuts from treated trees for food or feed.
- **DO NOT** use on Swedish Ivy (*Nephtrolepis exaltata*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).
- For cut flowers, **DO NOT** exceed the maximum application rate of 24 lbs. of **Tifton** (0.5 lb. a.i.) per acre per application.
- For ground applications, **DO NOT** exceed 300 pounds active ingredient from ALL thiofanate-methyl containing products per acre per crop season.

Precaution:

- Because it is impossible to test this product on all species and cultivars, perform a preliminary trial on a small scale before a full treatment is applied to any plant type not shown on this label. When conducting the trial, wait 5 - 7 days after treatment to evaluate results.

Follow additional **Use Precautions** and **Use Restrictions** provided under **PRODUCT INFORMATION** section of this label.

Application Instructions

Tifton may be applied as a surface, fly-on, broadcast or over-the-top application, or media incorporated.

- **Surface and Broadcast Applications to Established Plants:** Apply 6 lbs. of **Tifton** per 1,000 sq. feet. Apply under dry or moist leaf/soil conditions. For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 - 0.25 inch of water.
- **Media Incorporation:** Mix at a rate of 2 - 3 lbs. per cubic yards before seeding or transplanting of desired crop. If not activated by the application of moisture, media incorporated with **Tifton** may be stockpiled until needed for later use.
- **For Preventive Control:** Prior to planting, apply as a media incorporation using the rate listed above. After seeding or transplant to propagation beds, containers, pots, or nursery or landscape beds, apply as a broadcast, fly-on, surface or over-the-top application using the **Surface and Broadcast Applications** instructions above.
- **For Treatment:** When disease first appears after seeding or transplanting, apply using the **Surface and Broadcast Applications** instructions above. To maintain protection, make repeat applications every 21 - 28 days.

SPREADER SETTINGS (Lbs. of product per 1,000 Sq. Ft.)			
DROP SPREADERS	1.5 lbs. (Prevention)	3 lbs. (Treatment)	
Republic EZ #530	1.5 lbs. (0.03 lb. a.i.)	2.5 lbs. (0.05 lb. a.i.)	
Scotts AccuGreen	3 lbs. (0.06 lb. a.i.)	5 lbs. (0.10 lb. a.i.)	
Sears Drop	3.5 lbs. (0.07 lb. a.i.)	5 lbs. (0.10 lb. a.i.)	
ROTARY SPREADERS	1.5 lbs. (Prevention)	3 lbs. (Treatment)	
Earthway	7 lbs. (0.15 lb. a.i.)	9 lbs. (0.19 lb. a.i.)	
Republic EZ #540	1.5 lbs. (0.03 lb. a.i.)	3 lbs. (0.06 lb. a.i.)	
Scotts SpeedyGreen	3.5 lbs. (0.07 lb. a.i.)	6.5 lbs. (0.14 lb. a.i.)	
Sears Rotary	1.5 lbs. (0.03 lb. a.i.)	3 lbs. (0.06 lb. a.i.)	

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable Outer Bag; Do not reuse or refill the outer bag. Completely empty bag into application equipment. Offer bag for recycling if available. If recycling is not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Plastic Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Nonrefillable Plastic Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom, and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

THIOPHANATE-METHYL GROUP 1 FUNGICIDE

Tifton

Systemic Turf and Ornamental Fungicide

For the Prevention and Control of Turf Diseases and the Diseases of Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Groundcovers, plus Deciduous and Evergreen Trees and Shrubs.

ACTIVE INGREDIENT:	WT. BY %
Thiophanate-Methyl: (dimethyl 4,4'-o-phenylenebis[3-thioallophanate])	2.08%
OTHER INGREDIENTS:	97.92%
TOTAL:	100.00%

Tifton is a granular formulation that contains 0.0208 lb. active ingredient per pound of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu.</p>	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or areas where surface water is present or to intertidal areas below the mean high-water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticides. Intended for use only by certified applicators or those under their immediate supervision.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING:**

Nonrefillable Outer Bag: Do not reuse or refill the outer bag. Completely empty bag into application equipment. Offer bag for recycling if available. If recycling is not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Plastic Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

See inside for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-331

EPA Est. No. **CS** 70815-GA-001; **SC** 39578-TX-001; **MA** 83411-MN-001; **MC** 89332-GA-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 7.5 Lbs.* 25 Lbs.

* Unless alternate checked

OPEN HERE