

HELP 1/2020

REFERENZEN

Sepsis 4.0: Früherkennung und Behandlung im Notfallzentrum **med. pract. Daniel Knuchel, Dr. med. Christiane Rosin**

1. Rhodes, A., et al., *Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016*. Intensive care medicine, 2017. **43**(3): p. 304-377.
2. Singer, M. and M. Shankar-Hari, *qSOFA, cue confusion*. Annals of internal medicine, 2018. **168**(4): p. 293-295.
3. Daniels, R., *The Sepsis Manual*. Vol. 4. 2017-2018, London: UK Sepsis Trust.
4. Rodriguez, R.M., et al., *Comparison of qSOFA with current emergency department tools for screening of patients with sepsis for critical illness*. Emerg Med J, 2018. **35**(6): p. 350-356.
5. Ortega, R.N., et al., *Clinical Scores and Formal Triage for Screening of Sepsis and Adverse Outcomes on Arrival in an Emergency Department All-Comer Cohort*. The Journal of emergency medicine, 2019.
6. Levy, M.M., L.E. Evans, and A. Rhodes, *The surviving sepsis campaign bundle: 2018 update*. Intensive care medicine, 2018. **44**(6): p. 925-928.
7. Inada-Kim, M. and E. Nsutebu, *NEWS 2: an opportunity to standardise the management of deterioration and sepsis*. Bmj, 2018. **360**: p. k1260.
8. Grossmann, F.F., et al., *Transporting clinical tools to new settings: cultural adaptation and validation of the Emergency Severity Index in German*. Annals of emergency medicine, 2011. **57**(3): p. 257-264.

Sepsis beim Neugeborenen oder Säugling auf der Notfallstation **Dr. med. Julia Höffe**

Hanna, Greenes: *How Much Tachycardia in Infants Can Be Attributed to Fever?* Ann Emerg Med. 2004;43: 699-705.

The UK Sepsis Trust Pediatric Sepsis 6 (version 11.1 August 2015)

Children's Hospital of Philadelphia Sepsis Pathway (www.chop.edu)

Sepsis beim geriatrischen Notfallpatienten **Thomas Dreher-Hummel, Pflegeexperte APN**

- Angus, D. C., Linde-Zwirble, W. T., Lidicker, J., Clermont, G., Carcillo, J., & Pinsky, M. R. (2001). Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med*, 29(7), 1303-1310. doi: 10.1097/00003246-200107000-00002
- Burkett, E., Macdonald, S. P., Carpenter, C. R., Arendts, G., Hullick, C., Nagaraj, G., & Osborn, T. M. (2018). Sepsis in the older person: The ravages of time and bacteria. *Emerg Med Australas*, 30(2), 249-258. doi: 10.1111/1742-6723.12949

- Chassagne, P., Perol, M. B., Doucet, J., Trivalle, C., Menard, J. F., Manchon, N. D., . . . Bercoff, E. (1996). Is presentation of bacteremia in the elderly the same as in younger patients? *Am J Med*, 100(1), 65-70. doi: 10.1016/s0002-9343(96)90013-3
- de Groot, B., Stolwijk, F., Warmerdam, M., Lucke, J. A., Singh, G. K., Abbas, M., . . . Rijpsma, D. (2017). The most commonly used disease severity scores are inappropriate for risk stratification of older emergency department sepsis patients: an observational multi-centre study. *Scand J Trauma Resusc Emerg Med*, 25(1), 91. doi: 10.1186/s13049-017-0436-3
- Karakoumis, J., Nickel, C. H., Kirsch, M., Rohacek, M., Geigy, N., Muller, B., . . . Bingisser, R. (2015). Emergency Presentations With Nonspecific Complaints-the Burden of Morbidity and the Spectrum of Underlying Disease: Nonspecific Complaints and Underlying Disease. *Medicine (Baltimore)*, 94(26), e840. doi: 10.1097/md.0000000000000840
- Ko, S. Y., Esteve Cuevas, L. M., Willeboer, M., Ansems, A., Blomaard, L. C., Lucke, J. A., . . . de Groot, B. (2019). The association between intravenous fluid resuscitation and mortality in older emergency department patients with suspected infection. *Int J Emerg Med*, 12(1), 1. doi: 10.1186/s12245-018-0219-2
- Nasa, P., Juneja, D., Singh, O., Dang, R., & Arora, V. (2012). Severe sepsis and its impact on outcome in elderly and very elderly patients admitted in intensive care unit. *J Intensive Care Med*, 27(3), 179-183. doi: 10.1177/0885066610397116
- Perera, T., & Cortijo-Brown, A. (2016). Geriatric Resuscitation. *Emerg Med Clin North Am*, 34(3), 453-467. doi: 10.1016/j.emc.2016.04.002
- Poulsen, J. B., Moller, K., Kehlet, H., & Perner, A. (2009). Long-term physical outcome in patients with septic shock. *Acta Anaesthesiol Scand*, 53(6), 724-730. doi: 10.1111/j.1399-6576.2009.01921.x
- Rhodes, A., Evans, L. E., Alhazzani, W., Levy, M. M., Antonelli, M., Ferrer, R., . . . Dellinger, R. P. (2017). Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Med*, 43(3), 304-377. doi: 10.1007/s00134-017-4683-6
- Sehgal, V., Bajwa, S. J., Consalvo, J. A., & Bajaj, A. (2015). Clinical conundrums in management of sepsis in the elderly. *J Transl Int Med*, 3(3), 106-112. doi: 10.1515/jtim-2015-0010
- Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., . . . Angus, D. C. (2016). The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, 315(8), 801-810. doi: 10.1001/jama.2016.0287
- Warmerdam, M., Stolwijk, F., Boogert, A., Sharma, M., Tetteroo, L., Lucke, J., . . . de Groot, B. (2017). Initial disease severity and quality of care of emergency department sepsis patients who are older or younger than 70 years of age. *PLoS One*, 12(9), e0185214. doi: 10.1371/journal.pone.0185214
- Waterer, G. W., Kessler, L. A., & Wunderink, R. G. (2006). Delayed administration of antibiotics and atypical presentation in community-acquired pneumonia. *Chest*, 130(1), 11-15. doi: 10.1378/chest.130.1.11
- Yende, S., Austin, S., Rhodes, A., Finfer, S., Opal, S., Thompson, T., . . . Angus, D. C. (2016). Long-Term Quality of Life Among Survivors of Severe Sepsis: Analyses of Two International Trials. *Crit Care Med*, 44(8), 1461-1467. doi: 10.1097/ccm.0000000000001658