



CHA₂DS₂-VASc Score

RISK FACTORS FOR STROKE

Risk Factors for Stroke & Criteria	Points
<u>C</u> ongestive heart failure (Right-sided, left-sided, or biventricular HF confirmed by noninvasive or invasive metrics)	1
<u>H</u> ypertension (BP of > 140 mm Hg systolic and/or > 90 mm Hg diastolic on at least 2 occasions or on treatment)	1
<u>A</u> ge ≥ 75 years	2
<u>D</u> iabetes (Fasting plasma glucose ≥ 126 mg/dL or on treatment)	1
<u>S</u> troke/TIA [^] /VTE [^] (DVT or PE)	2
<u>V</u> ascular disease (Known CAD, PVD or PAD, PCI of vasculature)	1
<u>A</u> ge 65-74 years	1
<u>S</u> ex category: Female	1
Maximum Score	9

CHA ₂ DS ₂ -VASc Score		Recommendation
Male	Female	
0	0-1	No anticoagulation
1	2	Consider anticoagulation*
≥ 2	≥ 3	Start anticoagulation

Other Risk Factors that Increase Risk of Stroke for Consideration	
Increased AF burden & duration	Poorly controlled HTN
Persistent/permanent AF	eGFR < 45 mL/min/1.73m ²
Obesity	Proteinuria > 150 mg/24 hr
Hypertrophic cardiomyopathy	Enlarged LA