



Auranofin Capsules

3 mg 60 Caps.

NDC: 85437-093-06

INDICATIONS:

Auranofin Capsules (auranofin) is indicated in the management of adults with active classical or definite rheumatoid arthritis (ARA criteria) who have had an insufficient therapeutic response to, or are intolerant of, an adequate trial of full doses of one or more nonsteroidal anti-inflammatory drugs. Auranofin Capsules should be added to a comprehensive baseline program, including non-drug therapies.

Unlike anti-inflammatory drugs, Auranofin Capsules does not produce an immediate response. Therapeutic effects may be seen after three to four months of treatment, although improvement has not been seen in some patients before six months.

When cartilage and bone damage has already occurred, gold cannot reverse structural damage to joints caused by previous disease. The greatest potential benefit occurs in patients with active synovitis, particularly in its early stage.

In controlled clinical trials comparing Auranofin Capsules with injectable gold, Auranofin Capsules was associated with fewer dropouts due to adverse reactions, while injectable gold was associated with fewer dropouts for inadequate or poor therapeutic effect. Physicians should consider these findings when deciding on the use of Auranofin Capsules in patients who are candidates for chrysotherapy.

For full Prescribing Information, see attached Product Insert.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit fda.gov/medwatch or call 1-800-FDA-1088.

Adjustable Dosing for Adults Managing Rheumatoid Arthritis

Dosing typically ranges between two and four applications daily based on symptom severity, helping patients overcome corticosteroid-responsive dermatoses.

Starting dose can be prescribed once or twice daily

Usual adult starting dose begins at 6 mg per day

Dose can be increased to 9 mg (three-times daily) after six months



COPAY INFORMATION

0-60 Caps » Up to \$100 of Copay
61-120 Caps » Up to \$150 of Copay
121 & Over » Up to \$200 of Copay

PROCESSOR: Drex
BIN #: 017290
PCN #: 55101202
GROUP #: E1280
ID #: 1001001
PERSON CODE: 01

Help Desk: 877-274-3244 *Terms and conditions may apply.

HOW TO WRITE:

Auranofin Capsules

Take 1 capsule by mouth twice daily.

3 mg

Dispense as written.



Visit our website at:
FLEXDOSING.COM

©2026 Flexdosing.com
All rights reserved.
LB-FD-01 Rev. 04/2026

Auranofin Capsules (auranofin) contains gold and, like other gold-containing drugs, can cause gold toxicity, signs of which include: fall in hemoglobin, leukopenia below 4,000 WBC/cu mm, granulocytes below 1,500/cu mm, decrease in platelets below 150,000/cu mm, proteinuria, hematuria, pruritus, rash, stomatitis or persistent diarrhea. Therefore, the results of recommended laboratory work (See PRECAUTIONS) should be reviewed before writing each Auranofin Capsules prescription. Like other gold preparations, Auranofin Capsules is only indicated for use in selected patients with active rheumatoid arthritis. Physicians planning to use Auranofin Capsules should be experienced with chrysotherapy and should thoroughly familiarize themselves with the toxicity and benefits of Auranofin Capsules.

In addition, the following precautions should be routinely employed:

1. The possibility of adverse reactions should be explained to patients before starting therapy.
2. Patients should be advised to report promptly any symptoms suggesting toxicity.

AURANOFIN- auranofin capsule
Coral Way Pharma, LLC

PRESCRIBING INFORMATION

Auranofin Capsules
Rx Only


Auranofin Capsules (auranofin) contains gold and, like other gold-containing drugs, can cause gold toxicity, signs of which include: fall in hemoglobin, leukopenia below 4,000 WBC/cu mm, granulocytes below 1,500/cu mm, decrease in platelets below 150,000/cu mm, proteinuria, hematuria, pruritus, rash, stomatitis or persistent diarrhea. Therefore, the results of recommended laboratory work (See PRECAUTIONS) should be reviewed before writing each Auranofin Capsules prescription. Like other gold preparations, Auranofin Capsules is only indicated for use in selected patients with active rheumatoid arthritis. Physicians planning to use Auranofin Capsules should be experienced with chrysotherapy and should thoroughly familiarize themselves with the toxicity and benefits of Auranofin Capsules.

In addition, the following precautions should be routinely employed:

1. The possibility of adverse reactions should be explained to patients before starting therapy.
2. Patients should be advised to report promptly any symptoms suggesting toxicity. (See PRECAUTIONS—Information for Patients.)

DESCRIPTION

Auranofin Capsules (auranofin) is available in oral form as capsules containing 3 mg auranofin. Auranofin is (2,3,4,6-tetra-O-acetyl-1-thio-β-D glucopyranosato-S-) (triethylphosphine) gold. Auranofin contains 29% gold and has the following chemical structure:



Each Auranofin Capsule, with opaque brown cap and opaque tan body, contains auranofin, 3 mg, and is imprinted with the product name RIDAURA. Inactive ingredients consist of benzyl alcohol, cellulose, cetylpyridinium chloride, D&C Red No. 33, FD&C Blue No. 1, FD&C Red No. 40, FD&C Yellow No. 6, gelatin, lactose, magnesium stearate, povidone, sodium lauryl sulfate, sodium starch glycolate, starch, titanium dioxide and trace amounts of other inactive ingredients.

CLINICAL PHARMACOLOGY

The mechanism of action of auranofin is not understood. In patients with adult rheumatoid arthritis, Auranofin Capsules may modify disease activity as manifested by synovitis and associated symptoms, and reflected

by laboratory parameters such as ESR. There is no substantial evidence, however, that gold-containing compounds induce remission of rheumatoid arthritis.

Pharmacokinetics

Pharmacokinetic studies were performed in rheumatoid arthritis patients, not in normal volunteers. Auranofin is rapidly metabolized and intact auranofin has never been detected in the blood. Thus, studies of the pharmacokinetics of auranofin have involved measurement of gold concentrations. Approximately 25% of the gold in auranofin is absorbed. The mean terminal plasma half-life of auranofin gold at steady state was 26 days (range 21 to 31 days; n=5). The mean terminal body half-life was 80 days (range 42 to 128; n=5). Approximately 60% of the absorbed gold (15% of the administered dose) from a single dose of auranofin is excreted in urine; the remainder is excreted in the feces. In clinical studies, steady state blood-gold concentrations are achieved in about three months. In patients on 6 mg auranofin/day, mean steady state blood-gold concentrations were 0.68 ± 0.45 mcg/mL (n=63 patients). In blood, approximately 40% of auranofin gold is associated with red cells, and 60% associated with serum proteins. In contrast, 99% of injectable gold is associated with serum proteins. Mean blood-gold concentrations are proportional to dose; however, no correlation between blood-gold concentrations and safety or efficacy has been established.

INDICATIONS AND USAGE

Auranofin Capsules (auranofin) is indicated in the management of adults with active classical or definite rheumatoid arthritis (ARA criteria) who have had an insufficient therapeutic response to, or are intolerant of, an adequate trial of full doses of one or more nonsteroidal anti-inflammatory drugs. Auranofin Capsules should be added to a comprehensive baseline program, including non-drug therapies. Unlike anti-inflammatory drugs, Auranofin Capsules does not produce an immediate response. Therapeutic effects may be seen after three to four months of treatment, although improvement has not been seen in some patients before six months.

When cartilage and bone damage has already occurred, gold cannot reverse structural damage to joints caused by previous disease. The greatest potential benefit occurs in patients with active synovitis, particularly in its early stage. In controlled clinical trials comparing Auranofin Capsules with injectable gold, Auranofin Capsules was associated with fewer dropouts due to adverse reactions, while injectable gold was associated with fewer dropouts for inadequate or poor therapeutic effect. Physicians should consider these findings when deciding on the use of Auranofin Capsules in patients who are candidates for chrysotherapy.

CONTRAINDICATIONS

Auranofin Capsules (auranofin) is contraindicated in patients with a history of any of the following gold-induced disorders: anaphylactic reactions, necrotizing enterocolitis, pulmonary fibrosis, exfoliative dermatitis, bone marrow aplasia or other severe hematologic disorders.

WARNINGS

Danger signs of possible gold toxicity include fall in hemoglobin, leukopenia below 4,000 WBC/cu mm, granulocytes below 1,500/cu mm, decrease in platelets below 150,000/cu mm, proteinuria, hematuria, pruritus, rash, stomatitis or persistent diarrhea.

Thrombocytopenia has occurred in 1–3% of patients (See ADVERSE REACTIONS) treated with Auranofin Capsules (auranofin), some of whom developed bleeding. The thrombocytopenia usually appears to be peripheral in origin and is usually reversible upon withdrawal of Auranofin Capsules. Its onset bears no relationship to the duration of Auranofin Capsules therapy and its course may be rapid. While patients' platelet counts should normally be monitored at least monthly (See PRECAUTIONS—Laboratory Tests), the occurrence of a precipitous decline in platelets or a platelet count less than 100,000/cu mm or signs and symptoms (e.g., purpura, ecchymoses or petechiae) suggestive of thrombocytopenia indicates a need to immediately withdraw Auranofin Capsules and other therapies with the potential to cause thrombocytopenia, and to obtain additional platelet counts. No additional Auranofin Capsules should be given unless the thrombocytopenia resolves and further studies show it was not due to gold therapy.

Proteinuria has developed in 3-9% of patients (See ADVERSE REACTIONS) treated with Auranofin Capsules. If clinically significant proteinuria or microscopic hematuria is found (See PRECAUTIONS—Laboratory Tests), Auranofin Capsules and other therapies with the potential to cause proteinuria or microscopic hematuria should be stopped immediately.

PRECAUTIONS

General

The safety of concomitant use of Auranofin Capsules (auranofin) with injectable gold, hydroxychloroquine, penicillamine, immunosuppressive agents (e.g., cyclophosphamide, azathioprine, or methotrexate) or high doses of corticosteroids has not been established. Medical problems that might affect the signs or symptoms used to detect Auranofin Capsules toxicity should be under control before starting Auranofin Capsules (auranofin).

The potential benefits of using Auranofin Capsules in patients with progressive renal disease, significant hepatocellular disease, inflammatory bowel disease, skin rash or history of bone marrow depression should be weighed against 1) the potential risks of gold toxicity on organ systems previously compromised or with decreased reserve, and 2) the difficulty in quickly detecting and correctly attributing the toxic effect. The following adverse reactions have been reported with the use of gold preparations and require modification of Auranofin Capsules treatment or additional monitoring. See ADVERSE REACTIONS for the approximate incidence of those reactions specifically reported with Auranofin Capsules.

Gastrointestinal Reactions

Gastrointestinal reactions reported with gold therapy include diarrhea/loose stools, nausea, vomiting, anorexia and abdominal cramps. The most common reaction to

Auranofin Capsules is diarrhea/ loose stools reported in approximately 50% of the patients. This is generally manageable by reducing the dosage (e.g., from 6 mg daily to 3 mg) and in only 6% of the patients is it necessary to discontinue Auranofin Capsules (auranofin) permanently. Ulcerative enterocolitis is a rare serious gold reaction. Therefore, patients with gastrointestinal symptoms should be monitored for the appearance of gastrointestinal bleeding.

Cutaneous Reactions

Dermatitis is the most common reaction to injectable gold therapy and the second most common reaction to Auranofin Capsules. Any eruption, especially if pruritic, that develops during treatment should be considered a gold reaction until proven otherwise. Pruritus often exists before dermatitis becomes apparent, and therefore should be considered to be a warning signal of a cutaneous reaction. Gold dermatitis may be aggravated by exposure to sunlight or an actinic rash may develop. The most serious form of cutaneous reaction reported with injectable gold is generalized exfoliative dermatitis.

Mucous Membrane Reactions

Stomatitis, another common gold reaction, may be manifested by shallow ulcers on the buccal membranes, on the borders of the tongue, and on the palate or in the pharynx. Stomatitis may occur as the only adverse reaction or with a dermatitis. Sometimes diffuse glossitis or gingivitis develops. A metallic taste may precede these oral mucous membrane reactions and should be considered a warning signal.

Renal Reactions

Gold can produce a nephrotic syndrome or glomerulitis with proteinuria and hematuria. These renal reactions are usually relatively mild and subside completely if recognized early and treatment is discontinued. They may become severe and chronic if treatment is continued after the onset of the reaction. Therefore it is important to perform urinalyses regularly and to discontinue treatment promptly if proteinuria or hematuria develops.

Hematologic Reactions

Blood dyscrasias including leukopenia, granulocytopenia, thrombocytopenia and aplastic anemia have all been reported as reactions to injectable gold and Auranofin Capsules. These reactions may occur separately or in combination at anytime during treatment. Because they have potentially serious consequences, blood dyscrasias should be constantly watched for through regular monitoring (at least monthly) of the formed elements of the blood throughout treatment.

Miscellaneous Reactions

Rare reactions attributed to gold include cholestatic jaundice; gold bronchitis and interstitial pneumonitis and fibrosis; peripheral neuropathy; partial or complete hair loss; fever.

Information for Patients

Patients should be advised of the possibility of toxicity from Auranofin Capsules and of the signs and symptoms that they should report promptly. (Patient information sheets are available.)

Women of childbearing potential should be warned of the potential risks of Auranofin Capsules therapy during pregnancy (See PRECAUTIONS—Pregnancy).

Laboratory Tests

CBC with differential, platelet count, urinalysis, and renal and liver function tests should be performed prior to Auranofin Capsules (auranofin) therapy to establish a baseline and to identify any preexisting conditions.

CBC with differential, platelet count and urinalysis should then be monitored at least monthly; other parameters should be monitored as appropriate.

Drug Interactions

In a single patient-report, there is the suggestion that concurrent administration of Auranofin Capsules and phenytoin may have increased phenytoin blood levels.

Carcinogenesis/Mutagenesis

In a 24-month study in rats, animals treated with auranofin at 0.4, 1.0 or 2.5 mg/kg/day orally (3, 8 or 21 times the human dose) or gold sodium thiomalate at 2 or 6 mg/kg injected twice weekly (4 or 12 times the human dose) were compared to untreated control animals.

There was a significant increase in the frequency of renal tubular cell karyomegaly and cytomegaly and renal adenoma in the animals treated with 1.0 or 2.5 mg/kg/day of auranofin and 2 or 6 mg/kg twice weekly of gold sodium thiomalate. Malignant renal epithelial tumors were seen in the 1.0 mg/kg/day and the 2.5 mg/kg/day auranofin and in the 6 mg/kg twice weekly gold sodium thiomalate-treated animals.

In a 12-month study, rats treated with auranofin at 23 mg/kg/day (192 times the human dose) developed tumors of the renal tubular epithelium, whereas those treated with 3.6 mg/kg/day (30 times the human dose) did not.

In an 18-month study in mice given oral auranofin at doses of 1, 3 and 9 mg/kg/day (8, 24 and 72 times the human dose), there was no statistically significant increase above controls in the instances of tumors. In the mouse lymphoma forward mutation assay, auranofin at high concentrations (313 to 700 ng/mL) induced increases in the mutation frequencies in the presence of a rat liver microsomal preparation. Auranofin produced no mutation effects in the Ames test (Salmonella), in the in vitro assay (Forward and Reverse Mutation Inducement Assay with *Saccharomyces*), in the in vitro transformation of BALB/T3 cell mouse assay or in the Dominant Lethal Assay.

Pregnancy

Teratogenic Effects

Pregnancy Category C

Use of Auranofin Capsules (auranofin) by pregnant women is not recommended. Furthermore, women of childbearing potential should be warned of the potential risks of Auranofin Capsules therapy during pregnancy. (See below.)

Pregnant rabbits given auranofin at doses of 0.5, 3 or 6 mg/kg/day (4.2 to 50 times the human dose) had impaired

food intake, decreased maternal weights, decreased fetal weights and an increase above controls in the incidence of resorptions, abortions and congenital abnormalities, mainly abdominal defects such as gastroschisis and umbilical hernia. Pregnant rats given auranofin at a dose of 5 mg/kg/day (42 times the human dose) had an increase above controls in the incidence of resorptions and a decrease in litter size and weight linked to maternal toxicity. No such effects were found in rats given 2.5 mg/kg/day (21 times the human dose). Pregnant mice given auranofin at a dose of 5 mg/kg/day (42 times the human dose) had no teratogenic effects.

There are no adequate and well-controlled RIDAURA studies in pregnant women.

Nursing Mothers

Nursing during Auranofin Capsules therapy is not recommended.

Following auranofin administration to rats and mice, gold is excreted in milk. Following the administration of injectable gold, gold appears in the milk of nursing women; human data on auranofin are not available.

Pediatric Use

Auranofin Capsules (auranofin) is not recommended for use in pediatric patients because its safety and effectiveness have not been established.

ADVERSE REACTIONS

The adverse reactions incidences listed below are based on observations of 1) 4,784 Auranofin Capsules treated patients in clinical trials (2,474 U.S., 2,310 foreign), of whom 2,729 were treated more than one year and 573 for more than three years; and 2) postmarketing experience. The highest incidence is during the first six months of treatment; however, reactions can occur after many months of therapy. With rare exceptions, all patients were on concomitant nonsteroidal anti-inflammatory therapy; some of them were also taking low dosages of corticosteroids.

Reactions occurring in more than 1% of Auranofin Capsules-treated patients

Gastrointestinal

loose stools or diarrhea (47%); abdominal pain (14%); nausea with or without vomiting (10%); constipation; anorexia*; flatulence*; dyspepsia*; dysgeusia.

Dermatological

rash (24%); pruritus (17%); hair loss; urticaria.

Mucous Membrane

stomatitis (13%); conjunctivitis*; glossitis.

Hematological

anemia; leukopenia; thrombocytopenia; eosinophilia

Renal

proteinuria*; hematuria.

Hepatic

elevated liver enzymes.

*Reactions marked with an asterisk occurred in 3-9% of the patients. The other reactions listed occurred in 1-3%.

Reactions occurring in less than 1% of Auranofin Capsules-treated patients

Gastrointestinal

dysphagia; gastrointestinal bleeding†; melena†; positive stool for occult blood†; ulcerative enterocolitis.

Dermatological

angioedema.

Mucous Membrane

gingivitis†.

Hematological

aplastic anemia; neutropenia†; agranulocytosis; pure red cell aplasia; pancytopenia.

Hepatic

jaundice.

Respiratory

interstitial pneumonitis.

Neurological

peripheral neuropathy.

Ocular

gold deposits in the lens or cornea unassociated clinically with eye disorders or visual impairment.

† Reactions marked with a dagger occurred in 0.1-1% of the patients. The other reactions listed occurred in less than 0.1%.

Reactions reported with injectable gold preparations, but not with Auranofin Capsules (auranofin) (based on clinical trials and on postmarketing experience)

Cutaneous Reactions

generalized exfoliative dermatitis

Capsules overdosage with modalities such as chelating agents. However, they have been used with injectable gold and may be considered for Auranofin Capsules overdosage.

DOSAGE AND ADMINISTRATION

Usual Adult Dosage:The usual adult dosage of Auranofin Capsules (auranofin) is 6 mg daily, given either as 3 mg twice daily or 6 mg once daily. Initiation of therapy at dosages exceeding 6 mg daily is not recommended because it is associated with an increased incidence of diarrhea. If response is inadequate after six months, an increase to 9 mg (3 mg three times daily) may be tolerated. If response remains inadequate after a threemonth trial of 9 mg daily, Auranofin therapy should be discontinued. Safety at dosages exceeding 9 mg daily has not been studied.

Transferring from Injectable Gold:

In controlled clinical studies, patients on injectable gold have been transferred to Auranofin Capsules (auranofin) by discontinuing the injectable agent and starting oral therapy with Auranofin Capsules, 6 mg daily. When patients are transferred to Auranofin Capsules, they should be informed of its adverse reaction profile, in particular the gastrointestinal reactions. (See PRECAUTIONS—Information for Patients.) At six months, control of disease activity of patients transferred to Auranofin Capsules and those maintained on the injectable agent was not different. Data beyond six months are not available.

HOW SUPPLIED

Capsules, containing 3 mg auranofin, in bottles of 60. NDC 85427-093-06

STORAGE AND HANDLING

Store between 15° and 30°C (59° and 86°F). Dispense in a tight, light-resistant container.

Distributed by:

Coral Way Pharma, LLC
San Antonio, TX 78213
1-210-641-5334

Revised September 2025

PI 09306001

Principal Display Panel – 3 mg Bottle Label

3 mg NDC 85437-093-06

Auranofin Capsules

60 Capsules

Incidence of Adverse Reactions for Specific Categories - 18 Comparative Trials

	Ridaura (445 patients)	Injectable Gold (445 patients)
Proteinuria	0.9%	5.4%
Rash	26%	39%
Diarrhea	42.5%	13%
Stomatitis	13%	18%
Anemia	3.1%	2.7%
Leukopenia	1.3%	2.2%
Thrombocytopenia	0.9%	2.2%
Elevated liver function tests	1.9%	1.7%
Pulmonary	0.2%	0.2%

OVERDOSAGE

The acute oral LD50 for auranofin is 310 mg/kg in adult mice and 265 mg/kg in adult rats. The minimum lethal dose in rats is 30 mg/kg.

In case of acute overdosage, immediate induction of emesis or gastric lavage and appropriate supportive therapy are recommended.

Auranofin Capsules overdosage experience is limited. A 50-year-old female, previously on 6 mg Auranofin Capsules daily, took 27 mg (9 capsules) daily for 10 days and developed an encephalopathy and peripheral neuropathy. Auranofin Capsules was discontinued and she eventually recovered.

There has been no experience with treating Auranofin