

Histex™-DM Syrup

ANTI HISTAMINE • DECONGESTANT
COUGH SUPPRESSANT

Alcohol Free • Sugar Free • Dye Free • Gluten Free

Active ingredients (in each 5 mL teaspoonful)	Purpose
Dextromethorphan HBr 20 mg	Cough Suppressant
Phenylephrine HCl 10 mg	Decongestant
Triprolidine HCl 2.5 mg	Antihistamine

NDC 28595-804-16

Antihistamine • Nasal Decongestant
Cough Suppressant

HISTEX™-DM Syrup

Each teaspoonful (5 mL) contains:
Dextromethorphan HBr ... 20 mg
Phenylephrine HCl 10 mg
Triprolidine HCl 2.5 mg

Sugar-Free • Dye Free
Alcohol Free

Grape Flavor

Tamper evident by foil seal under cap.
Do not use if foil seal is missing or broken.

16 fl oz (473 mL)

Directions

Do not exceed recommended dosage.

Age	Dose
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Histex™-DM Syrup

NDC: 28595-804-16

USES:

Temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat or bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

ACTIVE INGREDIENTS:

(in each 5 mL teaspoonful)

Dextromethorphan HBr 20 mg — Cough Suppressant
Phenylephrine HCl 10 mg — Decongestant
Triprolidine HCl 2.5 mg — Antihistamine

See Full Prescribing Information on back.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit fda.gov/medwatch or call 1-800-FDA-1088.

Histex™-DM Syrup

HISTEX-DM- dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride syrup
Allegis Pharmaceuticals, LLC

HISTEX™-DM Syrup Drug Facts

Uses

Temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat or bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a persistent or chronic cough such occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages

HOW TO WRITE:

Histex™-DM Syrup

Adults and children 12 years or older:
1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours.

Children 6 to under 12: ½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours.

Children under 6: Consult a doctor.

16 fl oz (473 mL)

Dispense as written.

- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed recommended dosage.

AGE

DOSE

Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Other Information

Store at 15°-30°C (59°-86°F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Dispense in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients

Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol, Grape Flavor.

Questions? Comments?

Call 1-866-633-9033.

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 28595-804-16

Antihistamine • Decongestant

Cough Suppressant

HISTEX™-DM

Syrup

Each teaspoonful (5 mL)

contains:

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Triprolidine HCl 2.5 mg

Sugar free • Dye Free

Alcohol Free

Grape Flavor

16 fl oz (473 mL)



Visit our website at:

FLEXDOSING.COM

©2026 Flexdosing.com

All rights reserved.

LB-FD-02 Rev. 05/2026