

Bubble Gum Flavor

NDC 28595-809-30

NEW FORMULATION  
**HISTEX™ PD**  
**Drops**

Antihistamine

Each dropperful (1 mL) contains:

Triprolidine HCl .... 1.25 mg

Sugar-Free • Dye Free

Alcohol Free

Bubble Gum Flavor

Tempor Evident: Do not use if seal is broken or missing

1 fl oz (30 mL)

New Formulation

# Histex™ PD Drops

Upper Respiratory Allergy Relief

Alcohol Free • Sugar Free • Dye Free • Gluten Free



Runny  
Nose



Sneezing



Itchy Nose  
or Throat



Itchy,  
Watery  
Eyes

## Directions

Do not exceed recommended dosage.

- use only with enclosed dropper
- do not use enclosed dropper with any other drug product

## Age

Adults & Children  
12 years of age or older:

Children 6 to under  
12 years of age:

Children under 6  
years of age:

## Dose

2 mL (2.5 mg) every 4 to 6 hours, not to exceed 8 mL (10 mg) in 24 hours, or as directed by a doctor.

1 mL (1.25 mg) every 4 to 6 hours, not to exceed 4 mL (5 mg) in 24 hours, or as directed by a doctor.

Consult a doctor.

## Histex™ PD Drops

NDC: 28595-809-30

**Triprolidine** is an antihistamine that relieves runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes

### USES:

Temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

### ACTIVE INGREDIENTS:

(in each 1 mL dropperful)  
Triprolidine HCl 1.25 mg—Antihistamine

See Full Prescribing Information on back.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [fda.gov/medwatch](http://fda.gov/medwatch) or call 1-800-FDA-1088.

# Histex™ PD Drops

HISTEX™ PD DROPS NEW FORMULATION-  
Triprolidine HCl  
Allegis Pharmaceuticals, LLC

-----  
**HISTEX™ PD New Formulation**  
Drug Facts

**Active ingredient (in each 1 mL dropperful)**  
Triprolidine HCl 1.25 mg

**Purpose**  
Antihistamine

**Uses**  
Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

**Warnings**  
Do not exceed recommended dosage.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

**Ask a doctor before use if you are taking sedatives or tranquilizers**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- sedatives and tranquilizers may increase the drowsiness effect

**Stop use and ask a doctor if**

- new symptoms occur

**Keep out of the reach of children.**

**In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**  
Do not exceed recommended dosage.

- use only the enclosed dropper.
- do not use enclosed dropper for any other drug products.

AGE	DOSE
Adults and Children 12 years of age and older:	2 mL (2.5 mg) every 4 to 6 hours, not to exceed 8 mL (10 mg) in 24 hours or as directed by a doctor.
Children 6 to under 12 years of age:	1 mL (1.25 mg) every 4 to 6 hours, not to exceed 4 mL (5 mg) in 24 hours or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

**Other Information**  
Store at 15°-30°C (59°-86°F).  
Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

**Inactive ingredients**  
Bubble Gum Flavor, Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions? Comments?**  
Call 1-866-633-9033.

**PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton**  
NDC 28595-809-30  
HISTEX™ PD New Formulation

**Drops**  
Antihistamine  
Each dropperful (1 mL) contains:  
Triprolidine HCl  
1.25 mg  
Sugar free • Dye Free  
Alcohol Free  
Bubble Gum Flavor  
1 fl oz (30 mL) Bottle

## PROFESSIONAL LABELING\*: For Professional Use Only

Age	Dose
Children 4 to under 6 years of age:	¾ dropperful (0.75 mL) every 4 to 6 hours, not to exceed 4doses (3.0 mL) in 24 hours, or as directed by a Doctor.
Children 2 to under 4 years of age:	½ dropperful (0.5 mL) every 4 to 6 hours, not to exceed 4 doses (2 mL) in 24 hours, or as directed by a Doctor.
Infants 4 months to under 2 years of age:	¼ dropperful (0.25 mL) every 4 to 6 hours, not to exceed 4 doses (1 mL) in 24 hours, or as directed by a Doctor.

\*These directions are intended for use by health professionals only and not the general public 21CFR Sec. 341.90 (q)



Visit our website at:  
**FLEXDOSING.COM**

©2026 Flexdosing.com  
All rights reserved.  
LB-FD-02 Rev. 05/2026