



Histex™ Syrup

Upper Respiratory Allergy Relief

Alcohol Free • Sugar Free • Dye Free • Gluten Free



**Runny
Nose**



Sneezing



**Itchy Nose
or Throat**



**Itchy,
Watery
Eyes**

Directions

Do not exceed recommended dosage.

Age	Dose
Adults & Children 12 years of age or older:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	½ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Histex™ Syrup

NDC: 28595-802-08

Triprolidine is an antihistamine that relieves runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes

USES:

Temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

ACTIVE INGREDIENTS:

(in each 5 mL teaspoonful)
Triprolidine HCl 2.5 mg—Antihistamine

See Full Prescribing
Information on back

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit fda.gov/medwatch or call 1-800-FDA-1088.

Histex Syrup

HISTEX- triprolidine hydrochloride syrup
Allergis Pharmaceuticals, LLC

HISTEX™ Syrup Drug Facts

Active ingredients (in each 5 mL teaspoonful)
Triprolidine HCl 2.5 mg

Purpose
Antihistamine

Uses
Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Warnings
Do not exceed recommended dosage.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

HOW TO WRITE:

Histex Syrup

Adults and children 12 years or older:
1 teaspoonful (5 mL) every 4 to 6 hours,
not to exceed 4 teaspoonfuls (20 mL) in
24 hours

Children 6 to under 12: ½ teaspoonful (2.5
mL) every 4 to 6 hours, not to exceed 2
teaspoonfuls (10 mL) in 24 hours.

Children under 6: Consult a Doctor

8 fl oz (237 mL)
Dispense as written.

Directions
Do not exceed recommended dosage.

AGE	DOSE
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	½ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Other Information
Store at 15°-30°C (59°-86°F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Dispense in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients
Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol, Bubble Gum Flavor.

Questions? Comments?
Call 1-866-633-9033.

PRINCIPAL DISPLAY PANEL - 237 mL Bottle Label
NDC 28595-802-08

Antihistamine
HISTEX™
Syrup
Each teaspoonful (5 mL)
contains:
Triprolidine HCl 2.5 mg
Sugar free • Dye Free

Alcohol Free
Bubble Gum Flavor



Visit our website at:
FLEXDOSING.COM

©2026 Flexdosing.com
All rights reserved.
LB-FD-01 Rev. 04/2026