CTNG Central Texas Neurology Consultants
Diseases of the Brain, Spine, Nerves and Muscles

Ocrevus (ocrelizumab) / Ocrevus Zunovo Infusion Order Set

Phone: (512) 218-1222
Please fax signed order, clinicals, and recent labs to:
Fax: (512) 218-1393

Patient Information

Patient Name:		Order Date:
DOB:	Ht:	Wt:
Allergies:		
Diagnosis:		ICD-10 Code:
Patient Status: ☐ New to Therapy ☐ Continuing Therapy		Next Treatment Date:

Lab Orders

☐ CBC w/diff and Platelets ☐ CMP
☐ HBsAg and anti-HBc ☐ Immunoglobulin
☐ UA ☐ Pregnancy Test☐ Other: _____
Frequency: ☐ Every Infusion ☐ Other:

Other Orders & Notes

☐ IV to PO premedication conversion per nurse discretion after tolerance of loading doses

Pre-Medication Orders

☐ Methylprednisolone 125mg IVP over 5 min; Administer 30 min prior to infusion
Diphenhydramine PO 25 mg ☐ or 50 mg ☐ PO 30-60 min prior; may repeat x1 in 4 hours.
Diphenhydramine IVP 25 mg ☐ or 50 mg ☐ IVP 30-60 min prior; may repeat x1 in 4 hours.
Cetirizine ☐ 10 mg PO or ☐ 10 mg IVP 30 minutes prior
☐ Famotidine 20 mg PO or ☐ 20 mg IVP 30 minutes prior
☐ Dexamethasone 20 mg PO 30 minutes prior
Acetaminophen ☐ 650 mg or ☐ 1000 mg PO 30 minutes prior; may repeat x1 in 4 hours.

Treatment Order

☐ Ocrevus (ocrelizumab) IV	300 mg/10 mL vial	☐ Ocrevus Initial Dose: Infuse 300 mg IV over at least 2.5 hours on weeks 0 and 2 ☐ Ocrevus Maintenance Dose: Infuse 600 mg IV every 6 months x 1 year
☐ Ocrevus Zunovo (ocrelizumab) SC	920 mg/23,000 units injection	☐ 920 mg SC injection every 6 months

Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN

Nursing

Skilled RN to establish and maintain vascular access as needed, draw ordered labs, conduct patient assessments, and educate on infusion services, medication administration, self-monitoring and safety

Ocrevus IV Infusion Titration (per PI):

- Initial 300 mg infusion: start at 30 mL/hour, increase by 30 mL/hour every 30 minutes, max 180 mL/hour
- 600 mg infusion - minimum 3.5 hr duration: start at 40 mL/hour, increase by 40 mL/hour every 30 minutes, max 200 mL/hour
- 600 mg infusion - minimum 2 hr duration: Begin infusion at 100 mL/hour for first 15 minutes; increase to 200 mL/hour for the next 15 minutes; increase to 250 mL/hour for the next 30 minutes; increase to 300 mL/hour for the remaining 60 minutes.

Ocrevus Zunovo SC Guidance:

- Administer as a single 920 mg SC injection in the abdomen or thigh
- Observe patient for at least 1 hour post-injection for hypersensitivity or injection-related reactions for the first injection, and ≥15 minutes following subsequent injections

Adverse Event (AE) Protocol

Epinephrine (based on patient weight)
EpiPen® Auto-Injector 0.3mg (1:1000) Inject IM or SubQ to patients who weigh ≥ 30 kg (66 lbs); may repeat in 3-5 minutes x1 if necessary.
EpiPen Jr® Auto-Injector 0.15mg (1:2000) Inject IM or SubQ for patients who weigh 15 - 30 kg (33-66lbs); may repeat in 3-5 minutes x1 if necessary.
Diphenhydramine 50mg (1 mL) If not previously administered, give 50mg slow IVP, or administer IM if no IV access; May repeat x1 after 10 minutes, if necessary.
Solumedrol 125 mg IVP
Sodium Chloride 0.9% 500mL Infuse IV at a rate of 30mL/hr.
Zofran 4mg ODT or IVP for nausea
Pepcid 20mg IV for urticaria/rash
Acetaminophen 325-650 mg PO x1 PRN pain if not already administered OR ibuprofen 200-400mg PO prn pain

Provider Information

By signing this form and utilizing our services, you are authorizing Nira Medical and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient.

Office Contact Name:	Email Address:
Prescriber Name:	Date:
Prescriber NPI:	Phone: Fax:
Prescriber Signature:	

CONFIDENTIAL: Contains protected health information (PHI). If received in error, notify the sender and destroy all copies.