CTNC Central Texas Neurology Consultants
Diseases of the Brain, Spine, Nerves and Muscles

Immunoglobulin Order Set

Phone: (512) 218-1222
Please fax signed order, clinicals, and recent labs to:
Fax: (512) 218-1393

Patient Information

Patient Name:		Order Date:
DOB:	Ht:	Wt:
Allergies:		
Diagnosis:		ICD-10 Code:
Patient Status: ☐ New to Therapy ☐ Continuing Therapy		Next Treatment Date:

Lab Orders

☐ CBC w/diff and Platelets ☐ CMP ☐ BMP
☐ HBsAg and anti-HBc ☐ UA
☐ Pregnancy Test
☐ Other_____

Frequency: ☐ Every Infusion
☐ Other:

Other Orders & Notes

Infuse 0.9% Sodium Chloride ☐ 100 mL or ☐ 500 mL IV over 1 hr following completion of infusion
☐ IV to PO premedication conversion per nurse discretion after tolerance of loading doses

Pre-Medication Orders

☐Methylprednisolone 125mg IVP over 5 min; Administer 30 min prior to infusion
Diphenhydramine PO 25 mg ☐ or 50 mg ☐ PO 30-60 min prior; may repeat x1 in 4 hours.

Diphenhydramine IVP 25 mg ☐ or 50 mg ☐ IVP 30-60 min prior; may repeat x1 in 4 hours.

Cetirizine ☐ 10 mg PO or ☐ 10 mg IVP 30 minutes prior

☐ Famotidine 20 mg IVP 30 minutes prior

☐ Dexamethasone 20 mg PO 30 minutes prior

Acetaminophen ☐ 650 mg or ☐ 1000 mg PO 30 minutes prior; may repeat x1 in 4 hours.

Treatment Order

Immunoglobulin ☐IV ☐SubQ	**Loading Dose:** Administer ___ GM/kg OR (___) Total Dose IV divided over ___ day(s) every ___ weeks OR ☐ one time dose **Maintenance Dose:** Administer ___ GM/kg OR (___) total dose IV divided over ___ day(s) every ___ weeks for 12 months If needed, round dose to nearest whole 5 gm vial for IV doses and nearest single-use vial size for subQ doses.

☐Infusion team to select brand
☐Do not substitute. Administer brand:

Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN

Nursing

Skilled RN to establish and maintain vascular access as needed, draw ordered labs, conduct patient assessments, and educate on infusion services, medication administration, self-monitoring and safety

Adverse Event (AE) Protocol

Epinephrine (based on patient weight)
EpiPen® Auto-Injector 0.3mg (1:1000) Inject IM or SubQ to patients who weigh ≥ 30 kg (66 lbs); may repeat in 3-5 minutes x1 if necessary.
EpiPen Jr® Auto-Injector 0.15mg (1:2000) Inject IM or SubQ for patients who weigh 15 - 30 kg (33-66lbs); may repeat in 3-5 minutes x1 if necessary.
Diphenhydramine 50mg (1 mL) If not previously administered, give 50mg slow IVP, or administer IM if no IV access; May repeat x1 after 10 minutes, if necessary.
Solumedrol 125 mg IVP
Sodium Chloride 0.9% 500mL Infuse IV at a rate of 30mL/hr.
Zofran 4mg ODT or IVP for nausea
Pepcid 20mg IV for urticaria/rash
Acetaminophen 325-650 mg PO x1 PRN pain if not already administered OR ibuprofen 200-400mg PO prn pain

Provider Information

By signing this form and utilizing our services, you are authorizing Nira Medical and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient.

Office Contact Name:	Email Address:
Prescriber Name:	Date:
Prescriber NPI:	Phone: Fax:
Prescriber Signature:	

CONFIDENTIAL: Contains protected health information (PHI). If received in error, notify the sender and destroy all copies.