



Same Trusted Medication with More Flexibility

- Oral Formulation
- Precise Titration

Baclofen Oral Solution 5 mg/5mL

NDC: 73352-103-16

INDICATIONS:

Baclofen oral solution is indicated for the treatment of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity.

Baclofen oral solution may also be of some value in patients with spinal cord injuries and other spinal cord diseases.

AVERSE REACTIONS:

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Adverse Reactions from Abrupt Withdrawal of Baclofen
- Neonatal Withdrawal Symptoms
- Drowsiness and Sedation
- Poor Tolerability in Stroke Patients
- Exacerbation of Psychotic Disorders, Schizophrenia, or Confusional States
- Exacerbation of Autonomic Dysreflexia
- Exacerbation of Epilepsy
- Posture and Balance Effects
- Ovarian Cysts

For full Prescribing Information, see attached Product Insert.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit fda.gov/medwatch or call 1-800-FDA-1088.

When Tablets Aren't An Option

Baclofen oral solution is an appropriate option for patients who are unable to swallow tablets.

- Patients with dysphagia
- Pediatric patients
- Elderly patients



More Precise Titration Than Fixed Tablets

When titration is required, an oral solution helps patients achieve the dose they need.

- Baclofen often requires gradual titration
- The liquid solution facilitates easier administration

COPAY INFORMATION

UP TO \$50 OFF*

BIN #: 610600

PCN #: AS

GROUP #: 349

ID #: 34905139472

Help Desk: 877-274-3244

*Terms and conditions may apply.

HOW TO WRITE:

Baclofen

Take 20 mL by mouth four times daily

5 mg/5 mL

Dispense as written



Visit our website at:
TRIFLUENTPHARMA.COM

©2026 Trifluent Parma, LLC.
All rights reserved.
LB-TRI-01 Rev. 04/2026