



GABAPENTIN CAPSULES



Bridges the Dosing Gap for Gabapentin Treatment

Gabapentin Capsules 200 mg

NDC: 73352-231-01

ADVERSE REACTIONS:

Most common adverse reactions (incidence $\geq 8\%$ and at least twice that for placebo) were:

- Postherpetic neuralgia: Dizziness, somnolence, and peripheral edema
- Epilepsy in patients >12 years of age: Somnolence, dizziness, ataxia, fatigue, and nystagmus
- Epilepsy in patients 3 to 12 years of age: Viral infection, fever, nausea and/or vomiting, somnolence, and hostility

INDICATIONS:

Gabapentin is indicated for:

- Management of postherpetic neuralgia in adults
- Adjunctive therapy in the treatment of partial onset seizures, with and without secondary generalization, in adults and pediatric patients 3 years and older with epilepsy

WARNINGS & PRECAUTIONS:

- Drug Reaction with Eosinophilia and Systemic Symptoms (Multiorgan hypersensitivity): Discontinue if alternative etiology is not established
- Anaphylaxis and Angioedema: Discontinue and evaluate patient immediately
- Driving Impairment; Somnolence/Sedation and Dizziness: Warn patients not to drive until they have gained sufficient experience to assess whether their ability to drive or operate heavy machinery will be impaired
- Suicidal Behavior and Ideation: Monitor for suicidal thoughts/behavior
- Abrupt or rapid discontinuation may increase the risk for seizures. Withdrawal symptoms, or suicidal behavior and ideation have been observed after discontinuation
- Respiratory Depression: May occur with Gabapentin when used with concomitant central nervous system (CNS) depressants, including opioids, or in the setting of underlying respiratory impairment. Monitor patients and adjust dosage as appropriate
- Neuropsychiatric Adverse Reactions in Children 3 to 12 Years of Age: Monitor for such events

A Flexible Dosing Option For Gabapentin Treatment

Dose Titration

Stepping up or down
between standard doses

Renal Impairment

Slower doses are required
based on creatinine clearance

Pediatric Dosing

Weight-based dosing may
call for smaller increments



COPAY INFORMATION

PATIENTS **\$0***
May Pay As Little As

*Full terms and conditions at trifluentpharma.com/copay

BIN #: 637765

PCN #: CRX

GROUP #: 349

ID #: 34905139472

Help Desk: 347-503-7025

HOW TO WRITE:

Gabapentin

Take 1 capsule by mouth
up to three times daily.

☐ 200 mg

Dispense as written.



Visit our website at:
TRIFLUENTPHARMA.COM

©2026 Trifluent Pharma, LLC.
All rights reserved.
LB-TRI-01 Rev. 09/2026