

Patient Enrollment Form

Once complete, submit by fax 1-877-633-9522 or email GoutHBYS@horizontherapeutics.com



Complete all required fields, including prescriber's signature and date, to initiate patient enrollment process.

For patient support and/or assistance obtaining patient signature, call Horizon By Your Side at 1-877-633-9521.

Patient Information (*Indicates a required field)

First name*	Last name*
Sex*: <input type="radio"/> Male <input type="radio"/> Female	Date of birth*: (MM/DD/YYYY)
Primary language	Email address
Consent to leave voice message at patient and/or alternate contact telephone? <input type="radio"/> Yes <input type="radio"/> No	
Primary telephone*	Consent to send text message? <input type="radio"/> Yes <input type="radio"/> No
<input type="radio"/> Home <input type="radio"/> Cell	
Address*	
City*	State* ZIP code*
Alternate contact name	Alternate contact telephone

Insurance Information (*Indicates a required field) (Please include front and back copies of insurance card[s] with this form)

Primary insurance*	Secondary insurance, if applicable
Policy #*	Policy #
Policyholder's first and last name*	Policyholder's first and last name
Insurance company telephone*	Insurance company telephone
Group #*	Group #
Policyholder's DOB*: (MM/DD/YYYY)	Policyholder's DOB: (MM/DD/YYYY)
IPA/Medical group name	IPA/Medical group telephone

☐ Reverification request

☐ Patient is uninsured to my knowledge

Infusion Facility (*Indicates a required field)

Do you have a preferred infusion facility?* ☐ Yes ☐ No If yes, please fill out the preferred infusion facility information below. If no, Horizon By Your Side will help identify a facility in close proximity to your patient.

☐ The infusion facility is the same as the prescribing office

Sage Infusion

Facility name*	
Facility address*	
City*	State* ZIP code*
Telephone* 1013578442	Fax* 84-1874532
Facility NPI #*	Facility tax ID #*

Patient Authorization (Required – please see authorization language on page 2)

	Date: (MM/DD/YYYY)
Patient signature Please read page 2	
Printed full name	

Please see Important Safety Information on page 2 and see Full Prescribing Information, including Boxed Warning, at KRYSTEXXAhcp.com.

P-KRY-US-00253 07/22

Prescriber Information (*Indicates a required field)

First name*	Last name*
Address*	
City*	State* ZIP code*
NPI #*	Tax ID #* State license #*
Clinic/hospital affiliation	
Office contact name	
Office contact telephone*	Fax*
Email address*	
Preferred communication: <input type="radio"/> Telephone <input type="radio"/> Email	Prescriber specialty*:
Referring healthcare provider: Was this patient referred to you by another HCP? <input type="radio"/> Yes <input type="radio"/> No If yes, please populate:	
Name:	Specialty:
City:	State:
ZIP code:	Telephone:

Diagnosis (Required for benefits investigation) (*Indicates a required field)

Primary diagnosis code*: M1A. — Chronic Gout
(Use coding wheel or see full list of codes at ChronicGoutCodes.com)

Additional disease manifestation codes:

Co-administration Medication

Is there an immunomodulator prescribed? ☐ Yes ☐ No If yes, please indicate below:

☐ methotrexate ☐ Other

Prescription Information (Required for specialty pharmacy benefit) (*Indicates a required field)

Dose: KRYSTEXXA® (pegloticase) injection, 8 mg/mL, for intravenous infusion every two weeks

Vial quantity*: Refills*:

Allergies*: or ☐ No known drug allergies (NKDA)

☐ Authorize administration supplies as needed

Contraindications:

- Patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency
- Patients with a history of serious hypersensitivity reactions, including anaphylaxis, to KRYSTEXXA or any of its components

Administration: The KRYSTEXXA admixture should only be administered by intravenous infusion over no less than 120 minutes via gravity feed, syringe-type pump, or infusion pump. Do not administer as an intravenous push or bolus. Please refer to the KRYSTEXXA Full Prescribing Information on preinfusion medications and how to reconstitute and dilute KRYSTEXXA for intravenous (IV) infusion.

State requirements: The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Noncompliance with state-specific requirements could result in outreach to the prescriber.

Prescriber Certification (Required – please see certification language on page 2)

	Substitutions allowed
Prescriber signature / Dispense as written*	Written or e-signature only; stamps not acceptable.
Date*: (MM/DD/YYYY)	
<input type="checkbox"/> I certify that the above therapy is medically necessary for the treatment of documented uncontrolled gout.* The above signature grants permission to share records with the referring office and infusion facility.	

Prescriber Certification

Please read and provide signature in Prescriber Certification section on page 1

I certify that the above therapy is medically necessary, that the information provided is accurate to the best of my knowledge and that my patient is being administered KRYSTEXXA® (pegloticase) injection, 8 mg/mL, for intravenous infusion in accordance with the labeled use of the product. I understand that Horizon Therapeutics USA, Inc. and its affiliates and their respective employees or agents (collectively, "Horizon") will use this information to administer the Horizon By Your Side program (the "Program"), which provides a wide array of patient-focused services, including providing logistical and non-medical treatment support for KRYSTEXXA, as prescribed, and educating about the insurance process. By my signature, I also certify that (1) my patient or his/her personal representative has provided a signed HIPAA authorization that allows me to share protected health information with Horizon for purposes of the Program and (2) I have obtained the patient's authorization to release such information as may be required for AllCare Plus Pharmacy (or another party acting on behalf of Horizon) to assess insurance coverage for KRYSTEXXA and assistance in initiating or continuing KRYSTEXXA as prescribed. I further understand and agree that (a) any medication or service provided through the Program as a result of this form is for the named patient only and is not being made in exchange for any express or implied agreement or understanding that I would recommend, prescribe, or use KRYSTEXXA® or any other Horizon product or service, for any other person; (b) my decision to prescribe KRYSTEXXA® was based solely on my professional determination of medical necessity; and (c) I will not seek reimbursement for any medication or service provided by or through the Program from any government program or third-party insurer. I understand that Horizon may modify or terminate the Program at any time without notice. The completion and submission of coverage- or reimbursement-related documentation are the responsibility of the patient and healthcare provider. Horizon makes no representation or guarantee concerning coverage or reimbursement for any item or service. On behalf of the patient, Horizon expects the prescriber to coordinate with Horizon By Your Side to provide, to the best of the prescriber's ability, in-network infusion services and work with Horizon By Your Side to effectively communicate both in-network and out-of-network choices and the corresponding financial obligations of the patient connected to each choice. Should the prescriber knowingly perform out-of-network services without the knowledge and consent of the patient, the prescriber cannot balance bill the patient for the out-of-network services.

State requirements: The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Noncompliance with state-specific requirements could result in outreach to the prescriber.

By filling out and signing this form, the enrollment process in Horizon By Your Side has initiated; however, your patient must sign a Patient Authorization to complete enrollment in Horizon By Your Side. Please note that your patient will not benefit from the services and support offered by the Program unless your patient signs a Patient Authorization, consenting to receiving such services. If your patient does not sign the Patient Authorization contained within this form, Horizon will contact the patient to determine whether the patient is interested in signing a separate Patient Authorization.

Patient Consent for Patient Information, Enrolling in Services, and Accessing Financial Support (referred to as "Patient Authorization")

Please read and provide signature in Patient Authorization section on page 1

I hereby authorize my healthcare providers, my health insurance carriers, and my pharmacies to use and disclose my individually identifiable health information, including my medical records, insurance coverage information, and my name, address, and telephone number to Horizon Therapeutics USA, Inc. and its affiliates and their respective agents and representatives (collectively, "Horizon"), including third parties authorized by Horizon to administer drug support and to dispense drugs (collectively, "Horizon By Your Side") for the following purposes: (1) to establish eligibility for benefits; (2) to communicate with my healthcare providers and me about my treatment or condition and related products; (3) to facilitate the provision of products, supplies, or services by a third party including, but not limited to, specialty pharmacies; (4) to register me in any applicable product registration program required for my treatment; (5) to enroll me in eligible patient support programs offered by Horizon By Your Side and/or Horizon, including nursing or patient access support services (government-reimbursed programs may not be eligible for all support services offered; please contact Horizon By Your Side for determination); and (6) to send me marketing information or offer me products and services related to my treatment or condition (or other products or services in which I might be interested) and to contact me occasionally to obtain my feedback (for market research purposes only) about my treatment, my condition, or my experience with Horizon and/or Horizon By Your Side otherwise as required or permitted by law. Further, I appoint the Program, on my behalf, to proceed with Program services and to convey this prescription to the dispensing pharmacy, to the extent permitted under state law. I understand the pharmacies may receive a fee from Horizon in exchange for (1) providing me with certain materials and information described above, and (2) using or disclosing certain health information pursuant to this Authorization.

I understand that Horizon, as well as my healthcare providers, cannot require me, as a condition of having access to medications, prescription drugs, treatment, or other care, to sign this Authorization. I understand that I am entitled to a copy of this Authorization.

I understand that information disclosed pursuant to this Authorization in some cases may be redisclosed by the recipient and no longer protected by HIPAA or other privacy laws. But Horizon has agreed to use and disclose my information only for purposes of operating the Program. I understand that I may cancel this Authorization at any time by mailing a signed letter requesting such cancellation to Horizon By Your Side, 1 Horizon Way, Deerfield, IL 60015, but that this cancellation will not apply to any information used or disclosed by my healthcare providers and/or health insurance carriers based on this Authorization before they are notified that I have cancelled it. Unless required by state law, this Authorization is valid for whichever is greater: (a) the duration remaining on this treatment or (b) 10 years from the date signed above. A photocopy of this Authorization will be treated in the same manner as the original.

INDICATION

KRYSTEXXA® (pegloticase) is indicated for the treatment of chronic gout in adult patients who have failed to normalize serum uric acid and whose signs and symptoms are inadequately controlled with xanthine oxidase inhibitors at the maximum medically appropriate dose or for whom these drugs are contraindicated.

Limitations of Use: KRYSTEXXA is not recommended for the treatment of asymptomatic hyperuricemia.

IMPORTANT SAFETY INFORMATION

WARNING: ANAPHYLAXIS AND INFUSION REACTIONS, G6PD DEFICIENCY ASSOCIATED HEMOLYSIS AND METHEMOGLOBINEMIA

- Anaphylaxis and infusion reactions have been reported to occur during and after administration of KRYSTEXXA.
- Anaphylaxis may occur with any infusion, including a first infusion, and generally manifests within 2 hours of the infusion. Delayed hypersensitivity reactions have also been reported.
- KRYSTEXXA should be administered in healthcare settings and by healthcare providers prepared to manage anaphylaxis and infusion reactions.
- Premedicate with antihistamines and corticosteroids and closely monitor for anaphylaxis for an appropriate period after administration of KRYSTEXXA.
- Monitor serum uric acid levels prior to each infusion and discontinue treatment if levels increase to above 6 mg/dL, particularly when 2 consecutive levels above 6 mg/dL are observed.
- Screen patients at risk for glucose-6-phosphate dehydrogenase (G6PD) deficiency prior to starting KRYSTEXXA. Hemolysis and methemoglobinemia have been reported with KRYSTEXXA in patients with G6PD deficiency. KRYSTEXXA is contraindicated in patients with G6PD deficiency.

CONTRAINDICATIONS:

- In patients with G6PD deficiency.
- In patients with history of serious hypersensitivity reactions, including anaphylaxis, to KRYSTEXXA or any of its components.

WARNINGS AND PRECAUTIONS

Gout Flares: An increase in gout flares is frequently observed upon initiation of anti-hyperuricemic therapy, including KRYSTEXXA. Gout flare prophylaxis with a non-steroidal anti-inflammatory drug (NSAID) or colchicine is recommended starting at least 1 week before initiation of KRYSTEXXA therapy and lasting at least 6 months, unless medically contraindicated or not tolerated.

Congestive Heart Failure: KRYSTEXXA has not been formally studied in patients with congestive heart failure, but some patients in the pre-marketing placebo-controlled clinical trials experienced exacerbation. Exercise caution in patients who have congestive heart failure and monitor patients closely following infusion.

ADVERSE REACTIONS

The most commonly reported adverse reactions (≥5%) are:

KRYSTEXXA co-administration with methotrexate trial:

KRYSTEXXA with methotrexate: gout flares, arthralgia, COVID-19, nausea, and fatigue; KRYSTEXXA alone: gout flares, arthralgia, COVID-19, nausea, fatigue, infusion reaction, pain in extremity, hypertension, and vomiting.

KRYSTEXXA pre-marketing placebo-controlled trials:

gout flares, infusion reactions, nausea, contusion or ecchymosis, nasopharyngitis, constipation, chest pain, anaphylaxis, and vomiting.

For additional information on KRYSTEXXA, please see Full Prescribing Information, including Boxed Warning, at [KRYSTEXXAhcp.com](https://www.krystexxa.com).



KRYSTEXXA and the HORIZON logo are trademarks owned by or licensed to Horizon.

All other trademarks are the property of their respective owners.

© 2022 Horizon Therapeutics plc P-KRY-US-00253 07/22

Krystexxa (Pegloticase) Infusion Orders

Patient Name:	DOB:	<input type="checkbox"/> Male <input type="checkbox"/> Female
Diagnosis (please provide ICD10 code)	Other:	
<input type="checkbox"/> NKDA Allergies:		
<input type="checkbox"/> New Start Therapy	<input type="checkbox"/> Continuation of Therapy	Date of last dose (if applicable):

Ordering Provider:

Provider NPI:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

PRE-MEDICATION

- ☒ Protocol: Benadryl 25 mg PO, Solumedrol 125mg IV, and Tylenol 1000mg PO prior to infusions (*alternative premeds can be administered if ordered by referring physician*)

☐ Other: _____

KRYSTEXXA ORDERS

- ☒ Dose: 8mg IV in 250 mL 0.9% Sodium Chloride

Administered over 2 hours, with 1 hour observation post infusion

Frequency: Every 2 weeks

** It is recommended that Krystexxa be co-administered with weekly methotrexate 15 mg orally. Krystexxa alone may be used in patients for whom methotrexate is contraindicated or not clinically appropriate.*

LABS:

- ☒ Uric Acid Level q 2 weeks

** UA Point of Care Testing will be completed at Noble Infusion prior to each infusion. It is recommended that treatment be discontinued if levels increase to above 6 mg/dL, particularly when 2 consecutive levels above 6 mg/dL are observed.*

Noble Infusion Standing Orders:

- ☒ Provide treatment under Noble Infusion's Clinical Guidelines, Medication Safety Protocol, Emergency Guidelines, and Action Plan for Infusion Reactions.

Provider Name

Provider Signature

Date