

## **Reclast (zoledronic acid) Infusion Orders**

Patient Name:	DOB:	·	☐ Male ☐ Female
Diagnosis (please provide ICD10 code)			
□ Other:			
☐ New Start Therapy ☐ Continuation of Ther	apy Date of last dose (if applicab	le): 🔲 NKDA	Allergies:
Oudaria a Brasida a			
Ordering Provider:			
Provider NPI:	Phone:	Fax:	
Practice Address:	City:	State:	: Zip Code:
PREMEDICATIONS:		REQUIRED TE	ESTING/LABS:
☐ Acetaminophen 1000mg PO ☐ Solu-Medrol 125mg IVP		_	ss Notes supporting primary
☐ Diphenhydramine 25mg PO ☐ Solu-Co	rtef 100mg IVP	diagnosis (plea  DEXA scan resu	ise attach) ilts and date (please attach):
☐ Ceterizine 10mg PO ☐ Diphenh	nydramine 25mg IVP	 ✓ Most recent CN	 MP lab results (please attach)
☐ Other:		_	ν,
RECLAST ORDERS			
DOSING:		REFILLS:	
☑ Reclast 5mg/100ml IV infusion over at least 15 minutes		No Refills	
FREQUENCY:			
M. Once annually			
☑ Once annually			
Noble Infusion Standing Orders:			
Provide treatment under Noble Infusion's Cli Guidelines, and Action Plan for Infusion Read	inical Guidelines, Medication Safe ctions.	ty Protocol, Emergency	
Provider Name			
Provider Signature		 Date	

Hypocalcemia may worsen during treatment. Patients must be adequately supplemented with calcium and vitamin D. Renal Impairment: Monitor creatinine clearance before each dose.