

Orencia (Abatacept) Infusion OrderPatient Name: _____ DOB: _____ ☐ Male ☐ Female

Diagnosis (please provide ICD10 code) _____

☐ Other: _____☐ NKDA Allergies: _____☐ New Start Therapy☐ Continuation of Therapy

Date of last dose (if applicable): _____

Ordering Provider: _____

Provider NPI: _____

Phone: _____

Fax: _____

Practice Address: _____

City: _____

State: _____

Zip Code: _____

PRE-MEDICATION

- ☐ Acetaminophen 1000mg PO ☐ Solu-Medrol 125mg IVP
☐ Diphenhydramine 25mg PO ☐ Solu-Cortef 100mg IVP
☐ Ceterizine 10mg PO ☐ Diphenhydramine 25mg IVP

REQUIRED LABS

- ☒ TB status and date (please attach results):

☒ Hepatitis B status & date (please attach results):

ORENCIA ORDERS**DOSING:**

- ☒ Mix in 100ml 0.9% sodium chloride and administer intravenous infusion over 30 minutes

Dose: ☐ 500mg ☐ 750mg ☐ 1000mg ☐ Other: _____ mg**FREQUENCY:**

- ☐ **Induction and Maintenance:** Week 0, 2, and 4, then every 4 weeks
☐ **OR Maintenance Only:** every 4 weeks
☐ Other: _____

REFILLS:☐ _____

(if not indicated prescription will expire one year from date signed)

Noble Infusion Standing Orders:

- ☒ Provide treatment under Noble Infusion's Clinical Guidelines, Medication Safety Protocol, Emergency Guidelines, and Action Plan for Infusion Reactions.

Provider Name_____
Provider Signature_____
Date

Serious infections reported. Patients with a history of recurrent infections or underlying conditions predisposing to infections may experience more infections. Discontinue if a serious infection develops. Screen for latent TB infection prior to initiating therapy. Patients testing positive should be treated prior to initiating Orencia. Should screen for viral hepatitis prior to initiating Orencia. Update vaccinations prior to initiating Orencia. Live vaccines should not be given concurrently or within 3 months of discontinuation. Orencia may blunt the effectiveness of some immunizations. COPD patients may develop more frequent respiratory adverse events.