



MacroSol Ca Acetate

Ca

DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events. Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a jar test before mixing. Do not use with amine based products. Do not apply this product with copper fungicides by foliar application on all crops.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

Calcium (Ca): 70 g/L as calcium acetate.

A high analysis calcium acetate solution, free of any chloride, nitrogen and sulfur, specifically designed for foliar and fertigation applications.

CROPS	RATE	COMMENTS
BROADACRE CROPS	Foliar 2 - 10 L/ha	Apply as required by the crop
VEGETABLE AND ROW CROPS	Foliar 5 - 8 L/ha Fertigation 20 - 50 L/ha	Apply 4 to 5 times over the growing season as required. Apply with a min 500 L/ha.
APPLES	Foliar 5 - 8 L/ha Fertigation 40 - 80 L/ha	Apply 4 to 5 as required based on tissue analysis. Apply with a min 500 L/ha.
PEAR	Foliar 2 - 4 L/ha Fertigation 40 - 80 L/ha	Apply 4 to 5 as required based on tissue analysis from Petal fall. Apply with a min 500 L/ha.
CHERRIES	Foliar 2 - 3 L/ha Fertigation 40 - 80 L/ha	Apply 4 to 5 as required based on tissue analysis from Petal fall. Apply with a min 500 L/ha.
NUT CROPS	Foliar 3 - 4 L/ha Fertigation 10 - 30 L/ha	Apply during periods of active growth to assist with high calcium demand.
VINES	Foliar 3 - 5 L/ha Fertigation 20 - 50 L/ha	Apply three times per season, ideally at bud burst, flowering and early fruit fill to assist with times of high calcium demand.

CAUTION: Do not apply as a foliar application in conjunction with copper and EC fungicides. Applications in adverse spraying conditions or in conjunctions with some crop protection products may result in reductions in efficacy or phytotoxicity which may not be acceptable to growers. For most effective results apply early morning or late afternoon.

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant.
Avoid inhalation and contact with the eyes and skin.

Batch No:

Date of Manufacture:

This product is NOT subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container

Specialist Advice In Emergency Only
1800 033 111
All Hours - Australia Wide