

Fruit Fly Control in Stonefruit

For the control of Queensland Fruit Fly and Mediterranean Fruit Fly on stone fruit

MAIN BENEFITS

- Fast Knockdown of QFF and MFF
- Short Witholding Period

APPLICATION

- Perfect Pre-Harvest clean up spray
- 50ml /100L, Maximum 3 applications per season
- 3 Day Witholding Period

ACTIVE INGREDIENT

- ACTIVE CONSTITUENT: 287.5 g/L ETOFENPROX
- Etofenprox is a non ester pyrethroid with GROUP 3A mode of action.

WHY USE TREBON?

- WHP 3 days vs 7 days for clothianidin.
- Should be used as part of a full orchard hygiene program that includes pest monitoring, lure & kill and mating disruption products.
- Rotate with other insecticides for resisitance management.
- Fast knock-down action plus residual activity.
- The perfect clean-up spray before harvest.
- Trebon is used in many other countries in a wide range of crops for control of various insect pests.
- Trebon is the leading product for Mediterranean Fruit Fly control in Europe.
- Trebon controls insects on contact and by ingestion, it has repellancy activity and inhibits egg laying.
- The mammalian toxicity of Etofenprox is very low.



Mediterranean Fruit Fly (Ceratitis capitata)



Queensland Fruit Fly (Bactrocera tryoni)


















CONTACTS FOR FURTHER INFORMATION Dean Smith (STH EAST AUS)0407 601 717 Steve Lacy (WA)......0408 125 679 Stewart Frankling (NORTHERN AUS)......0447 558 219

Queensland Fruit Fly (Bactrocera tryoni) Stanthorpe 1 QLD 2017 MEAN INCIDENCE TREE FRUIT **MEAN SEVERITY TREE FRUIT** Malathion 200 mL Trebon 50 mL Clothianidin 20gai







DIRECTIONS FOR USE:

CROP	INSECT PEST	RATE	WHP	CRITICAL COMMENTS
Stone fruit	Queensland Fruit	50-100	3 days	Spray to wet foliage and fruit to near the point of run-off
(except	Fly (Bactrocera	mL/100L		when adult fruit fly numbers warrant spraying. Trebon will not
cherries)	tryoni) and			control eggs or larvae present in fruit at spraying. Use the
	Mediterranean			higher rate for heavy infestations. If re-application is needed,
	Fruit Fly			a minimum of 7 days is required between treatments.
	(Ceratitis capitata)			Do not apply more than 3 applications per season.

Always read the entire label prior to use.

® Trebon is a registered trademark of Mitsui AgriScience International. For further information please call Sipcam or visit our website: sipcam.com.au

Phone: +61 3 5223 3746

