

Clothes, Sizing and Body Changes in Recovery

Body image, self-perception and eating-disorder recovery

Clothing labels are inconsistent, but they can become emotionally loaded. Tight, uncomfortable or “goal” clothes may turn getting dressed into a daily test and intensify eating-disorder thoughts.

What may be happening

- A clothing size is manufacturer information, not a measure of character or health.
- Bodies can change with nourishment, hormones, medication, age, disability, stress and recovery.
- Comfortable clothes can reduce body monitoring and free attention for living.

Your body does not need to shrink to deserve clothes that feel safe and comfortable.

A gentle reminder

Body-image distress can feel convincing and urgent. You do not need to solve your relationship with your body in one moment. The next helpful step may simply be to pause, reduce checking or comparison, eat according to your support plan, and tell someone that the thoughts are loud today.

Ways to respond with care

Try these small steps

- Put away items that punish, pinch or invite repeated checking.
- Choose clothes that fit your body today and allow breathing, sitting and eating.
- If shopping is difficult, plan support, limit fitting-room time and remove labels at home.

A short reflection

What triggered me? Think about the situation, conversation, image, sensation or emotion that came first.

What did the eating-disorder voice ask me to do? Name the urge without automatically following it.

What would protect my recovery? Choose one small action involving nourishment, rest, connection, boundaries or professional support.

Words you can borrow

- “Your body does not need to shrink to deserve clothes that feel safe and comfortable.”
- “A thought is not an instruction.”
- “My body deserves care even when my feelings are difficult.”

Use this resource alongside care

This guide is supportive information, not a diagnosis or a replacement for individual medical, nutritional or psychological care. Eating disorders can affect people at any body size and can become medically serious. If you are worried about your eating, compensatory behaviours or physical symptoms, contact your GP or eating-disorder team as soon as possible.

Support and next steps

You deserve support before things become “bad enough”. A GP can ask about eating and wellbeing, check physical health and refer to a specialist service when appropriate. You can take someone you trust to the appointment.

Seek urgent medical help now if you are

- fainting or close to fainting, confused, extremely weak or unable to stay awake
- experiencing chest pain, breathing difficulty, a very fast or irregular heartbeat
- unable to keep fluids down, severely dehydrated, or feeling at immediate risk
- thinking you may harm yourself or cannot keep yourself safe

In an emergency, call 999 or go to A&E.; For urgent mental-health help in England, call NHS 111 and select the mental-health option. Beat offers UK eating-disorder information and support; its helpline is 0808 801 0677. Availability can change, so check the organisation’s website.

Trusted information

NHS: Eating disorders overview -

www.nhs.uk/mental-health/feelings-symptoms-behaviours/behaviours/eating-disorders/overview/

NICE guideline NG69: Eating disorders - recognition and treatment - www.nice.org.uk/guidance/ng69

Beat: www.beateatingdisorders.org.uk

Until next time, give yourself space to breathe.

Maggie Learoyd
Space to Breathe Therapy