

To Fill Boredom, Emptiness or Lack of Purpose

When the behaviour provides stimulation, routine or something to look forward to

People rarely turn to substances or addictive behaviours for no reason. The behaviour may begin as curiosity, relief, belonging, stimulation, pain management or a way to survive difficult feelings. Understanding the reason does not excuse harm; it helps us work out what support and replacement coping strategies are actually needed.

Boredom can be intensely uncomfortable

Removing an addictive behaviour can expose long periods of emptiness, isolation or lack of structure. Sustainable recovery is easier when something meaningful replaces what has been lost: connection, activity, creativity, movement, volunteering, learning, rest or purposeful routine.

Addictive behaviours provide anticipation and reward

The brain learns from what works quickly. If a behaviour repeatedly reduces distress or produces relief, stimulation or reward, it becomes easier to reach for it automatically the next time the same feeling appears. Understanding this learning loop helps replace shame with practical intervention.

Isolation can make the pattern stronger

People usually have more than one reason for turning towards a substance or addictive behaviour. Curiosity about what happens before, during and after use can reveal the need underneath it and point towards safer ways of meeting that need.

Removing the behaviour leaves a gap

Removing an addictive behaviour can expose long periods of emptiness, isolation or lack of structure. Sustainable recovery is easier when something meaningful replaces what has been lost: connection, activity, creativity, movement, volunteering, learning, rest or purposeful routine.

Recovery needs meaningful replacement, not only restriction

People usually have more than one reason for turning towards a substance or addictive behaviour. Curiosity about what happens before, during and after use can reveal the need underneath it and point towards safer ways of meeting that need.

Questions to explore

What happens immediately before the urge? What changes while using? What feeling, sensation or situation becomes easier? What happens afterwards? What might offer even 10% of the same relief with less harm?

Moving from the reason to a safer response

Understanding the 'why' is most useful when it leads to a practical alternative.

If I need relief	Grounding, breathing, changing environment, talking to someone, professional support.
If I need stimulation	Movement, music, novelty, structured activity, ADHD-informed strategies where appropriate.
If I need connection	A trusted person, peer group, recovery community, shared activity or therapy.
If I need sleep	Sleep assessment, routine, reducing triggers and discussing persistent insomnia with a clinician.
If I need escape	Create immediate safety and stabilisation; trauma work should be paced and professionally supported.
If I am physically dependent	Seek medical or specialist substance-use advice; do not assume abrupt stopping is safe.

Books & places for support

Changing for Good

James O. Prochaska, John C. Norcross & Carlo C. DiClemente

Helps explain stages of change and why motivation can fluctuate.

The Unexpected Joy of Being Sober

Catherine Gray

An accessible lived-experience account focused on alcohol and building a life beyond drinking.

The Biology of Desire

Marc Lewis

Explores addiction through learning, motivation and brain change and offers a useful alternative to moral explanations.

UK help

NHS / GP	A GP can discuss alcohol or drug concerns and local treatment. You can ask for help before things reach crisis point.
FRANK	Confidential drug information and local-service finder. Helpline: 0300 123 6600.
Adfam	Support for families affected by another person's drinking, drug use or gambling.
SMART Recovery	Practical meetings and tools supporting motivation, coping with urges, thoughts/behaviours and balanced living. Family & Friends support is also available.
Local drug & alcohol services	Community services can offer assessment, harm-reduction advice, psychological support, recovery groups and treatment pathways.
Emergency help	Call 999 for suspected overdose, collapse, seizure, severe breathing difficulty or immediate danger. Physical dependence on alcohol or benzodiazepines can make abrupt withdrawal dangerous.

Educational resource - not a substitute for individual medical, psychological or emergency care.