

# Rejection Sensitivity

## Moments that can intensify the experience — understanding the load underneath

Rejection sensitivity is a descriptive term, not a separate diagnosis. Some people with ADHD describe criticism, ambiguity or perceived rejection as especially intense and difficult to shake.

**The key idea:** Slow the meaning-making down. Separate what is known from what is feared, ask for clarification where appropriate, and give the nervous system time to settle before responding.

## The six moments in this resource

### 1. A brief or delayed reply

When this happens, the brain may have to hold, shift or regulate more than is visible from the outside.

### 2. Unexpected criticism

This can add uncertainty or interrupt the mental thread that was already being managed.

### 3. A change in someone's tone

The effect is often cumulative: what looks small may arrive on top of an already busy nervous system.

### 4. Not being invited

Without an external support or clear next step, the demand can become harder to organise or tolerate.

### 5. Making a visible mistake

Stress, fatigue, sensory load and previous experiences can reduce available capacity further.

### 6. Unclear feedback

Recovery matters. Repeated demands without enough recovery can make the same situation harder later in the day.

#### NOTICE THE LOAD

Look for the build-up, not only the final trigger.

#### REDUCE DEMANDS

Simplify, externalise, clarify or postpone what can safely wait.

#### PAUSE

Create enough space to think before adding another demand.

#### RECOVER

Give the brain and body time to settle before expecting full capacity.

## Practical ways to lower the load

The aim is not to force the difficulty away. It is to change the environment, timing or amount of information so that the task asks less of the person's executive and nervous-system capacity.

### **Make the invisible visible**

Use written steps, timers, visual cues, labels, checklists or a single capture place rather than relying on memory alone.

### **Reduce simultaneous demands**

One question, one instruction or one decision at a time can be easier to process than several competing demands.

### **Build transition space**

Warnings, finishing points and a short buffer between activities reduce the cost of switching attention.

### **Protect recovery**

Quiet time, food, drink, movement, sensory reduction and fewer social demands may restore capacity.

### **Use collaborative language**

Ask what would make the next step easier. Support works better when it reduces shame rather than adding pressure.

### **Review patterns, not character**

Look for repeatable conditions: time of day, noise, fatigue, ambiguity, interruptions, hunger, urgency or social demand.

## For partners, family and supporters

Try not to interpret difficulty as laziness, rudeness or lack of care. Reduce repeated questioning when the person is overloaded, offer information in a usable form, and agree in advance how to signal that a pause is needed. Support can still include boundaries; the difference is that boundaries are clear and calm rather than shaming.

## At work or in education

Helpful adjustments may include written instructions, quieter working areas, reduced distraction, predictable deadlines, task chunking, advance warning of changes, extra processing time and agreed ways to protect focused work. Adjustments should be individual rather than assumed.

# My Rejection Sensitivity Map

Use this page after an experience, or when you are calm enough to notice patterns. Short answers are enough.

**What happened?**

|  |
|--|
|  |
|  |
|  |

**What did my mind immediately decide it meant?**

|  |
|--|
|  |
|  |
|  |

**What facts do I actually have?**

|  |
|--|
|  |
|  |
|  |

**What else could be true?**

|  |
|--|
|  |
|  |
|  |

**What response can wait until I am steadier?**

|  |
|--|
|  |
|  |
|  |

## A message from Maggie

“When something that looks small becomes the last straw, I want you to look underneath it before you judge yourself. Ask what your brain and body were already carrying. The goal is not to become better at pushing through. It is to notice your load earlier, make the next step more workable and give yourself enough recovery to begin again.”

# Understanding, support and further reading

This resource is psychoeducation, not a diagnostic checklist. ADHD presentations vary widely, and the terms people use for lived experiences do not always map neatly onto formal diagnostic language. Difficulties can also overlap with autism, anxiety, depression, sleep problems, trauma, pain, medication effects and other health conditions.

## Useful professional context

**NICE guideline NG87 — Attention deficit hyperactivity disorder: diagnosis and management.** NICE recommends a comprehensive approach to ADHD and recognises the importance of environmental modifications and support across everyday settings.

**NHS — ADHD in adults.** NHS information describes common difficulties with attention, organisation, forgetfulness and impulsivity and points to support and reasonable adjustments.

## Book recommendations

- **Taking Charge of Adult ADHD** — Russell A. Barkley
- **How to ADHD** — Jessica McCabe
- **Smart but Stuck** — Thomas E. Brown

## When to seek more support

If these difficulties are causing significant distress, affecting safety, work, education, relationships or daily care, consider speaking with a GP or appropriately qualified clinician. If anger, overwhelm or hopelessness creates an immediate risk of harm, seek urgent support rather than relying on a self-help resource.

## Sources

National Institute for Health and Care Excellence (NICE), NG87: Attention deficit hyperactivity disorder: diagnosis and management.  
NHS: ADHD in adults.

These sources provide professional context; the practical reflection exercises in this guide are original psychoeducational material from Space to Breathe Therapy.